• Solution A consists of a 0.050 M aqueous solution of benzoic acid,  $C_6H_5COOH$ , at 25 °C. Calculate the pH of Solution A. The p $K_a$  of benzoic acid is 4.20.

As benzoic acid is a weak acid,  $[H_3O^+]$  must be calculated using a reaction table:

	C <sub>6</sub> H <sub>5</sub> COOH	<del>~`</del>	$\mathbf{H}^{+}$	C <sub>6</sub> H <sub>5</sub> COO <sup>-</sup>
initial	0.050		0	0
change	- <i>x</i>		+x	+x
final	0.050 <i>-x</i>		x	x

The equilibrium constant  $K_a$  is given by:

$$K_{\rm a} = \frac{[{\rm H}^+][{\rm C_6H_5C00^-}]}{[{\rm C_6H_5C00H}]} = \frac{x^2}{0.050 - x}$$

As  $pK_a = -\log_{10}K_a$ ,  $K_a = 10^{-4.20}$  and is very small,  $0.050 - x \sim 0.050$  and hence:

$$x^2 = 0.050 \times 10^{-4.2}$$
 or  $x = 1.78 \times 10^{-3} \text{ M} = [\text{H}^+]$ 

Hence, the pH is given by:

$$pH = -log_{10}[H^+] = -log_{10}(1.78 \times 10^{-3}) = 2.75$$

pH = **2.75** 

Other than water, what are the major species present in solution A?

 $K_{\rm a}$  is very small and the equilibrium lies almost completely to the left. The major species present are water and the undissociated acid: C<sub>6</sub>H<sub>5</sub>COOH

Solution B consists of a 0.050 M aqueous solution of ammonia, NH<sub>3</sub>, at 25 °C. Calculate the pH of Solution B. The  $pK_a$  of NH<sub>4</sub><sup>+</sup> is 9.24.

## NH<sub>3</sub> is a weak base so [OH<sup>-</sup>] must be calculated by considering the equilibrium:

	NH <sub>3</sub>	H <sub>2</sub> O	 NH4 <sup>+</sup>	OH-
initial	0.050	large	0	0
change	- <i>y</i>	negligible	+ <i>y</i>	+ <i>y</i>
final	0.050 - y	large	у	у

The equilibrium constant  $K_b$  is given by:

## ANSWER CONTINUES ON THE NEXT PAGE

Marks 6



Other than water, what are the major species present in solution B?

 $K_{\rm b}$  is very small and the equilibrium lies almost completely to the left. The major species present are water and the unprotonated weak base: NH<sub>3</sub>

## THIS QUESTION CONTINUES ON THE NEXT PAGE.