

The University of Sydney

COMBINATION CHEM1001/1101 EXAM

FIRST SEMESTER EXAMINATION

CONFIDENTIAL

JUN 2003 / NOV 2003

TIME ALLOWED: THREE HOURS

GIVE THE FOLLOWING INFORMATION IN BLOCK LETTERS

FAMILY NAME		SID NUMBER	
OTHER NAMES		TABLE NUMBER	

INSTRUCTIONS TO CANDIDATES

- All questions are to be attempted. There are 18 pages of examinable material.
- Complete the written section of the examination paper in **INK**.
- Read each question carefully. Report the appropriate answer and show all relevant working in the space provided.
- The total score for this paper is 100. The possible score per page is shown in the adjacent tables.
- Each new short answer question begins with a •.
- Electronic calculators, including programmable calculators, may be used. Students are warned, however, that credit may not be given, even for a correct answer, where there is insufficient evidence of the working required to obtain the solution.
- Numerical values required for any question and a Periodic Table may be found on a separate data sheet.
- Pages 5, 12, 15, 17 and 24 are for rough working only.

OFFICIAL USE ONLY

Multiple choice section

	Marks	
Pages	Max	Gained
2-11	37	

Short answer section

Page	Marks		Marker
	Max	Gained	
13	6		
14	8		
16	8		
18	7		
19	7		
20	8		
21	6		
22	5		
23	8		
Total	63		
Check Total			

Marks
2

- The relative atomic mass of magnesium is reported as 24.3. Show how this figure is calculated given the natural abundances of the following isotopes of magnesium: ^{24}Mg (79.0 %); ^{25}Mg (10.0 %); ^{26}Mg (11.0 %).

- With examples, briefly explain what allotropes are.

2

- Complete the following table.

2

Formula	Name
Na_2CO_3	
	iron(III) oxide
PCl_3	
	ammonia

Marks
4

- Explain the physical characteristics of an ionic solid, such as brittleness, high melting point and conductivity when molten, in terms of the bonding present.

- Draw Lewis diagrams of the following species. Show both bonding and non-bonding pairs of valence electrons.

4

NH_4^+	CO_2
-----------------	---------------

Marks
5

- The active ingredient in superphosphate fertilizer is calcium dihydrogenphosphate, $\text{Ca}(\text{H}_2\text{PO}_4)_2$. It is made by treating insoluble rock phosphate, $\text{Ca}_3(\text{PO}_4)_2$ with sulfuric acid. The other product of the reaction is calcium sulfate. Write the molecular equation for the reaction.

What mass of sulfuric acid is needed to convert 1.0 tonne (1000 kg) of rock phosphate to superphosphate?

Answer:

- Analysis of an unknown compound returned the following percentage composition by weight:

nitrogen: 26.2%; chlorine: 66.4% hydrogen 7.5%


What is the empirical formula of this compound?

Answer:

3

Marks
5

- The structure of morphine is given below.



Name the functional groups in morphine that have been highlighted by the boxes.

a =

b =

c =

What are the approximate bond angles at the labelled atoms?

Atom	Bond angles
#1 N	
#2 C	
#3 C	
#4 O	

THE REMAINDER OF THIS PAGE IS FOR ROUGH WORKING ONLY

Marks
3

- In the chlor-alkali process, three useful products are formed, including two of the “top ten” chemicals. Write the overall reaction, identify the two “top ten” chemicals, and propose why the third useful product is not usually harnessed in this process.

Explain why the $\text{Na}^+(\text{aq})$ is not reduced to $\text{Na}(\text{s})$ in this process.

2


- How does nitric oxide, $\text{NO}(\text{g})$, form in a car engine? What happens to the NO once emitted from the tailpipe? Make sure you include the appropriate chemical reactions in your answer.

Marks
4

- A 50.0 mL solution contained 10.00 g of NaOH in water at 25.00 °C. When it was added to a 250.0 mL solution of 0.200 M HCl at 25.00 °C in a “coffee cup” calorimeter, the temperature of the solution rose to 33.95 °C. Assuming the specific heat of the solution is $4.18 \text{ J K}^{-1} \text{ g}^{-1}$, that the calorimeter absorbs a negligible amount of heat, and that the density of the solution is 1.00 g mL^{-1} , calculate ΔH_r (in kJ mol^{-1}) for the following reaction. $\text{H}^+(\text{aq}) + \text{OH}^-(\text{aq}) \rightarrow \text{H}_2\text{O}(\text{l})$

When the experiment was repeated using 12.00 g of NaOH in water, the temperature increase was the same. Explain.

- You are a member of a research team of industrial chemists who are discussing the operation of an ammonia plant. Ammonia is formed from nitrogen and hydrogen according to the following equilibrium reaction.




The plant operates close to 700 K, at which K_p is $1.00 \times 10^{-4} \text{ atm}^{-2}$ and employs the stoichiometric ratio 1:3 of $\text{N}_2:\text{H}_2$. At equilibrium the partial pressure of NH_3 is 50 atm. Calculate the partial pressures of each reactant and hence the total pressure under these conditions.

Marks
3

--	--	--

p(N ₂) =	p(H ₂) =	p(total) =
----------------------	----------------------	------------

- Ammonium carbamate ($\text{NH}_2\text{COONH}_4$) is a salt of carbamic acid that is found in the blood and urine of mammals. At 250 °C, $K_c = 1.58 \times 10^{-8} \text{ M}^3$ for the following equilibrium:



If 7.81 g of $\text{NH}_2\text{COONH}_4$ is introduced into a 0.500 L evacuated container, what is the total pressure inside the container at equilibrium at 250 °C?

3

--	--

Answer:

Marks
2

- Diborane (B_2H_6) is a highly reactive compound, which was once considered as a possible rocket fuel for the US space program. Calculate the heat of formation of diborane at 298 K from the following reactions.

Reaction	ΔH_r (kJ mol ⁻¹)
$2B(s) + \frac{3}{2}O_2(g) \rightarrow B_2O_3(s)$	-1273
$B_2H_6(g) + 3O_2(g) \rightarrow B_2O_3(s) + 3H_2O(g)$	-2035
$H_2(g) + \frac{1}{2}O_2(g) \rightarrow H_2O(l)$	-286
$H_2O(l) \rightarrow H_2O(g)$	+44

Answer:


- What is meant by “cathodic protection”? Which of the following metals can provide cathodic protection to iron and why?

Zn, Ni, Al, Sn

2

Marks
6

- In the refining of copper, impure copper electrodes are electrolysed in a manner such as described in the following figure. Indicate in the boxes on the figure, which electrode is the anode and which is the cathode.



Why are noble metals left as a mud on the bottom of the reaction cell?


Explain why Zn^{2+} and Fe^{2+} are not deposited from solution during this reaction.

How many kilograms of pure copper will be obtained when the electrolytic cell is operated for 24.0 hours at a constant current of 100.0 A?

Answer:

Marks
3

- Corn is a valuable source of industrial chemicals. For example, furfural is prepared from corncobs. It is an important reactant in plastics manufacture and a key solvent in the production of cellulose acetate, which is used to make products such as videotape and waterproof fabric. Furfural can be reduced to furfuryl alcohol or oxidised to 2-furoic acid. The structures of these three compounds are shown below.




Explain, in terms of oxidation numbers, why we say that furfural is *oxidised* to 2-furoic acid and *reduced* to furfuryl alcohol.

Which of these three compounds can form hydrogen bonds? Draw the structure in each case.

THE REMAINDER OF THIS PAGE IS FOR ROUGH WORKING ONLY

Marks
3

- The figure below shows the boiling points of Group 14 and 17 hydrides as a function of the period (row) of the periodic table.




A number of trends are apparent from this figure, including:

- the tetrahydrides have lower boiling points than the monohydrides,
- the boiling point increases with period, with the exception of HF.

Explain these two trends, and the reason that HF is exceptional.

Marks
5

- Styrene is the monomer from which the important polymer, polystyrene, is manufactured. The formula of styrene is shown below.



Draw the repeating unit for polystyrene.

The average C–C bond length in the backbone of polystyrene is 0.154 nm and the C–C–C bond angle is 109.5°. Calculate the total extended end-to-end distance of the polymer chain, and the average radius of gyration in a sample of polystyrene that has a molar mass of 100,000 g mol⁻¹.

Unlike polystyrene, which exhibits free rotation about the C-C single bonds, a polypeptide exhibits restricted rotation in its backbone because of the partial double bond character of the peptide bond. Explain this feature of polypeptides using resonance structures of the peptide bond.

The University of Sydney

FIRST SEMESTER EXAMINATION**JUNE 2003**

Numerical Data

Physical constants

$$\text{Planck constant} = h = 6.626 \times 10^{-34} \text{ J s}$$

$$\text{Speed of light in vacuum} = c_0 = 2.998 \times 10^8 \text{ m s}^{-1}$$

$$\text{Avogadro constant} = N_A = 6.022 \times 10^{23} \text{ mol}^{-1}$$

$$\text{Faraday constant} = F = 96485 \text{ C mol}^{-1}$$

$$\begin{aligned} \text{Ideal gas constant} &= R = 8.314 \text{ J K}^{-1} \text{ mol}^{-1} \\ &= 0.08206 \text{ L atm K}^{-1} \text{ mol}^{-1} \end{aligned}$$

$$\text{Volume of 1 mol of ideal gas at 1 atm, } 0 \text{ }^\circ\text{C} = 22.4 \text{ L}$$

$$\text{Volume of 1 mol of ideal gas at 1 atm, } 25 \text{ }^\circ\text{C} = 24.5 \text{ L}$$

Conversion factors

$$0 \text{ }^\circ\text{C} = 273 \text{ K}$$

$$1 \text{ atm} = 101.3 \text{ kPa} = 760.0 \text{ mmHg}$$

$$1 \text{ nm} = 10^{-9} \text{ m}$$

$$1 \text{ MHz} = 10^6 \text{ Hz} = 10^6 \text{ s}^{-1}$$

$$1 \text{ L} = 10^{-3} \text{ m}^3$$

Formulas

$$E = h\nu$$

$$c = \lambda\nu$$

$$PV = nRT$$

$$\Delta H = C_p m \Delta T$$

$$\text{pH} = -\log[\text{H}^+]$$

$$\text{pH} + \text{pOH} = 14.00$$

$$K_p = K_c(RT)^{\Delta n}$$

**A periodic table is printed on the other side of this data sheet.
Atomic weights are included in the periodic table.**

PERIODIC TABLE OF THE ELEMENTS

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
1 HYDROGEN H 1.008																	2 HELIUM He 4.003
3 LITHIUM Li 6.941	4 BERYLLIUM Be 9.012											5 BORON B 10.81	6 CARBON C 12.01	7 NITROGEN N 14.01	8 OXYGEN O 16.00	9 FLUORINE F 19.00	10 NEON Ne 20.18
11 SODIUM Na 22.99	12 MAGNESIUM Mg 24.31											13 ALUMINIUM Al 26.98	14 SILICON Si 28.09	15 PHOSPHORUS P 30.97	16 SULFUR S 32.07	17 CHLORINE Cl 35.45	18 ARGON Ar 39.95
19 POTASSIUM K 39.10	20 CALCIUM Ca 40.08	21 SCANDIUM Sc 44.96	22 TITANIUM Ti 47.88	23 VANADIUM V 50.94	24 CHROMIUM Cr 52.00	25 MANGANESE Mn 54.94	26 IRON Fe 55.85	27 COBALT Co 58.93	28 NICKEL Ni 58.69	29 COPPER Cu 63.55	30 ZINC Zn 65.39	31 GALLIUM Ga 69.72	32 GERMANIUM Ge 72.59	33 ARSENIC As 74.92	34 SELENIUM Se 78.96	35 BROMINE Br 79.90	36 KRYPTON Kr 83.80
37 RUBIDIUM Rb 85.47	38 STRONTIUM Sr 87.62	39 YTRIUM Y 88.91	40 ZIRCONIUM Zr 91.22	41 NIوبيUM Nb 92.91	42 MOLYBDENUM Mo 95.94	43 TECHNETIUM Tc [98.91]	44 RUTHENIUM Ru 101.07	45 RHODIUM Rh 102.91	46 PALLADIUM Pd 106.4	47 SILVER Ag 107.87	48 CADMIUM Cd 112.40	49 INDIUM In 114.82	50 TIN Sn 118.69	51 ANTIMONY Sb 121.75	52 TELLURIUM Te 127.60	53 IODINE I 126.90	54 XENON Xe 131.30
55 CAESIUM Cs 132.91	56 BARIUM Ba 137.34	57-71	72 HAFNIUM Hf 178.49	73 TANTALUM Ta 180.95	74 TUNGSTEN W 183.85	75 RHENIUM Re 186.2	76 OSMIUM Os 190.2	77 IRIDIUM Ir 192.22	78 PLATINUM Pt 195.09	79 GOLD Au 196.97	80 MERCURY Hg 200.59	81 THALLIUM Tl 204.37	82 LEAD Pb 207.2	83 BISMUTH Bi 208.98	84 POLONIUM Po [210.0]	85 ASTATINE At [210.0]	86 RADON Rn [222.0]
87 FRANCIUM Fr [223.0]	88 RADIUM Ra [226.0]	89-103	104 RUTHERFORDIUM Rf [261]	105 DUBNIUM Db [262]	106 SEABORGIUM Sg [266]	107 BOHRIUM Bh [262]	108 HASSIUM Hs [265]	109 MEITNERIUM Mt [266]									

LANTHANIDES	57 LANTHANUM La 138.91	58 CERIUM Ce 140.12	59 PRASEODYMIUM Pr 140.91	60 NEODYMIUM Nd 144.24	61 PROMETHIUM Pm [144.9]	62 SAMARIUM Sm 150.4	63 EUROPIUM Eu 151.96	64 GADOLINIUM Gd 157.25	65 TERBIUM Tb 158.93	66 DYSPROSIUM Dy 162.50	67 HOLMIUM Ho 164.93	68 ERBIUM Er 167.26	69 THULIUM Tm 168.93	70 YTTERBIUM Yb 173.04	71 LUTETIUM Lu 174.97
ACTINIDES	89 ACTINIUM Ac [227.0]	90 THORIUM Th 232.04	91 PROTACTINIUM Pa [231.0]	92 URANIUM U 238.03	93 NEPTUNIUM Np [237.0]	94 PLUTONIUM Pu [239.1]	95 AMERICIUM Am [243.1]	96 CURIUM Cm [247.1]	97 BERKELIUM Bk [247.1]	98 CALIFORNIUM Cf [252.1]	99 EINSTEINIUM Es [252.1]	100 FERMIUM Fm [257.1]	101 MENDELEVIUM Md [256.1]	102 NOBELIUM No [259.1]	103 LAWRENCIUM Lr [260.1]