

Topics in the November 2006 Exam Paper for CHEM1002

Click on the links for resources on each topic.

2006-N-2:

2006-N-3:

- [Strong Acids and Bases](#)
- [Weak Acids and Bases](#)

2006-N-4:

- [Weak Acids and Bases](#)
- [Calculations Involving \$pK_a\$](#)

2006-N-5:

- [Coordination Chemistry](#)

2006-N-6:

- [Alkenes](#)
- [Alcohols](#)
- [Organic Halogen Compounds](#)

2006-N-7:

- [Representations of Molecular Structure](#)
- [Stereochemistry](#)

2006-N-8:

- [Alcohols](#)
- [Amines](#)
- [Aldehydes and Ketones](#)
- [Carboxylic Acids and Derivatives](#)

2006-N-9:

- [Stereochemistry](#)

FUNDAMENTALS OF CHEMISTRY 1B - CHEM1002

SECOND SEMESTER EXAMINATION

CONFIDENTIAL

NOVEMBER 2006

TIME ALLOWED: THREE HOURS

GIVE THE FOLLOWING INFORMATION IN BLOCK LETTERS

FAMILY NAME		SID NUMBER	
OTHER NAMES		TABLE NUMBER	

INSTRUCTIONS TO CANDIDATES

- All questions are to be attempted. There are 19 pages of examinable material.
- Complete the written section of the examination paper in **INK**.
- Read each question carefully. Report the appropriate answer and show all relevant working in the space provided.
- The total score for this paper is 100. The possible score per page is shown in the adjacent tables.
- Each new question of the short answer section begins with a •.
- Electronic calculators, including programmable calculators, may be used. Students are warned, however, that credit may not be given, even for a correct answer, where there is insufficient evidence of the working required to obtain the solution.
- Numerical values required for any question, standard electrode reduction potentials, a Periodic Table and some useful formulas may be found on the separate data sheet.
- Pages 16, 18, 21 and 24 are for rough working only.

OFFICIAL USE ONLY

~~Multiple choice section~~

		Marks	
Pages	Max	Gained	
2-12	50		

Short answer section

Page	Marks		Marker
	Max	Gained	
13	4		
14	9		
15	6		
17	6		
19	8		
20	5		
22	6		
23	6		
Total	50		


- Limestone caves can be found near Sydney. How have these caves been formed? Use appropriate chemical equations in your explanation.

Marks
4

Stalactites and stalagmites can be found in many limestone caves. How do these form? Use appropriate chemical equations in your explanation.

- The titration curves for a titration of a weak acid with a strong base and for a strong acid with a strong base are distinctly different. Draw a diagram for each case.

Marks
7



List the main differences.

Blank space for listing the main differences between the titration curves.

Explain these differences.

Blank space for explaining the differences between the titration curves.

- What is the difference between the 'end point' and the 'equivalence point' in a titration.

2

Blank space for answering the question about the difference between the end point and the equivalence point.

Marks
2

- What is the pH of a 0.020 M solution of HF? The pK_a of HF is 3.17.

pH =

2

- What is the pH of a solution that is 0.075 M in acetic acid and 0.150 M in sodium acetate? The pK_a of CH_3COOH is 4.76.

pH =

2

- What is the pH of a 0.010 M solution of $Ba(OH)_2$?

pH =

- Complete the following table.






Marks
6

Formula	Oxidation state of transition metal	Coordination number of transition metal	Number of <i>d</i> -electrons in transition metal	Species formed upon dissolving in water
$\text{Na}_2[\text{Ni}(\text{CN})_4]$				
$[\text{Cr}(\text{NH}_3)_5\text{Cl}]\text{Cl}_2$				
$[\text{Cu}(\text{en})_3]\text{Br}_2$				

en = ethylenediamine = $\text{NH}_2\text{CH}_2\text{CH}_2\text{NH}_2$


THE REMAINDER OF THIS PAGE IS FOR ROUGH WORKING ONLY.

- Give the name of the starting material where indicated and the constitutional formula of the major organic product formed in each of the following reactions.

Marks
8**Name:****Name:****Name:****Name:**

- Classify the starting materials for the following reaction as nucleophile or electrophile in the boxes provided and draw the structure of the product.

Marks
3



--	--

- Draw the constitutional formula for each of the following compounds.

2

(E)-5-methylhex-2-ene


cis-1,2-dichlorocyclopentane

THE REMAINDER OF THIS PAGE IS FOR ROUGH WORKING ONLY.

- Consider the following reaction sequence.

Marks

6




Give the reagents **A** and **D** and draw the structures of the major organic products, **B**, **C**, **E** and **F**, formed in these reactions.

A	D
B	E
C	F

- Phenylalanine is a naturally occurring amino acid. Only the enantiomer (**X**) is commonly produced in nature.

Marks
6



What is the molecular formula of (**X**)?

List the substituents attached to the stereogenic centre in descending order of priority according to the sequence rules.

highest priority

lowest priority

--	--	--	--

What is the absolute stereochemistry of (**X**)? Write (*R*) or (*S*).

Name the functional groups, highlighted by the boxes **a** and **b**, present in (**X**).

a =	b =
------------	------------

THE REMAINDER OF THIS PAGE IS FOR ROUGH WORKING ONLY.

CHEM1002 - CHEMISTRY 1B**DATA SHEET***Physical constants*Avogadro constant, $N_A = 6.022 \times 10^{23} \text{ mol}^{-1}$ Faraday constant, $F = 96485 \text{ C mol}^{-1}$ Planck constant, $h = 6.626 \times 10^{-34} \text{ J s}$ Speed of light in vacuum, $c = 2.998 \times 10^8 \text{ m s}^{-1}$ Rydberg constant, $E_R = 2.18 \times 10^{-18} \text{ J}$ Boltzmann constant, $k_B = 1.381 \times 10^{-23} \text{ J K}^{-1}$ Gas constant, $R = 8.314 \text{ J K}^{-1} \text{ mol}^{-1}$
 $= 0.08206 \text{ L atm K}^{-1} \text{ mol}^{-1}$ Charge of electron, $e = 1.602 \times 10^{-19} \text{ C}$ Mass of electron, $m_e = 9.1094 \times 10^{-31} \text{ kg}$ Mass of proton, $m_p = 1.6726 \times 10^{-27} \text{ kg}$ Mass of neutron, $m_n = 1.6749 \times 10^{-27} \text{ kg}$ *Properties of matter*

Volume of 1 mole of ideal gas at 1 atm and 25 °C = 24.5 L

Volume of 1 mole of ideal gas at 1 atm and 0 °C = 22.4 L

Density of water at 298 K = 0.997 g cm^{-3} *Conversion factors*

1 atm = 760 mmHg = 101.3 kPa

0 °C = 273 K

1 L = 10^{-3} m^3 1 Å = 10^{-10} m 1 eV = $1.602 \times 10^{-19} \text{ J}$ 1 Ci = $3.70 \times 10^{10} \text{ Bq}$ 1 Hz = 1 s^{-1} *Decimal fractions*

Fraction	Prefix	Symbol
10^{-3}	milli	m
10^{-6}	micro	μ
10^{-9}	nano	n
10^{-12}	pico	p

Decimal multiples

Multiple	Prefix	Symbol
10^3	kilo	k
10^6	mega	M
10^9	giga	G

CHEM1002 - CHEMISTRY 1B**Standard Reduction Potentials, E°**

Reaction	E° / V
$\text{Co}^{3+}(\text{aq}) + \text{e}^- \rightarrow \text{Co}^{2+}(\text{aq})$	+1.82
$\text{Ce}^{4+}(\text{aq}) + \text{e}^- \rightarrow \text{Ce}^{3+}(\text{aq})$	+1.72
$\text{Au}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Au}(\text{s})$	+1.50
$\text{Cl}_2 + 2\text{e}^- \rightarrow 2\text{Cl}^-(\text{aq})$	+1.36
$\text{O}_2 + 4\text{H}^+(\text{aq}) + 4\text{e}^- \rightarrow 2\text{H}_2\text{O}$	+1.23
$\text{Br}_2 + 2\text{e}^- \rightarrow 2\text{Br}^-(\text{aq})$	+1.10
$\text{MnO}_2(\text{s}) + 4\text{H}^+(\text{aq}) + \text{e}^- \rightarrow \text{Mn}^{3+} + 2\text{H}_2\text{O}$	+0.96
$\text{Pd}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Pd}(\text{s})$	+0.92
$\text{Ag}^+(\text{aq}) + \text{e}^- \rightarrow \text{Ag}(\text{s})$	+0.80
$\text{Fe}^{3+}(\text{aq}) + \text{e}^- \rightarrow \text{Fe}^{2+}(\text{aq})$	+0.77
$\text{Cu}^+(\text{aq}) + \text{e}^- \rightarrow \text{Cu}(\text{s})$	+0.53
$\text{Cu}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Cu}(\text{s})$	+0.34
$\text{Sn}^{4+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Sn}^{2+}(\text{aq})$	+0.15
$2\text{H}^+(\text{aq}) + 2\text{e}^- \rightarrow \text{H}_2(\text{g})$	0 (by definition)
$\text{Fe}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Fe}(\text{s})$	-0.04
$\text{Pb}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Pb}(\text{s})$	-0.13
$\text{Sn}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Sn}(\text{s})$	-0.14
$\text{Ni}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Ni}(\text{s})$	-0.24
$\text{Co}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Co}(\text{s})$	-0.28
$\text{Fe}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Fe}(\text{s})$	-0.44
$\text{Cr}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Cr}(\text{s})$	-0.74
$\text{Zn}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Zn}(\text{s})$	-0.76
$2\text{H}_2\text{O} + 2\text{e}^- \rightarrow \text{H}_2(\text{g}) + 2\text{OH}^-(\text{aq})$	-0.83
$\text{Cr}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Cr}(\text{s})$	-0.89
$\text{Al}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Al}(\text{s})$	-1.68
$\text{Mg}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Mg}(\text{s})$	-2.36
$\text{Na}^+(\text{aq}) + \text{e}^- \rightarrow \text{Na}(\text{s})$	-2.71
$\text{Ca}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Ca}(\text{s})$	-2.87
$\text{Li}^+(\text{aq}) + \text{e}^- \rightarrow \text{Li}(\text{s})$	-3.04

CHEM1002 - CHEMISTRY 1B

Useful formulas

<p>Quantum Chemistry</p> $E = h\nu = hc/\lambda$ $\lambda = h/mv$ $4.5k_B T = hc/\lambda$ $E = Z^2 E_R (1/n^2)$ $\Delta x \cdot \Delta(mv) \geq h/4\pi$ $q = 4\pi r^2 \times 5.67 \times 10^{-8} \times T^4$	<p>Electrochemistry</p> $\Delta G^\circ = -nFE^\circ$ $\text{Moles of } e^- = It/F$ $E = E^\circ - (RT/nF) \times 2.303 \log Q$ $= E^\circ - (RT/nF) \times \ln Q$ $E^\circ = (RT/nF) \times 2.303 \log K$ $= (RT/nF) \times \ln K$ $E = E^\circ - \frac{0.0592}{n} \log Q \text{ (at } 25^\circ \text{C)}$
<p>Acids and Bases</p> $pK_w = \text{pH} + \text{pOH} = 14.00$ $pK_w = \text{p}K_a + \text{p}K_b = 14.00$ $\text{pH} = \text{p}K_a + \log\{[A^-] / [HA]\}$	<p>Gas Laws</p> $PV = nRT$ $(P + n^2 a/V^2)(V - nb) = nRT$
<p>Colligative properties</p> $\pi = cRT$ $P_{\text{solution}} = X_{\text{solvent}} \times P^\circ_{\text{solvent}}$ $p = kc$ $\Delta T_f = K_f m$ $\Delta T_b = K_b m$	<p>Kinetics</p> $t_{1/2} = \ln 2/k$ $k = Ae^{-E_a/RT}$ $\ln[A] = \ln[A]_0 - kt$ $\ln \frac{k_2}{k_1} = \frac{E_a}{R} \left(\frac{1}{T_1} - \frac{1}{T_2} \right)$
<p>Radioactivity</p> $t_{1/2} = \ln 2/\lambda$ $A = \lambda N$ $\ln(N_0/N_t) = \lambda t$ $^{14}\text{C age} = 8033 \ln(A_0/A_t)$	<p>Thermodynamics & Equilibrium</p> $\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$ $\Delta G = \Delta G^\circ + RT \ln Q$ $\Delta G^\circ = -RT \ln K$ $K_p = K_c (RT)^{\Delta n}$
<p>Polymers</p> $R_g = \sqrt{\frac{nl_0^2}{6}}$	<p>Mathematics</p> $\text{If } ax^2 + bx + c = 0, \text{ then } x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ $\ln x = 2.303 \log x$

PERIODIC TABLE OF THE ELEMENTS

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
1 HYDROGEN H 1.008																	2 HELIUM He 4.003
3 LITHIUM Li 6.941	4 BERYLLIUM Be 9.012											5 BORON B 10.81	6 CARBON C 12.01	7 NITROGEN N 14.01	8 OXYGEN O 16.00	9 FLUORINE F 19.00	10 NEON Ne 20.18
11 SODIUM Na 22.99	12 MAGNESIUM Mg 24.31											13 ALUMINIUM Al 26.98	14 SILICON Si 28.09	15 PHOSPHORUS P 30.97	16 SULFUR S 32.07	17 CHLORINE Cl 35.45	18 ARGON Ar 39.95
19 POTASSIUM K 39.10	20 CALCIUM Ca 40.08	21 SCANDIUM Sc 44.96	22 TITANIUM Ti 47.88	23 VANADIUM V 50.94	24 CHROMIUM Cr 52.00	25 MANGANESE Mn 54.94	26 IRON Fe 55.85	27 COBALT Co 58.93	28 NICKEL Ni 58.69	29 COPPER Cu 63.55	30 ZINC Zn 65.39	31 GALLIUM Ga 69.72	32 GERMANIUM Ge 72.59	33 ARSENIC As 74.92	34 SELENIUM Se 78.96	35 BROMINE Br 79.90	36 KRYPTON Kr 83.80
37 RUBIDIUM Rb 85.47	38 STRONTIUM Sr 87.62	39 YTRIUM Y 88.91	40 ZIRCONIUM Zr 91.22	41 NIOBIUM Nb 92.91	42 MOLYBDENUM Mo 95.94	43 TECHNETIUM Tc [98.91]	44 RUTHENIUM Ru 101.07	45 RHODIUM Rh 102.91	46 PALLADIUM Pd 106.4	47 SILVER Ag 107.87	48 CADMIUM Cd 112.40	49 INDIUM In 114.82	50 TIN Sn 118.69	51 ANTIMONY Sb 121.75	52 TELLURIUM Te 127.60	53 IODINE I 126.90	54 XENON Xe 131.30
55 CAESIUM Cs 132.91	56 BARIUM Ba 137.34	57-71	72 HAFNIUM Hf 178.49	73 TANTALUM Ta 180.95	74 TUNGSTEN W 183.85	75 RHENIUM Re 186.2	76 OSMIUM Os 190.2	77 IRIDIUM Ir 192.22	78 PLATINUM Pt 195.09	79 GOLD Au 196.97	80 MERCURY Hg 200.59	81 THALLIUM Tl 204.37	82 LEAD Pb 207.2	83 BISMUTH Bi 208.98	84 POLONIUM Po [210.0]	85 ASTATINE At [210.0]	86 RADON Rn [222.0]
87 FRANCIUM Fr [223.0]	88 RADIUM Ra [226.0]	89-103	104 RUTHERFORDIUM Rf [261]	105 DUBNIUM Db [262]	106 SEABORGIUM Sg [266]	107 BOHRIUM Bh [262]	108 HASSIUM Hs [265]	109 MEITNERIUM Mt [266]									
LANTHANIDES																	
	57 LANTHANUM La 138.91	58 CERIUM Ce 140.12	59 PRASEODYMIUM Pr 140.91	60 NEODYMIUM Nd 144.24	61 PROMETHIUM Pm [144.9]	62 SAMARIUM Sm 150.4	63 EUROPIUM Eu 151.96	64 GADOLINIUM Gd 157.25	65 TERBIUM Tb 158.93	66 DYSPROSIUM Dy 162.50	67 HOLMIUM Ho 164.93	68 ERBIUM Er 167.26	69 THULIUM Tm 168.93	70 YTTERBIUM Yb 173.04	71 LUTETIUM Lu 174.97		
ACTINIDES																	
	89 ACTINIUM Ac [227.0]	90 THORIUM Th 232.04	91 PROTACTINIUM Pa [231.0]	92 URANIUM U 238.03	93 NEPTUNIUM Np [237.0]	94 PLUTONIUM Pu [239.1]	95 AMERICIUM Am [243.1]	96 CURIUM Cm [247.1]	97 BERKELIUM Bk [247.1]	98 CALIFORNIUM Cf [252.1]	99 EINSTEINIUM Es [252.1]	100 FERMIUM Fm [257.1]	101 MENDELEVIUM Md [256.1]	102 NOBELIUM No [259.1]	103 LAWRENCIUM Lr [260.1]		