

**Topics in the November 2007 Exam Paper for CHEM1002**

Click on the links for resources on each topic.

2007-N-2:

- [Solubility Equilibrium](#)
- [Metal Complexes](#)

2007-N-3:

- [Kinetics](#)

2007-N-4:

- [Weak Acids and Bases](#)
- [Calculations Involving  \$pK\_a\$](#)

2007-N-5:

- [Weak Acids and Bases](#)
- [Periodic Trends](#)
- [Coordination Chemistry](#)

2007-N-6:

- [Alkenes](#)
- [Alcohols](#)
- [Organic Halogen Compounds](#)

2007-N-7:

- [Representations of Molecular Structure](#)
- [Stereochemistry](#)

2007-N-8:

- [Alcohols](#)
- [Amines](#)
- [Organic Halogen Compounds](#)
- [Aldehydes and Ketones](#)
- [Carboxylic Acids and Derivatives](#)

2007-N-9:

- [Representations of Molecular Structure](#)
- [Stereochemistry](#)
- [Carboxylic Acids and Derivatives](#)

2007-N-10:

- [Synthetic Strategies](#)

## FUNDAMENTALS OF CHEMISTRY 1B - CHEM1002

### SECOND SEMESTER EXAMINATION

**CONFIDENTIAL**

**NOVEMBER 2007**

**TIME ALLOWED: THREE HOURS**

GIVE THE FOLLOWING INFORMATION IN BLOCK LETTERS

<b>FAMILY NAME</b>		<b>SID NUMBER</b>	
<b>OTHER NAMES</b>		<b>TABLE NUMBER</b>	

#### INSTRUCTIONS TO CANDIDATES

- All questions are to be attempted. There are 18 pages of examinable material.
- Complete the written section of the examination paper in **INK**.
- Read each question carefully. Report the appropriate answer and show all relevant working in the space provided.
- The total score for this paper is 100. The possible score per page is shown in the adjacent tables.
- Each new question of the short answer section begins with a •.
- Electronic calculators, including programmable calculators, may be used. Students are warned, however, that credit may not be given, even for a correct answer, where there is insufficient evidence of the working required to obtain the solution.
- Numerical values required for any question, standard electrode reduction potentials, a Periodic Table and some useful formulas may be found on the separate data sheet.
- Pages 14, 16, 19, 23 and 24 are for rough working only.

#### OFFICIAL USE ONLY

##### ~~Multiple choice section~~

		Marks	
Pages	Max	Gained	
2-10	36		

##### Short answer section

Page	Marks		Marker
	Max	Gained	
11	8		
12	4		
13	8		
15	10		
17	8		
18	7		
20	6		
21	7		
22	6		
<b>Total</b>	<b>64</b>		

**Marks**  
**3**

- Define what is meant by an “allotrope”. Give an example of a *pair* of allotropes involving (i) carbon and (ii) oxygen.

**5**

- The  $K_{sp}$  of  $\text{Fe}(\text{OH})_3$  is  $2.0 \times 10^{-39} \text{ M}^4$ . What is the solubility of  $\text{Fe}(\text{OH})_3$  in  $\text{g L}^{-1}$ ?

Answer:

What effect does lowering the pH have on the solubility of  $\text{Fe}(\text{OH})_3$ ? Explain your answer.



**Marks**  
**8**

- Solution A consists of a 0.50 M aqueous solution of HF at 25 °C. Calculate the pH of Solution A. The  $pK_a$  of HF is 3.17.

pH =

At 25 °C, 1.00 L of Solution B consists of 12.97 g of lithium fluoride, LiF, dissolved in water. Calculate the pH of Solution B.

pH =

Solution B (1.00 L) is poured into Solution A (1.00 L) and allowed to equilibrate at 25 °C. Calculate the pH of the final solution.

pH =

If you wanted to adjust the pH of the mixture of Solution A and Solution B to be exactly equal to 4.00, which component in the mixture would you need to increase in concentration?

- Briefly explain why H<sub>2</sub>S is a stronger Brønsted acid than H<sub>2</sub>O.

**Marks**  
2

- Compounds of *d*-block elements are frequently paramagnetic. Using the box notation to represent atomic orbitals, account for this property in compounds of Ni<sup>2+</sup>.

2






- Complete the following table.

6

Formula	Oxidation state of transition metal	Coordination number of transition metal	Number of <i>d</i> -electrons in the complex ion	Species formed upon dissolving in water
K <sub>3</sub> [Mn(CN) <sub>6</sub> ]				
[Ru(NH <sub>3</sub> ) <sub>5</sub> (OH <sub>2</sub> )](NO <sub>3</sub> ) <sub>2</sub>				
[Cr(en) <sub>3</sub> ]Cl <sub>3</sub>				



en = ethylenediamine = NH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>

- Give the name of the starting material where indicated and the constitutional formula of the major organic product formed in each of the following reactions.

**Marks**  
**8****Name:****Name:****Name:**

**Marks**  
**4**

- Classify the starting materials for each of the following reactions as nucleophile or electrophile in the boxes provided and indicate with  $\delta^+$  and  $\delta^-$  the polarisation of the H-Br and C-Br bonds in the starting materials.



**3**

- Draw the constitutional formula for each of the following compounds.

(Z)-4-methylhex-2-ene

*trans*-1,3-dichlorocyclohexane


(*R*)-butan-2-ol



Marks

6

- Consider the following reaction sequence.




Give the reagents **B** and **D** and draw the structures of the major organic products, **A**, **C**, **E** and **F**, formed in these reactions.

<b>A</b>	<b>D</b>
<b>B</b>	<b>E</b>
<b>C</b>	<b>F</b>

**Marks**  
**7**

- Dopa is a non-proteinogenic amino acid used to treat Parkinson's disease. Only the enantiomer (**X**) is effective in restoring nerve function. The other enantiomer is highly toxic.



What is the molecular formula of (**X**)?

List the substituents attached to the stereogenic centre in descending order of priority according to the sequence rules.

highest priority

lowest priority

--	--	--	--

What is the absolute stereochemistry of (**X**)? Write (*R*) or (*S*).


Name the functional groups, highlighted by the boxes **a** and **b**, present in (**X**).

<b>a</b> =	<b>b</b> =
------------	------------

Give the constitutional formula of the product obtained when (**X**) is treated with  $\text{NaHCO}_3$ .

**Marks**  
**6**

- Show clearly the reagents you would use to carry out the following chemical conversion. Exactly one intermediate compound and hence two steps are required. Give the constitutional formula of the intermediate compound.



How could you distinguish between the starting material, the intermediate compound and the final product using infrared spectroscopy?

**CHEM1002 - CHEMISTRY 1B****DATA SHEET***Physical constants*Avogadro constant,  $N_A = 6.022 \times 10^{23} \text{ mol}^{-1}$ Faraday constant,  $F = 96485 \text{ C mol}^{-1}$ Planck constant,  $h = 6.626 \times 10^{-34} \text{ J s}$ Speed of light in vacuum,  $c = 2.998 \times 10^8 \text{ m s}^{-1}$ Rydberg constant,  $E_R = 2.18 \times 10^{-18} \text{ J}$ Boltzmann constant,  $k_B = 1.381 \times 10^{-23} \text{ J K}^{-1}$ Gas constant,  $R = 8.314 \text{ J K}^{-1} \text{ mol}^{-1}$   
 $= 0.08206 \text{ L atm K}^{-1} \text{ mol}^{-1}$ Charge of electron,  $e = 1.602 \times 10^{-19} \text{ C}$ Mass of electron,  $m_e = 9.1094 \times 10^{-31} \text{ kg}$ Mass of proton,  $m_p = 1.6726 \times 10^{-27} \text{ kg}$ Mass of neutron,  $m_n = 1.6749 \times 10^{-27} \text{ kg}$ *Properties of matter*

Volume of 1 mole of ideal gas at 1 atm and 25 °C = 24.5 L

Volume of 1 mole of ideal gas at 1 atm and 0 °C = 22.4 L

Density of water at 298 K =  $0.997 \text{ g cm}^{-3}$ *Conversion factors*

1 atm = 760 mmHg = 101.3 kPa

0 °C = 273 K

1 L =  $10^{-3} \text{ m}^3$ 1 Å =  $10^{-10} \text{ m}$ 1 eV =  $1.602 \times 10^{-19} \text{ J}$ 1 Ci =  $3.70 \times 10^{10} \text{ Bq}$ 1 Hz =  $1 \text{ s}^{-1}$ *Decimal fractions*

Fraction	Prefix	Symbol
$10^{-3}$	milli	m
$10^{-6}$	micro	μ
$10^{-9}$	nano	n
$10^{-12}$	pico	p

*Decimal multiples*

Multiple	Prefix	Symbol
$10^3$	kilo	k
$10^6$	mega	M
$10^9$	giga	G

**CHEM1002 - CHEMISTRY 1B****Standard Reduction Potentials,  $E^\circ$** 

Reaction	$E^\circ / \text{V}$
$\text{S}_2\text{O}_8^{2-} + 2\text{e}^- \rightarrow 2\text{SO}_4^{2-}$	+2.01
$\text{Co}^{3+}(\text{aq}) + \text{e}^- \rightarrow \text{Co}^{2+}(\text{aq})$	+1.82
$\text{Ce}^{4+}(\text{aq}) + \text{e}^- \rightarrow \text{Ce}^{3+}(\text{aq})$	+1.72
$\text{Au}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Au}(\text{s})$	+1.50
$\text{Cl}_2 + 2\text{e}^- \rightarrow 2\text{Cl}^-(\text{aq})$	+1.36
$\text{O}_2 + 4\text{H}^+(\text{aq}) + 4\text{e}^- \rightarrow 2\text{H}_2\text{O}$	+1.23
$\text{Br}_2 + 2\text{e}^- \rightarrow 2\text{Br}^-(\text{aq})$	+1.10
$\text{MnO}_2(\text{s}) + 4\text{H}^+(\text{aq}) + \text{e}^- \rightarrow \text{Mn}^{3+} + 2\text{H}_2\text{O}$	+0.96
$\text{Pd}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Pd}(\text{s})$	+0.92
$\text{Ag}^+(\text{aq}) + \text{e}^- \rightarrow \text{Ag}(\text{s})$	+0.80
$\text{Fe}^{3+}(\text{aq}) + \text{e}^- \rightarrow \text{Fe}^{2+}(\text{aq})$	+0.77
$\text{Cu}^+(\text{aq}) + \text{e}^- \rightarrow \text{Cu}(\text{s})$	+0.53
$\text{Cu}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Cu}(\text{s})$	+0.34
$\text{Sn}^{4+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Sn}^{2+}(\text{aq})$	+0.15
$2\text{H}^+(\text{aq}) + 2\text{e}^- \rightarrow \text{H}_2(\text{g})$	0 (by definition)
$\text{Fe}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Fe}(\text{s})$	-0.04
$\text{Pb}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Pb}(\text{s})$	-0.13
$\text{Sn}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Sn}(\text{s})$	-0.14
$\text{Ni}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Ni}(\text{s})$	-0.24
$\text{Co}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Co}(\text{s})$	-0.28
$\text{Fe}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Fe}(\text{s})$	-0.44
$\text{Cr}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Cr}(\text{s})$	-0.74
$\text{Zn}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Zn}(\text{s})$	-0.76
$2\text{H}_2\text{O} + 2\text{e}^- \rightarrow \text{H}_2(\text{g}) + 2\text{OH}^-(\text{aq})$	-0.83
$\text{Cr}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Cr}(\text{s})$	-0.89
$\text{Al}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Al}(\text{s})$	-1.68
$\text{Mg}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Mg}(\text{s})$	-2.36
$\text{Na}^+(\text{aq}) + \text{e}^- \rightarrow \text{Na}(\text{s})$	-2.71
$\text{Ca}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Ca}(\text{s})$	-2.87
$\text{Li}^+(\text{aq}) + \text{e}^- \rightarrow \text{Li}(\text{s})$	-3.04

## CHEM1002 - CHEMISTRY 1B

## Useful formulas

<p><b>Quantum Chemistry</b></p> $E = h\nu = hc/\lambda$ $\lambda = h/m\nu$ $4.5k_B T = hc/\lambda$ $E = -Z^2 E_R (1/n^2)$ $\Delta x \cdot \Delta(mv) \geq h/4\pi$ $q = 4\pi r^2 \times 5.67 \times 10^{-8} \times T^4$	<p><b>Electrochemistry</b></p> $\Delta G^\circ = -nFE^\circ$ <p>Moles of <math>e^- = It/F</math></p> $E = E^\circ - (RT/nF) \times 2.303 \log Q$ $= E^\circ - (RT/nF) \times \ln Q$ $E^\circ = (RT/nF) \times 2.303 \log K$ $= (RT/nF) \times \ln K$ $E = E^\circ - \frac{0.0592}{n} \log Q \text{ (at 25 }^\circ\text{C)}$
<p><b>Acids and Bases</b></p> $pK_w = \text{pH} + \text{pOH} = 14.00$ $pK_w = \text{p}K_a + \text{p}K_b = 14.00$ $\text{pH} = \text{p}K_a + \log \{ [A^-] / [HA] \}$	<p><b>Gas Laws</b></p> $PV = nRT$ $(P + n^2 a/V^2)(V - nb) = nRT$
<p><b>Colligative properties</b></p> $\pi = cRT$ $P_{\text{solution}} = X_{\text{solvent}} \times P^\circ_{\text{solvent}}$ $p = kc$ $\Delta T_f = K_f m$ $\Delta T_b = K_b m$	<p><b>Kinetics</b></p> $t_{1/2} = \ln 2/k$ $k = Ae^{-E_a/RT}$ $\ln[A] = \ln[A]_0 - kt$ $\ln \frac{k_2}{k_1} = \frac{E_a}{R} \left( \frac{1}{T_1} - \frac{1}{T_2} \right)$
<p><b>Radioactivity</b></p> $t_{1/2} = \ln 2/\lambda$ $A = \lambda N$ $\ln(N_0/N_t) = \lambda t$ $^{14}\text{C age} = 8033 \ln(A_0/A_t) \text{ years}$	<p><b>Thermodynamics &amp; Equilibrium</b></p> $\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$ $\Delta G = \Delta G^\circ + RT \ln Q$ $\Delta G^\circ = -RT \ln K$ $K_p = K_c (RT)^{\Delta n}$
<p><b>Miscellaneous</b></p> $A = -\log_{10} \frac{I}{I_0}$ $A = \epsilon cl$ $E = -A \frac{e^2}{4\pi\epsilon_0 r} N_A$	<p><b>Mathematics</b></p> <p>If <math>ax^2 + bx + c = 0</math>, then <math>x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}</math></p> $\ln x = 2.303 \log x$

## PERIODIC TABLE OF THE ELEMENTS

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
1 HYDROGEN <b>H</b> 1.008																	2 HELIUM <b>He</b> 4.003
3 LITHIUM <b>Li</b> 6.941	4 BERYLLIUM <b>Be</b> 9.012											5 BORON <b>B</b> 10.81	6 CARBON <b>C</b> 12.01	7 NITROGEN <b>N</b> 14.01	8 OXYGEN <b>O</b> 16.00	9 FLUORINE <b>F</b> 19.00	10 NEON <b>Ne</b> 20.18
11 SODIUM <b>Na</b> 22.99	12 MAGNESIUM <b>Mg</b> 24.31											13 ALUMINIUM <b>Al</b> 26.98	14 SILICON <b>Si</b> 28.09	15 PHOSPHORUS <b>P</b> 30.97	16 SULFUR <b>S</b> 32.07	17 CHLORINE <b>Cl</b> 35.45	18 ARGON <b>Ar</b> 39.95
19 POTASSIUM <b>K</b> 39.10	20 CALCIUM <b>Ca</b> 40.08	21 SCANDIUM <b>Sc</b> 44.96	22 TITANIUM <b>Ti</b> 47.88	23 VANADIUM <b>V</b> 50.94	24 CHROMIUM <b>Cr</b> 52.00	25 MANGANESE <b>Mn</b> 54.94	26 IRON <b>Fe</b> 55.85	27 COBALT <b>Co</b> 58.93	28 NICKEL <b>Ni</b> 58.69	29 COPPER <b>Cu</b> 63.55	30 ZINC <b>Zn</b> 65.39	31 GALLIUM <b>Ga</b> 69.72	32 GERMANIUM <b>Ge</b> 72.59	33 ARSENIC <b>As</b> 74.92	34 SELENIUM <b>Se</b> 78.96	35 BROMINE <b>Br</b> 79.90	36 KRYPTON <b>Kr</b> 83.80
37 RUBIDIUM <b>Rb</b> 85.47	38 STRONTIUM <b>Sr</b> 87.62	39 YTRIUM <b>Y</b> 88.91	40 ZIRCONIUM <b>Zr</b> 91.22	41 NIوبيUM <b>Nb</b> 92.91	42 MOLYBDENUM <b>Mo</b> 95.94	43 TECHNETIUM <b>Tc</b> [98.91]	44 RUTHENIUM <b>Ru</b> 101.07	45 RHODIUM <b>Rh</b> 102.91	46 PALLADIUM <b>Pd</b> 106.4	47 SILVER <b>Ag</b> 107.87	48 CADMIUM <b>Cd</b> 112.40	49 INDIUM <b>In</b> 114.82	50 TIN <b>Sn</b> 118.69	51 ANTIMONY <b>Sb</b> 121.75	52 TELLURIUM <b>Te</b> 127.60	53 IODINE <b>I</b> 126.90	54 XENON <b>Xe</b> 131.30
55 CAESIUM <b>Cs</b> 132.91	56 BARIUM <b>Ba</b> 137.34	57-71	72 HAFNIUM <b>Hf</b> 178.49	73 TANTALUM <b>Ta</b> 180.95	74 TUNGSTEN <b>W</b> 183.85	75 RHENIUM <b>Re</b> 186.2	76 OSMIUM <b>Os</b> 190.2	77 IRIDIUM <b>Ir</b> 192.22	78 PLATINUM <b>Pt</b> 195.09	79 GOLD <b>Au</b> 196.97	80 MERCURY <b>Hg</b> 200.59	81 THALLIUM <b>Tl</b> 204.37	82 LEAD <b>Pb</b> 207.2	83 BISMUTH <b>Bi</b> 208.98	84 POLONIUM <b>Po</b> [210.0]	85 ASTATINE <b>At</b> [210.0]	86 RADON <b>Rn</b> [222.0]
87 FRANCIUM <b>Fr</b> [223.0]	88 RADIUM <b>Ra</b> [226.0]	89-103	104 RUTHERFORDIUM <b>Rf</b> [261]	105 DUBNIUM <b>Db</b> [262]	106 SEABORGIUM <b>Sg</b> [266]	107 BOHRIUM <b>Bh</b> [262]	108 HASSIUM <b>Hs</b> [265]	109 MEITNERIUM <b>Mt</b> [266]	110 DARMSTADIUM <b>Ds</b> [271]	111 ROENTGENIUM <b>Rg</b> [272]							

LANTHANIDES	57 LANTHANUM <b>La</b> 138.91	58 CERIUM <b>Ce</b> 140.12	59 PRASEODYMIUM <b>Pr</b> 140.91	60 NEODYMIUM <b>Nd</b> 144.24	61 PROMETHIUM <b>Pm</b> [144.9]	62 SAMARIUM <b>Sm</b> 150.4	63 EUROPIUM <b>Eu</b> 151.96	64 GADOLINIUM <b>Gd</b> 157.25	65 TERBIUM <b>Tb</b> 158.93	66 DYSPROSIUM <b>Dy</b> 162.50	67 HOLMIUM <b>Ho</b> 164.93	68 ERBIUM <b>Er</b> 167.26	69 THULIUM <b>Tm</b> 168.93	70 YTTERBIUM <b>Yb</b> 173.04	71 LUTETIUM <b>Lu</b> 174.97
ACTINIDES	89 ACTINIUM <b>Ac</b> [227.0]	90 THORIUM <b>Th</b> 232.04	91 PROTACTINIUM <b>Pa</b> [231.0]	92 URANIUM <b>U</b> 238.03	93 NEPTUNIUM <b>Np</b> [237.0]	94 PLUTONIUM <b>Pu</b> [239.1]	95 AMERICIUM <b>Am</b> [243.1]	96 CURIUM <b>Cm</b> [247.1]	97 BERKELIUM <b>Bk</b> [247.1]	98 CALIFORNIUM <b>Cf</b> [252.1]	99 EINSTEINIUM <b>Es</b> [252.1]	100 FERMIUM <b>Fm</b> [257.1]	101 MENDELEVIUM <b>Md</b> [256.1]	102 NOBELIUM <b>No</b> [259.1]	103 LAWRENCIUM <b>Lr</b> [260.1]