

CHEMISTRY 1B - CHEM1102FIRST SEMESTER EXAMINATION**CONFIDENTIAL****JUNE 2005****TIME ALLOWED: THREE HOURS**

GIVE THE FOLLOWING INFORMATION IN BLOCK LETTERS

FAMILY NAME		SID NUMBER	
OTHER NAMES		TABLE NUMBER	

INSTRUCTIONS TO CANDIDATES

- All questions are to be attempted. There are 19 pages of examinable material.
- Complete the written section of the examination paper in **INK**.
- Read each question carefully. Report the appropriate answer and show all relevant working in the space provided.
- The total score for this paper is 100. The possible score per page is shown in the adjacent tables.
- Each new question of the short answer section begins with a •.
- Electronic calculators, including programmable calculators, may be used. Students are warned, however, that credit may not be given, even for a correct answer, where there is insufficient evidence of the working required to obtain the solution.
- Numerical values required for any question, standard electrode reduction potentials, a Periodic Table and some useful formulas may be found on the separate data sheet.
- Pages 17, 18, 21 & 24 are for rough working only.

OFFICIAL USE ONLY**Multiple choice section**

	Marks	
Pages	Max	Gained
2-12	50	

Short answer section

Page	Marks		Marker
	Max	Gained	
13	6		
14	6		
15	7		
16	6		
19	8		
20	6		
22	7		
23	4		
Total	50		
Check Total			

Marks
3

- Many elemental metals crystallise in one of three cubic forms, either with a face-centred cubic, a body-centred cubic or a simple cubic unit cell. Explain the main differences and similarities between these different crystalline forms.

3

- Teeth are made from hydroxyapatite, $\text{Ca}_5(\text{PO}_4)_3\text{OH}$. Why does an acidic medium promote tooth decay? Use chemical equations where appropriate.

How does the fluoridation of drinking water aid the prevention of tooth decay?

Marks
3

- The molecular formulas of the two water soluble compounds, Na_2O and Cl_2O , look very similar, but their chemical properties are very different. Classify each of these compounds as either acidic or basic in aqueous solution.

What happens to the pH value if you dissolve a significant amount of either of these compounds in neutral water (at 25 °C)?

Explain why these compounds react so differently.

- In a major industrial process alumina, Al_2O_3 , is isolated from bauxite, a mineral consisting of mainly Al_2O_3 and Fe_2O_3 . The first step of the process is treatment of the solid ore with concentrated aqueous NaOH solution. Describe how this step enables the separation of the two compounds. Use chemical equations as part of your explanation.

Later on in the process, CO_2 is used to precipitate Al_2O_3 from solution. Write a chemical equation for this step.

What property of Al_2O_3 is exploited in these two steps?

3

Marks
3


- Consider the compound with formula $[\text{Ni}(\text{en})_2(\text{H}_2\text{O})_2]\text{Br}_2 \cdot 2\text{H}_2\text{O}$.
(en = ethylenediamine = $\text{NH}_2\text{CH}_2\text{CH}_2\text{NH}_2$)

Write the formula of the complex ion.

Write the symbols of the ligand donor atoms.

What is the *d* electron configuration of the metal ion in this complex?**4**

- The data given in the table below were obtained for the reaction between nitric oxide and chlorine at 1400 K.



Experiment Number	Initial $[\text{Cl}_2]$ (mol L^{-1})	Initial $[\text{NO}]$ (mol L^{-1})	Initial Reaction Rate ($\text{mol L}^{-1} \text{ s}^{-1}$)
1	0.10	0.10	0.18
2	0.20	0.10	0.36
3	0.10	0.20	0.72

Deduce the rate law for this reaction.

Rate =

Calculate the value of the rate constant.

k =

- Calculate the pH a 0.200 M solution of acetic acid, CH_3COOH , at 25 °C. (The $\text{p}K_a$ of acetic acid is 4.76).

Marks
6

pH =

Solid sodium acetate, NaCH_3CO_2 , (0.15 mol) was dissolved in 0.500 L of 0.200 M acetic acid and the volume made up to 750 mL with water. What is the pH of the resulting solution?









pH =

How much more NaCH_3CO_2 needs to be dissolved in the above solution to give a final pH of 5.00?

Answer:



Marks
8

- Draw the constitutional structure of the major organic product formed in the following reactions. Indicate the correct isomer where appropriate.




Marks
6

- Devise a synthesis of the following compounds from the starting material indicated. Note that more than one step may be required and you should indicate all necessary steps and the constitutional formulas of any intermediate compounds.



Marks
5

- Consider the following reaction sequences.



List the reagents **A - E**.

A

B

C

D


E

- Compare the acidity of a phenol to that of a carboxylic acid.



2

- Draw the repeating unit of the polymers formed in the following reactions.

Marks
2




+



- Write the constitutional formulas of the major products from the acid catalysed hydrolysis of the following dipeptide.

2



CHEM1102 - CHEMISTRY 1B
DATA SHEET

Physical constants

Avogadro constant, $N_A = 6.022 \times 10^{23} \text{ mol}^{-1}$

Faraday constant, $F = 96485 \text{ C mol}^{-1}$

Planck constant, $h = 6.626 \times 10^{-34} \text{ J s}$

Speed of light in vacuum, $c = 2.998 \times 10^8 \text{ m s}^{-1}$

Rydberg constant, $E_R = 2.18 \times 10^{-18} \text{ J}$

Boltzmann constant, $k_B = 1.381 \times 10^{-23} \text{ J K}^{-1}$

Gas constant, $R = 8.314 \text{ J K}^{-1} \text{ mol}^{-1}$
 $= 0.08206 \text{ L atm K}^{-1} \text{ mol}^{-1}$

Charge of electron, $e = 1.602 \times 10^{-19} \text{ C}$

Mass of electron, $m_e = 9.1094 \times 10^{-31} \text{ kg}$

Mass of proton, $m_p = 1.6726 \times 10^{-27} \text{ kg}$

Mass of neutron, $m_n = 1.6749 \times 10^{-27} \text{ kg}$

Properties of matter

Volume of 1 mole of ideal gas at 1 atm and 25 °C = 24.5 L

Volume of 1 mole of ideal gas at 1 atm and 0 °C = 22.4 L

Density of water at 298 K = 0.997 g cm⁻³

Conversion factors

1 atm = 760 mmHg = 101.3 kPa

0 °C = 273 K

1 L = 10⁻³ m³

1 Å = 10⁻¹⁰ m

1 eV = 1.602 × 10⁻¹⁹ J

1 Ci = 3.70 × 10¹⁰ Bq

1 Hz = 1 s⁻¹

Decimal fractions

Fraction	Prefix	Symbol
10 ⁻³	milli	m
10 ⁻⁶	micro	μ
10 ⁻⁹	nano	n
10 ⁻¹²	pico	p

Decimal multiples

Multiple	Prefix	Symbol
10 ³	kilo	k
10 ⁶	mega	M
10 ⁹	giga	G

CHEM1102 - CHEMISTRY 1B*Standard Reduction Potentials, E°*

Reaction	E° / V
$\text{Co}^{3+}(\text{aq}) + \text{e}^- \rightarrow \text{Co}^{2+}(\text{aq})$	+1.82
$\text{Ce}^{4+}(\text{aq}) + \text{e}^- \rightarrow \text{Ce}^{3+}(\text{aq})$	+1.72
$\text{Cl}_2 + 2\text{e}^- \rightarrow 2\text{Cl}^-(\text{aq})$	+1.36
$\text{O}_2 + 4\text{H}^+(\text{aq}) + 4\text{e}^- \rightarrow 2\text{H}_2\text{O}$	+1.23
$\text{Pd}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Pd}(\text{s})$	+0.92
$\text{Ag}^+(\text{aq}) + \text{e}^- \rightarrow \text{Ag}(\text{s})$	+0.80
$\text{Fe}^{3+}(\text{aq}) + \text{e}^- \rightarrow \text{Fe}^{2+}(\text{aq})$	+0.77
$\text{Cu}^+(\text{aq}) + \text{e}^- \rightarrow \text{Cu}(\text{s})$	+0.53
$\text{Cu}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Cu}(\text{s})$	+0.34
$\text{Sn}^{4+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Sn}^{2+}(\text{aq})$	+0.15
$2\text{H}^+(\text{aq}) + 2\text{e}^- \rightarrow \text{H}_2(\text{g})$	0 (by definition)
$\text{Fe}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Fe}(\text{s})$	-0.04
$\text{Pb}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Pb}(\text{s})$	-0.13
$\text{Sn}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Sn}(\text{s})$	-0.14
$\text{Ni}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Ni}(\text{s})$	-0.24
$\text{Co}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Co}(\text{s})$	-0.28
$\text{Fe}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Fe}(\text{s})$	-0.44
$\text{Cr}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Cr}(\text{s})$	-0.74
$\text{Zn}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Zn}(\text{s})$	-0.76
$2\text{H}_2\text{O} + 2\text{e}^- \rightarrow \text{H}_2(\text{g}) + 2\text{OH}^-(\text{aq})$	-0.83
$\text{Cr}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Cr}(\text{s})$	-0.89
$\text{Al}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Al}(\text{s})$	-1.68
$\text{Mg}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Mg}(\text{s})$	-2.36
$\text{Na}^+(\text{aq}) + \text{e}^- \rightarrow \text{Na}(\text{s})$	-2.71

CHEM1102 - CHEMISTRY 1B

Useful formulas

<p>Quantum Chemistry</p> $E = h\nu = hc/\lambda$ $\lambda = h/mv$ $4.5k_B T = hc/\lambda$ $E = Z^2 E_R (1/n^2)$	<p>Radioactivity</p> $t_{1/2} = \ln 2 / \lambda$ $A = \lambda N$ $\ln(N_0/N_t) = \lambda t$ $^{14}\text{C age} = 8033 \ln(A_0/A_t)$
<p>Acids and Bases</p> $pK_w = \text{pH} + \text{pOH} = 14.00$ $pK_w = pK_a + pK_b = 14.00$ $\text{pH} = pK_a + \log\{[A^-] / [\text{HA}]\}$	<p>Gas Laws</p> $PV = nRT$ $(P + n^2 a/V^2)(V - nb) = nRT$
<p>Colligative properties</p> $\pi = cRT$ $P_{\text{solution}} = X_{\text{solvent}} \times P_{\text{solvent}}^\circ$ $p = kc$ $\Delta T_f = K_f m$ $\Delta T_b = K_b m$	<p>Kinetics</p> $t_{1/2} = \ln 2 / k$ $k = Ae^{-E_a/RT}$ $\ln[A] = \ln[A]_0 - kt$ $\ln \frac{k_2}{k_1} = \frac{E_a}{R} \left(\frac{1}{T_1} - \frac{1}{T_2} \right)$
<p>Electrochemistry</p> $\Delta G^\circ = -nFE^\circ$ <p>Moles of $e^- = It/F$</p> $E = E^\circ - (RT/nF) \times 2.303 \log Q$ $= E^\circ - (RT/nF) \times \ln Q$ $E^\circ = (RT/nF) \times 2.303 \log K$ $= (RT/nF) \times \ln K$ $E = E^\circ - \frac{0.0592}{n} \log Q \text{ (at } 25^\circ\text{C)}$	<p>Thermodynamics & Equilibrium</p> $\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$ $\Delta G = \Delta G^\circ + RT \ln Q$ $\Delta G^\circ = -RT \ln K$ $K_p = K_c (RT)^{\Delta n}$
<p>Polymers</p> $R_g = \sqrt{\frac{nl_0^2}{6}}$	<p>Mathematics</p> <p>If $ax^2 + bx + c = 0$, then $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$</p> $\ln x = 2.303 \log x$

PERIODIC TABLE OF THE ELEMENTS

June 2005

CHEM1102 - CHEMISTRY 1B

1		2		3		4		5		6		7		8		9		10		11		12		13		14		15		16		17		18			
1 HYDROGEN H 1.008																								5 BORON B 10.81		6 CARBON C 12.01		7 NITROGEN N 14.01		8 OXYGEN O 16.00		9 FLUORINE F 19.00		10 NEON Ne 20.18			
3 LITHIUM Li 6.941		4 BERYLLIUM Be 9.012																						13 ALUMINIUM Al 26.98		14 SILICON Si 28.09		15 PHOSPHORUS P 30.97		16 SULFUR S 32.07		17 CHLORINE Cl 35.45		18 ARGON Ar 39.95			
11 SODIUM Na 22.99		12 MAGNESIUM Mg 24.31																																			
19 POTASSIUM K 39.10		20 CALCIUM Ca 40.08		21 SCANDIUM Sc 44.96		22 TITANIUM Ti 47.88		23 VANADIUM V 50.94		24 CHROMIUM Cr 52.00		25 MANGANESE Mn 54.94		26 IRON Fe 55.85		27 COBALT Co 58.93		28 NICKEL Ni 58.69		29 COPPER Cu 63.55		30 ZINC Zn 65.39		31 GALLIUM Ga 69.72		32 GERMANIUM Ge 72.59		33 ARSENIC As 74.92		34 SELENIUM Se 78.96		35 BROMINE Br 79.90		36 KRYPTON Kr 83.80			
37 RUBIDIUM Rb 85.47		38 STRONTIUM Sr 87.62		39 YTRIUM Y 88.91		40 ZIRCONIUM Zr 91.22		41 NIوبيUM Nb 92.91		42 MOLYBDENUM Mo 95.94		43 TECHNETIUM Tc [98.91]		44 RUTHENIUM Ru 101.07		45 RHODIUM Rh 102.91		46 PALLADIUM Pd 106.4		47 SILVER Ag 107.87		48 CADMIUM Cd 112.40		49 INDIUM In 114.82		50 TIN Sn 118.69		51 ANTIMONY Sb 121.75		52 TELLURIUM Te 127.60		53 IODINE I 126.90		54 XENON Xe 131.30			
55 CAESIUM Cs 132.91		56 BARIUM Ba 137.34		57-71		72 HAFNIUM Hf 178.49		73 TANTALUM Ta 180.95		74 TUNGSTEN W 183.85		75 RHENIUM Re 186.2		76 OSMIUM Os 190.2		77 IRIDIUM Ir 192.22		78 PLATINUM Pt 195.09		79 GOLD Au 196.97		80 MERCURY Hg 200.59		81 THALLIUM Tl 204.37		82 LEAD Pb 207.2		83 BISMUTH Bi 208.98		84 POLONIUM Po [210.0]		85 ASTATINE At [210.0]		86 RADON Rn [222.0]			
87 FRANCIUM Fr [223.0]		88 RADIUM Ra [226.0]		89-103		104 RUTHERFORDIUM Rf [261]		105 DUBNIUM Db [262]		106 SEABORGIUM Sg [266]		107 BOHRIUM Bh [262]		108 HASSIUM Hs [265]		109 MEITNERIUM Mt [266]																					

LANTHANIDES

57 LANTHANUM La 138.91	58 CERIUM Ce 140.12	59 PRASEODYMIUM Pr 140.91	60 NEODYMIUM Nd 144.24	61 PROMETHIUM Pm [144.9]	62 SAMARIUM Sm 150.4	63 EUROPIUM Eu 151.96	64 GADOLINIUM Gd 157.25	65 TERBIUM Tb 158.93	66 DYSPROSIUM Dy 162.50	67 HOLMIUM Ho 164.93	68 ERBIUM Er 167.26	69 THULIUM Tm 168.93	70 YTTERBIUM Yb 173.04	71 LUTETIUM Lu 174.97
--	-------------------------------------	---	--	--	--------------------------------------	---------------------------------------	---	--------------------------------------	---	--------------------------------------	-------------------------------------	--------------------------------------	--	---------------------------------------

ACTINIDES

89 ACTINIUM Ac [227.0]	90 THORIUM Th 232.04	91 PROTACTINIUM Pa [231.0]	92 URANIUM U 238.03	93 NEPTUNIUM Np [237.0]	94 PLUTONIUM Pu [239.1]	95 AMERICIUM Am [243.1]	96 CURIUM Cm [247.1]	97 BERKELIUM Bk [247.1]	98 CALIFORNIUM Cf [252.1]	99 EINSTEINIUM Es [252.1]	100 FERMIUM Fm [257.1]	101 MENDELEVIUM Md [256.1]	102 NOBELIUM No [259.1]	103 LAWRENCIUM Lr [260.1]
--	--------------------------------------	--	-------------------------------------	---	---	---	--------------------------------------	---	---	---	--	--	---	---

22/07(b)