

2206(a)

THE UNIVERSITY OF SYDNEY

CHEMISTRY 1B - CHEM1102

FIRST SEMESTER EXAMINATION

CONFIDENTIAL

JUNE 2014
TIME ALLOWED: THREE HOURS

GIVE THE FOLLOWING INFORMATION IN BLOCK LETTERS

FAMILY NAME		SID NUMBER	
OTHER NAMES		TABLE NUMBER	

INSTRUCTIONS TO CANDIDATES

- All questions are to be attempted. There are 18 pages of examinable material.
- Complete the written section of the examination paper in **INK**.
- Read each question carefully. Report the appropriate answer and show all relevant working in the space provided.
- The total score for this paper is 100. The possible score per page is shown in the adjacent tables.
- Each new short answer question begins with a •.
- Only non-programmable, University-approved calculators may be used.
- Students are warned that credit may not be given, even for a correct answer, where there is insufficient evidence of the working required to obtain the solution.
- Numerical values required for any question, standard electrode reduction potentials, a Periodic Table and some useful formulas may be found on the separate data sheets.
- Page 20 is for rough working only.

OFFICIAL USE ONLY


Multiple choice section

		Marks	
Pages	Max	Gained	
2-8	28		

Short answer section

Page	Marks		Marker
	Max	Gained	
9	6		
10	5		
11	7		
12	4		
13	8		
14	6		
15	11		
16	7		
17	8		
18	4		
19	6		
Total	72		
Check Total			

- The diagram below shows the structure of an alloy of copper and gold with a gold atom at each of the corners and a copper atom in the centre of each of the faces.



● = Au

● = Cu

What is the chemical formula of the alloy?

	Answer:
--	---------

- Compounds of *d*-block elements are frequently paramagnetic. Using the box notation to represent atomic orbitals, account for this property in compounds of Co^{2+} .



	2
--	---

- Briefly explain how a catalyst works.

	2
--	---

- The structures of the drugs aspirin and benzocaine are shown below.
 - Draw the conjugate base of aspirin and the conjugate acid of benzocaine.
 - Circle the form of each that will be present in a highly acidic environment.

Marks
5

 aspirin	conjugate base of aspirin
 benzocaine	conjugate acid of benzocaine

Ions are less likely to cross cell membranes than uncharged molecules. One of the drugs above is absorbed in the acid environment of the stomach and the other is absorbed in the basic environment of the intestine. Identify which is absorbed in each environment below and *briefly* explain your answers.

Drug absorbed in the stomach:

aspirin / benzocaine

Drug absorbed in the intestine:

aspirin / benzocaine

Aspirin, $C_9H_8O_4$ is not very soluble in water. "Soluble aspirin", the sodium salt $NaC_9H_7O_4$, is often administered instead. Is a solution of "soluble aspirin" acidic or basic? Briefly explain your answer.

THIS QUESTION CONTINUES ON THE NEXT PAGE.

Calculate the pH of a 0.010 M solution of aspirin at 25 °C. The pK_a of aspirin is 3.5 at this temperature.

Marks
7

pH =

Ammonia, NH_3 , is a weak base in water. Write the equation for the acid/base reaction between aspirin and ammonia.

What is the expression for the equilibrium constant, K , for this reaction?

Rewrite this expression in terms of the K_a of aspirin and the K_a of NH_4^+ . (Hint: multiply by $[H^+]/[H^+] = 1$) Hence calculate the value of K . The pK_a of NH_4^+ is 9.2.

Answer:

Would aspirin dissolve in a solution of ammonia? Explain your answer.

- Name the complex $[\text{CoCl}_2(\text{en})_2]$. en = ethylenediamine = $\text{NH}_2\text{CH}_2\text{CH}_2\text{NH}_2$

Marks
4

Draw all possible isomers of this complex.

THE REMAINDER OF THIS PAGE IS FOR ROUGH WORKING ONLY.

- A solution is prepared that contains sodium chloride and sodium chromate (both 0.10 M). When a concentrated solution of silver nitrate is added slowly, white AgCl(s) begins to precipitate. After most of the $\text{Cl}^{\text{-(aq)}}$ has been consumed, red $\text{Ag}_2\text{CrO}_4\text{(s)}$ starts to precipitate.

Ignoring dilution, what is the concentration of silver ions when silver chloride solid first starts to precipitate? $K_{\text{sp}}(\text{AgCl})$ is 1.8×10^{-10} .

Answer:

Ignoring dilution, what is the concentration of silver ions when silver chromate solid first starts to precipitate? $K_{\text{sp}}(\text{Ag}_2\text{CrO}_4)$ is 3.6×10^{-12} .

Answer:

What is the concentration of chloride ions when silver chromate solid first starts to precipitate?

Answer:

What percentage of the chloride ion is precipitated before any silver chromate is precipitated?


Answer:

Marks
8

- Solid sulfur can exist in two forms, rhombic sulfur and monoclinic sulfur. A portion of the phase diagram for sulfur is reproduced schematically below. The pressure and temperature axes are not drawn to scale.

Complete the diagram by adding the labels “vapour” and “liquid” to the appropriate regions.

Marks
6



Which form of solid sulfur is stable at 25 °C and 1 atm?

Describe what happens when sulfur at 25 °C is slowly heated to 200 °C at a constant pressure of 1 atm.










How many triple points are there in the phase diagram?

What phases are in equilibrium at the triple points?

Which solid form of sulfur is more dense? Explain your reasoning.

- Complete the following table. Make sure you give the name of the starting material where indicated.


Marks
11

STARTING MATERIAL	REAGENTS/ CONDITIONS	STRUCTURAL FORMULA(S) OF MAJOR ORGANIC PRODUCT(S)
 Name:	HBr / CCl ₄ (solvent)	
	NaOH	
	KCN / ethanol (solvent)	
 Name:		
	(CH ₃) ₂ NH	
	hot 3 M NaOH	
		


- Methylphenidate, also known as Ritalin, is a psychostimulant drug approved for the treatment of attention-deficit disorder. Identify all stereogenic (chiral) centres in methylphenidate by clearly marking each with an asterisk (*) on the structure below.

Marks
7

methylphenidate




Using one stereogenic centre you have identified, draw the (*R*)-configuration of that centre.



How many stereoisomers are there of methylphenidate? Describe the relationships between these isomers.

Give the products formed when methylphenidate is hydrolysed with 4 M HCl.

- The structure of (–)-linalool, a commonly occurring natural product, is shown below.



Marks
8

What is the molecular formula of (–)-linalool?

Which of the following best describes (–)-linalool?
achiral compound, racemic mixture,
(*R*)-enantiomer, or (*S*)-enantiomer

What functional groups are present in (–)-linalool?


Is it possible to obtain (*Z*) and (*E*) isomers of (–)-linalool? Give a reason for your answer.

Give the structural formula of the organic product formed from (–)-linalool in each of the following reactions. NB: If there is no reaction, write "no reaction".

Reagents / Conditions	Structural Formula of Product
Br ₂ (in CCl ₄ as solvent)	
Na ₂ Cr ₂ O ₇ in aqueous acid	
Na, then CH ₃ Br	
H ₂ / Pd-C catalyst	

- Concentrated HCl reacts with 2-methyl-2-propanol in an S_N1 reaction to give 2-chloro-2-methylpropane as shown below. Complete the reaction mechanism by adding curly arrows and formal charges on the intermediates as appropriate.

Marks
4



Explain what each part of the abbreviation S_N1 means.

S =



N =

1 =

THE REMAINDER OF THIS PAGE IS FOR ROUGH WORKING ONLY.

- Show clearly the reagents you would use to carry out the following chemical conversions. More than one step is required in each case. Give the structure of any intermediate compounds formed.

Marks
6



CHEM1102 –CHEMISTRY 1B

DATA SHEET

*Physical constants*Avogadro constant, $N_A = 6.022 \times 10^{23} \text{ mol}^{-1}$ Faraday constant, $F = 96485 \text{ C mol}^{-1}$ Planck constant, $h = 6.626 \times 10^{-34} \text{ J s}$ Speed of light in vacuum, $c = 2.998 \times 10^8 \text{ m s}^{-1}$ Rydberg constant, $E_R = 2.18 \times 10^{-18} \text{ J}$ Boltzmann constant, $k_B = 1.381 \times 10^{-23} \text{ J K}^{-1}$ Permittivity of a vacuum, $\epsilon_0 = 8.854 \times 10^{-12} \text{ C}^2 \text{ J}^{-1} \text{ m}^{-1}$ Gas constant, $R = 8.314 \text{ J K}^{-1} \text{ mol}^{-1}$
 $= 0.08206 \text{ L atm K}^{-1} \text{ mol}^{-1}$ Charge of electron, $e = 1.602 \times 10^{-19} \text{ C}$ Mass of electron, $m_e = 9.1094 \times 10^{-31} \text{ kg}$ Mass of proton, $m_p = 1.6726 \times 10^{-27} \text{ kg}$ Mass of neutron, $m_n = 1.6749 \times 10^{-27} \text{ kg}$ *Properties of matter*

Volume of 1 mole of ideal gas at 1 atm and 25 °C = 24.5 L

Volume of 1 mole of ideal gas at 1 atm and 0 °C = 22.4 L

Density of water at 298 K = 0.997 g cm⁻³*Conversion factors*

1 atm = 760 mmHg = 101.3 kPa

1 Ci = 3.70 × 10¹⁰ Bq

0 °C = 273 K

1 Hz = 1 s⁻¹1 L = 10⁻³ m³1 tonne = 10³ kg1 Å = 10⁻¹⁰ m1 W = 1 J s⁻¹1 eV = 1.602 × 10⁻¹⁹ J1 J = 1 kg m² s⁻²*Decimal fractions*

Fraction	Prefix	Symbol
10 ⁻³	milli	m
10 ⁻⁶	micro	μ
10 ⁻⁹	nano	n
10 ⁻¹²	pico	p

Decimal multiples

Multiple	Prefix	Symbol
10 ³	kilo	k
10 ⁶	mega	M
10 ⁹	giga	G
10 ¹²	tera	T

CHEM1102 –CHEMISTRY 1B*Standard Reduction Potentials, E°*

Reaction	E° / V
$\text{Co}^{3+}(\text{aq}) + \text{e}^- \rightarrow \text{Co}^{2+}(\text{aq})$	+1.82
$\text{Ce}^{4+}(\text{aq}) + \text{e}^- \rightarrow \text{Ce}^{3+}(\text{aq})$	+1.72
$\text{MnO}_4^-(\text{aq}) + 8\text{H}^+(\text{aq}) + 5\text{e}^- \rightarrow \text{Mn}^{2+}(\text{aq}) + 4\text{H}_2\text{O}$	+1.51
$\text{Au}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Au}(\text{s})$	+1.50
$\text{Cr}_2\text{O}_7^{2-}(\text{aq}) + 14\text{H}^+(\text{aq}) + 6\text{e}^- \rightarrow 2\text{Cr}^{3+}(\text{g}) + 7\text{H}_2\text{O}$	+1.36
$\text{Cl}_2(\text{g}) + 2\text{e}^- \rightarrow 2\text{Cl}^-(\text{aq})$	+1.36
$\text{O}_2(\text{g}) + 4\text{H}^+(\text{aq}) + 4\text{e}^- \rightarrow 2\text{H}_2\text{O}$	+1.23
$\text{Pt}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Pt}(\text{s})$	+1.18
$\text{MnO}_2(\text{s}) + 4\text{H}^+(\text{aq}) + \text{e}^- \rightarrow \text{Mn}^{3+} + 2\text{H}_2\text{O}$	+0.96
$\text{NO}_3^-(\text{aq}) + 4\text{H}^+(\text{aq}) + 3\text{e}^- \rightarrow \text{NO}(\text{g}) + 2\text{H}_2\text{O}$	+0.96
$\text{Pd}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Pd}(\text{s})$	+0.92
$\text{Ag}^+(\text{aq}) + \text{e}^- \rightarrow \text{Ag}(\text{s})$	+0.80
$\text{Fe}^{3+}(\text{aq}) + \text{e}^- \rightarrow \text{Fe}^{2+}(\text{aq})$	+0.77
$\text{Cu}^+(\text{aq}) + \text{e}^- \rightarrow \text{Cu}(\text{s})$	+0.53
$\text{Cu}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Cu}(\text{s})$	+0.34
$\text{Sn}^{4+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Sn}^{2+}(\text{aq})$	+0.15
$2\text{H}^+(\text{aq}) + 2\text{e}^- \rightarrow \text{H}_2(\text{g})$	0 (by definition)
$\text{Fe}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Fe}(\text{s})$	-0.04
$\text{Pb}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Pb}(\text{s})$	-0.13
$\text{Sn}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Sn}(\text{s})$	-0.14
$\text{Ni}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Ni}(\text{s})$	-0.24
$\text{Cd}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Cd}(\text{s})$	-0.40
$\text{Fe}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Fe}(\text{s})$	-0.44
$\text{Cr}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Cr}(\text{s})$	-0.74
$\text{Zn}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Zn}(\text{s})$	-0.76
$2\text{H}_2\text{O} + 2\text{e}^- \rightarrow \text{H}_2(\text{g}) + 2\text{OH}^-(\text{aq})$	-0.83
$\text{Cr}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Cr}(\text{s})$	-0.89
$\text{Al}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Al}(\text{s})$	-1.68
$\text{Sc}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Sc}(\text{s})$	-2.09
$\text{Mg}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Mg}(\text{s})$	-2.36
$\text{Na}^+(\text{aq}) + \text{e}^- \rightarrow \text{Na}(\text{s})$	-2.71
$\text{Ca}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Ca}(\text{s})$	-2.87
$\text{Li}^+(\text{aq}) + \text{e}^- \rightarrow \text{Li}(\text{s})$	-3.04

CHEM1102 –CHEMISTRY 1B

Useful formulas

<p>Quantum Chemistry</p> $E = h\nu = hc/\lambda$ $\lambda = h/mv$ $E = -Z^2 E_R(1/n^2)$ $\Delta x \cdot \Delta(mv) \geq h/4\pi$ $q = 4\pi r^2 \times 5.67 \times 10^{-8} \times T^4$ $T\lambda = 2.898 \times 10^6 \text{ K nm}$	<p>Electrochemistry</p> $\Delta G^\circ = -nFE^\circ$ <p>Moles of $e^- = It/F$</p> $E = E^\circ - (RT/nF) \times 2.303 \log Q$ $= E^\circ - (RT/nF) \times \ln Q$ $E^\circ = (RT/nF) \times 2.303 \log K$ $= (RT/nF) \times \ln K$ $E = E^\circ - \frac{0.0592}{n} \log Q \text{ (at } 25^\circ \text{C)}$
<p>Acids and Bases</p> $pK_w = \text{pH} + \text{pOH} = 14.00$ $pK_w = \text{p}K_a + \text{p}K_b = 14.00$ $\text{pH} = \text{p}K_a + \log \{[A^-] / [\text{HA}]\}$	<p>Gas Laws</p> $PV = nRT$ $(P + n^2a/V^2)(V - nb) = nRT$ $E_k = \frac{1}{2}mv^2$
<p>Radioactivity</p> $t_{1/2} = \ln 2 / \lambda$ $A = \lambda N$ $\ln(N_0/N_t) = \lambda t$ $^{14}\text{C age} = 8033 \ln(A_0/A_t) \text{ years}$	<p>Kinetics</p> $t_{1/2} = \ln 2 / k$ $k = Ae^{-E_a/RT}$ $\ln[A] = \ln[A]_0 - kt$ $\ln \frac{k_2}{k_1} = \frac{E_a}{R} \left(\frac{1}{T_1} - \frac{1}{T_2} \right)$
<p>Mathematics</p> <p>If $ax^2 + bx + c = 0$, then $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$</p> $\ln x = 2.303 \log x$ <p>Area of circle = πr^2</p> <p>Surface area of sphere = $4\pi r^2$</p> <p>Volume of sphere = $\frac{4}{3} \pi r^3$</p>	<p>Thermodynamics & Equilibrium</p> $\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$ $\Delta G = \Delta G^\circ + RT \ln Q$ $\Delta G^\circ = -RT \ln K$ $\Delta_{\text{univ}} S^\circ = R \ln K$ $\ln \frac{K_2}{K_1} = \frac{-\Delta H^\circ}{R} \left(\frac{1}{T_2} - \frac{1}{T_1} \right)$
<p>Miscellaneous</p> $A = -\log \frac{I}{I_0}$ $A = \epsilon cl$ $E = -A \frac{e^2}{4\pi\epsilon_0 r} N_A$	<p>Colligative Properties & Solutions</p> $\Pi = cRT$ $P_{\text{solution}} = X_{\text{solvent}} \times P^\circ_{\text{solvent}}$ $c = kp$ $\Delta T_f = K_f m$ $\Delta T_b = K_b m$

PERIODIC TABLE OF THE ELEMENTS

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
	1 HYDROGEN H 1.008	4 BERYLLIUM Be 9.012											5 BORON B 10.81	6 CARBON C 12.01	7 NITROGEN N 14.01	8 OXYGEN O 16.00	9 FLUORINE F 19.00	10 NEON Ne 20.18
	3 LITHIUM Li 6.941												13 ALUMINUM Al 26.98	14 SILICON Si 28.09	15 PHOSPHORUS P 30.97	16 SULFUR S 32.07	17 CHLORINE Cl 35.45	18 ARGON Ar 39.95
	11 SODIUM Na 22.99	12 MAGNESIUM Mg 24.31											26.98	28.09	30.97	32.07	35.45	39.95
	19 POTASSIUM K 39.10	20 CALCIUM Ca 40.08	21 SCANDIUM Sc 44.96	22 TITANIUM Ti 47.88	23 VANADIUM V 50.94	24 CHROMIUM Cr 52.00	25 MANGANESE Mn 54.94	26 IRON Fe 55.85	27 COBALT Co 58.93	28 NICKEL Ni 58.69	29 COPPER Cu 63.55	30 ZINC Zn 65.39	31 GALLIUM Ga 69.72	32 GERMANIUM Ge 72.59	33 ARSENIC As 74.92	34 SELENIUM Se 78.96	35 BROMINE Br 79.90	36 KRYPTON Kr 83.80
	37 BROMINE Rb 85.47	38 STRONTIUM Sr 87.62	39 YTRBIUM Y 88.91	40 ZIRCONIUM Zr 91.22	41 NIOBIUM Nb 92.91	42 MOLYBDENUM Mo 95.94	43 TECHNETIUM Tc [98.91]	44 RHENIUM Ru 101.07	45 RHODIUM Rh 102.91	46 PALLADIUM Pd 106.4	47 SILVER Ag 107.87	48 Cadmium Cd 112.40	49 INDIUM In 114.82	50 TIN Sn 118.69	51 ANTIMONY Sb 121.75	52 TELLURIUM Te 127.60	53 IODINE I 126.90	54 XENON Xe 131.30
	55 CAESIUM Cs 132.91	56 BARIUM Ba 137.34	57-71 LANTHANIDS	72 HAFNIUM Hf 178.49	73 TANTALUM Ta 180.95	74 TUNGSTEN W 183.85	75 RHENIUM Re 186.2	76 OSMIUM Os 190.2	77 IRIDIUM Ir 192.22	78 PLATINUM Pt 195.09	79 GOLD Au 196.97	80 MERCURY Hg 200.59	81 THALLIUM Tl 204.37	82 LEAD Pb 207.2	83 BISMUTH Bi 208.98	84 POLONIUM Po [210.0]	85 ASTATINE At [210.0]	86 RADON Rn [222.0]
	87 FRANCIUM Fr [223.0]	88 RADIUM Ra [226.0]		104 RIFERBERORIUM Rf [263]	105 DUBNIUM Db [268]	106 SEABERGIUM Sg [271]	107 BOHRIUM Bh [274]	108 HASSIUM Hs [270]	109 MEITNERIUM Mt [278]	110 DARMSTADTIUM Ds [281]	111 ROSTENIUM Rg [281]	112 COOPERIUM Cn [285]		114 FLEROVIUM Fl [289]		116 LIVERMORIUM Lv [293]		

	57 LANTHANUM La 138.91	58 CEURIUM Ce 140.12	59 PRASEODYMIUM Pr 140.91	60 NEODYMIUM Nd 144.24	61 PROMETHIUM Pm [144.91]	62 SAMARIUM Sm 150.4	63 EUROPIUM Eu 151.96	64 GADOLINIUM Gd 157.25	65 TERBIUM Tb 158.93	66 DYSPROSIUM Dy 162.50	67 HOIMIUM Ho 164.93	68 ERBIUM Er 167.26	69 THULIUM Tm 168.93	70 YTERBIUM Yb 173.04	71 LUTETIUM Lu 174.97
	89 ACTINIUM Ac [227.0]	90 THORIUM Th 232.04	91 PROTACTINIUM Pa [231.0]	92 URANIUM U 238.03	93 NEPTUNIUM Np [237.0]	94 PLUTONIUM Pu [239.1]	95 AMERICIUM Am [243.1]	96 CURIUM Cm [247.1]	97 BERKELIUM Bk [247.1]	98 CALIFORNIUM Cf [252.1]	99 EINSTEINIUM Es [252.1]	100 FERMIUM Fm [257.1]	101 MENDELEVIUM Md [256.1]	102 NOBELIUM No [259.1]	103 LAWRENCIUM Lr [260.1]

ACTINOIDS

LANTHANIDS