

CHEMISTRY 1B - CHEM1102SECOND SEMESTER EXAMINATION**CONFIDENTIAL****NOVEMBER 2006****TIME ALLOWED: THREE HOURS**

GIVE THE FOLLOWING INFORMATION IN BLOCK LETTERS

FAMILY NAME		SID NUMBER	
OTHER NAMES		TABLE NUMBER	

INSTRUCTIONS TO CANDIDATES

- All questions are to be attempted. There are 18 pages of examinable material.
- Complete the written section of the examination paper in **INK**.
- Read each question carefully. Report the appropriate answer and show all relevant working in the space provided.
- The total score for this paper is 100. The possible score per page is shown in the adjacent tables.
- Each new question of the short answer section begins with a •.
- Electronic calculators, including programmable calculators, may be used. Students are warned, however, that credit may not be given, even for a correct answer, where there is insufficient evidence of the working required to obtain the solution.
- Numerical values required for any question, standard electrode reduction potentials, a Periodic Table and some useful formulas may be found on the separate data sheet.
- Page 20 is for rough working only.

OFFICIAL USE ONLY~~Multiple choice section~~

		Marks	
Pages	Max	Gained	
2-10	44		

Short answer section

Page	Marks		Marker
	Max	Gained	
11	2		
12	4		
13	10		
14	8		
15	3		
16	7		
17	6		
18	10		
19	6		
Total	56		


- Oxalic acid, $\text{H}_2\text{C}_2\text{O}_4$, found in rhubarb, causes muscle spasms by precipitating Ca^{2+} ions from the blood as calcium oxalate, $\text{CaC}_2\text{O}_4 \cdot \text{H}_2\text{O}$. Given the solubility product constant for calcium oxalate is $2.3 \times 10^{-9} \text{ M}^2$, calculate the concentration of calcium ions in g L^{-1} formed by dissolving $\text{CaC}_2\text{O}_4 \cdot \text{H}_2\text{O}$ in water at 25°C to give a saturated solution.

Marks
2

Answer:

THE REMAINDER OF THIS PAGE IS FOR ROUGH WORKING ONLY.

- The following initial rate data have been obtained for the gas phase reaction of nitrogen dioxide, $\text{NO}_2(\text{g})$, and ozone, $\text{O}_3(\text{g})$, at 300 K.



$[\text{NO}_2(\text{g})]$ M	$[\text{O}_3(\text{g})]$ M	Rate M s^{-1}
0.65	0.80	2.61×10^4
1.10	0.80	4.40×10^4
1.10	1.60	8.80×10^4

What is the order of this reaction with respect to each reagent?

Marks
4

What is the rate constant of the reaction?

Answer:

- Briefly explain why HF is a weaker Brønsted acid than HI and a stronger acid than H₂O.

Marks
2

- Compounds of *d*-block elements are frequently paramagnetic. Using the box notation to represent atomic orbitals, account for this property in compounds of Cu²⁺.

2

- Complete the following table.

6

Formula	Oxidation state of transition metal	Coordination number of transition metal	Number of <i>d</i> -electrons in the transition metal	Species formed upon dissolving in water
Na ₂ [CoCl ₄]				
[Ni(NH ₃) ₅ (H ₂ O)]SO ₄				
[Cr(en) ₃]Br ₃				

en = ethylenediamine = NH₂CH₂CH₂NH₂

Marks
8

- Solution A consists of a 0.25 M aqueous solution of hydrazoic acid, HN_3 , at 25 °C. Calculate the pH of Solution A. The $\text{p}K_a$ of HN_3 is 4.63.

Answer:

At 25 °C, 1.00 L of Solution B consists of 13.0 g of sodium azide (NaN_3) dissolved in water. Calculate the pH of Solution B.

Answer:

Solution B (1.00 L) is poured into Solution A (1.00 L) and allowed to equilibrate at 25 °C to give Solution C. Calculate the pH of Solution C.

Answer:


If you wanted to adjust the pH of Solution C to be exactly equal to 4.00, which component in the mixture would you need to increase in concentration?

- Define what is meant by an “allotrope”. Give an example of a pair of allotropes involving oxygen and a pair *not* involving oxygen.

Marks
3

THE REMAINDER OF THIS PAGE IS FOR ROUGH WORKING ONLY.

- The structure of lignocaine, a local anaesthetic, is given below.

Marks**7**

Give the molecular formula of lignocaine.

Name the functional groups in lignocaine indicated by boxes “a” and “b”.

a:	b:
-----------	-----------

Give the structure(s) of all organic products formed when lignocaine is treated with the following reagents. If no reaction occurs, write “NO REACTION”.

cold HCl (1 M)







hot NaOH (4 M)

hot HCl (4 M)

THE REMAINDER OF THIS PAGE IS FOR ROUGH WORKING ONLY.


Marks
6

- Draw the constitutional formula of the major organic product formed in each of the following reactions.

**THE REMAINDER OF THIS PAGE IS FOR ROUGH WORKING ONLY.**

- Propionaldehyde (propanal) is treated first with phenylmagnesium bromide in dry diethyl ether and then with dilute aqueous acid, to yield alcohol **G**.

Marks
5



State whether **G** is obtained as the (*R*)-enantiomer, the (*S*)-enantiomer, a racemic mixture, or is achiral.

--

List below, the substituents on the stereogenic carbon atom in **G**, in decreasing priority (*i.e.* from highest to lowest priority), as determined by the sequence rules.

highest priority

lowest priority


--	--	--	--

Draw the (*R*) enantiomer of **G**, showing the correct absolute stereochemistry.

--

- The incomplete proposed mechanism for the reaction of (*E*)-but-2-ene with aqueous acid is shown below. Complete the mechanism by adding curly arrows and relevant lone pairs to illustrate the bonding changes that take place.

5




What two-word description may be used for the name of this mechanism?

--

- Devise a synthesis of 2-phenyl-2-propanol, starting from propene and bromobenzene. Note that your synthetic route will require more than one step from each starting material. Show clearly the reagents you would use and draw constitutional formulas for all intermediate compounds.

Marks
6



THE REMAINDER OF THIS PAGE IS FOR ROUGH WORKING ONLY.

CHEM1102 - CHEMISTRY 1B
DATA SHEET

Physical constants

Avogadro constant, $N_A = 6.022 \times 10^{23} \text{ mol}^{-1}$

Faraday constant, $F = 96485 \text{ C mol}^{-1}$

Planck constant, $h = 6.626 \times 10^{-34} \text{ J s}$

Speed of light in vacuum, $c = 2.998 \times 10^8 \text{ m s}^{-1}$

Rydberg constant, $E_R = 2.18 \times 10^{-18} \text{ J}$

Boltzmann constant, $k_B = 1.381 \times 10^{-23} \text{ J K}^{-1}$

Gas constant, $R = 8.314 \text{ J K}^{-1} \text{ mol}^{-1}$
 $= 0.08206 \text{ L atm K}^{-1} \text{ mol}^{-1}$

Charge of electron, $e = 1.602 \times 10^{-19} \text{ C}$

Mass of electron, $m_e = 9.1094 \times 10^{-31} \text{ kg}$

Mass of proton, $m_p = 1.6726 \times 10^{-27} \text{ kg}$

Mass of neutron, $m_n = 1.6749 \times 10^{-27} \text{ kg}$

Properties of matter

Volume of 1 mole of ideal gas at 1 atm and 25 °C = 24.5 L

Volume of 1 mole of ideal gas at 1 atm and 0 °C = 22.4 L

Density of water at 298 K = 0.997 g cm⁻³

Conversion factors

1 atm = 760 mmHg = 101.3 kPa

0 °C = 273 K

1 L = 10⁻³ m³

1 Å = 10⁻¹⁰ m

1 eV = 1.602 × 10⁻¹⁹ J

1 Ci = 3.70 × 10¹⁰ Bq

1 Hz = 1 s⁻¹

Decimal fractions

Fraction	Prefix	Symbol
10 ⁻³	milli	m
10 ⁻⁶	micro	μ
10 ⁻⁹	nano	n
10 ⁻¹²	pico	p

Decimal multiples

Multiple	Prefix	Symbol
10 ³	kilo	k
10 ⁶	mega	M
10 ⁹	giga	G

CHEM1102 - CHEMISTRY 1B**Standard Reduction Potentials, E°**

Reaction	E° / V
$\text{Co}^{3+}(\text{aq}) + \text{e}^- \rightarrow \text{Co}^{2+}(\text{aq})$	+1.82
$\text{Ce}^{4+}(\text{aq}) + \text{e}^- \rightarrow \text{Ce}^{3+}(\text{aq})$	+1.72
$\text{Au}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Au}(\text{s})$	+1.50
$\text{Cl}_2 + 2\text{e}^- \rightarrow 2\text{Cl}^-(\text{aq})$	+1.36
$\text{O}_2 + 4\text{H}^+(\text{aq}) + 4\text{e}^- \rightarrow 2\text{H}_2\text{O}$	+1.23
$\text{Br}_2 + 2\text{e}^- \rightarrow 2\text{Br}^-(\text{aq})$	+1.10
$\text{MnO}_2(\text{s}) + 4\text{H}^+(\text{aq}) + \text{e}^- \rightarrow \text{Mn}^{3+} + 2\text{H}_2\text{O}$	+0.96
$\text{Pd}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Pd}(\text{s})$	+0.92
$\text{Ag}^+(\text{aq}) + \text{e}^- \rightarrow \text{Ag}(\text{s})$	+0.80
$\text{Fe}^{3+}(\text{aq}) + \text{e}^- \rightarrow \text{Fe}^{2+}(\text{aq})$	+0.77
$\text{Cu}^+(\text{aq}) + \text{e}^- \rightarrow \text{Cu}(\text{s})$	+0.53
$\text{Cu}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Cu}(\text{s})$	+0.34
$\text{Sn}^{4+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Sn}^{2+}(\text{aq})$	+0.15
$2\text{H}^+(\text{aq}) + 2\text{e}^- \rightarrow \text{H}_2(\text{g})$	0 (by definition)
$\text{Fe}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Fe}(\text{s})$	-0.04
$\text{Pb}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Pb}(\text{s})$	-0.13
$\text{Sn}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Sn}(\text{s})$	-0.14
$\text{Ni}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Ni}(\text{s})$	-0.24
$\text{Co}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Co}(\text{s})$	-0.28
$\text{Fe}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Fe}(\text{s})$	-0.44
$\text{Cr}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Cr}(\text{s})$	-0.74
$\text{Zn}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Zn}(\text{s})$	-0.76
$2\text{H}_2\text{O} + 2\text{e}^- \rightarrow \text{H}_2(\text{g}) + 2\text{OH}^-(\text{aq})$	-0.83
$\text{Cr}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Cr}(\text{s})$	-0.89
$\text{Al}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Al}(\text{s})$	-1.68
$\text{Mg}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Mg}(\text{s})$	-2.36
$\text{Na}^+(\text{aq}) + \text{e}^- \rightarrow \text{Na}(\text{s})$	-2.71
$\text{Ca}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Ca}(\text{s})$	-2.87
$\text{Li}^+(\text{aq}) + \text{e}^- \rightarrow \text{Li}(\text{s})$	-3.04

CHEM1102 - CHEMISTRY 1B

Useful formulas

<p>Quantum Chemistry</p> $E = h\nu = hc/\lambda$ $\lambda = h/m\nu$ $4.5k_B T = hc/\lambda$ $E = Z^2 E_R (1/n^2)$ $\Delta x \cdot \Delta(mv) \geq h/4\pi$ $q = 4\pi r^2 \times 5.67 \times 10^{-8} \times T^4$	<p>Electrochemistry</p> $\Delta G^\circ = -nFE^\circ$ <p>Moles of $e^- = It/F$</p> $E = E^\circ - (RT/nF) \times 2.303 \log Q$ $= E^\circ - (RT/nF) \times \ln Q$ $E^\circ = (RT/nF) \times 2.303 \log K$ $= (RT/nF) \times \ln K$ $E = E^\circ - \frac{0.0592}{n} \log Q \text{ (at 25 }^\circ\text{C)}$
<p>Acids and Bases</p> $pK_w = \text{pH} + \text{pOH} = 14.00$ $pK_w = \text{p}K_a + \text{p}K_b = 14.00$ $\text{pH} = \text{p}K_a + \log \{ [A^-] / [HA] \}$	<p>Gas Laws</p> $PV = nRT$ $(P + n^2 a/V^2)(V - nb) = nRT$
<p>Colligative properties</p> $\pi = cRT$ $P_{\text{solution}} = X_{\text{solvent}} \times P^\circ_{\text{solvent}}$ $p = kc$ $\Delta T_f = K_f m$ $\Delta T_b = K_b m$	<p>Kinetics</p> $t_{1/2} = \ln 2/k$ $k = Ae^{-E_a/RT}$ $\ln[A] = \ln[A]_0 - kt$ $\ln \frac{k_2}{k_1} = \frac{E_a}{R} \left(\frac{1}{T_1} - \frac{1}{T_2} \right)$
<p>Radioactivity</p> $t_{1/2} = \ln 2/\lambda$ $A = \lambda N$ $\ln(N_0/N_t) = \lambda t$ $^{14}\text{C age} = 8033 \ln(A_0/A_t)$	<p>Thermodynamics & Equilibrium</p> $\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$ $\Delta G = \Delta G^\circ + RT \ln Q$ $\Delta G^\circ = -RT \ln K$ $K_p = K_c (RT)^{\Delta n}$
<p>Polymers</p> $R_g = \sqrt{\frac{nl_0^2}{6}}$	<p>Mathematics</p> <p>If $ax^2 + bx + c = 0$, then $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$</p> $\ln x = 2.303 \log x$

PERIODIC TABLE OF THE ELEMENTS

November 2006

CHEM1102 - CHEMISTRY 1B

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
1 HYDROGEN H 1.008																	2 HELIUM He 4.003
3 LITHIUM Li 6.941	4 BERYLLIUM Be 9.012											5 BORON B 10.81	6 CARBON C 12.01	7 NITROGEN N 14.01	8 OXYGEN O 16.00	9 FLUORINE F 19.00	10 NEON Ne 20.18
11 SODIUM Na 22.99	12 MAGNESIUM Mg 24.31											13 ALUMINIUM Al 26.98	14 SILICON Si 28.09	15 PHOSPHORUS P 30.97	16 SULFUR S 32.07	17 CHLORINE Cl 35.45	18 ARGON Ar 39.95
19 POTASSIUM K 39.10	20 CALCIUM Ca 40.08	21 SCANDIUM Sc 44.96	22 TITANIUM Ti 47.88	23 VANADIUM V 50.94	24 CHROMIUM Cr 52.00	25 MANGANESE Mn 54.94	26 IRON Fe 55.85	27 COBALT Co 58.93	28 NICKEL Ni 58.69	29 COPPER Cu 63.55	30 ZINC Zn 65.39	31 GALLIUM Ga 69.72	32 GERMANIUM Ge 72.59	33 ARSENIC As 74.92	34 SELENIUM Se 78.96	35 BROMINE Br 79.90	36 KRYPTON Kr 83.80
37 RUBIDIUM Rb 85.47	38 STRONTIUM Sr 87.62	39 YTRIUM Y 88.91	40 ZIRCONIUM Zr 91.22	41 NOBIUM Nb 92.91	42 MOLYBDENUM Mo 95.94	43 TECHNETIUM Tc [98.91]	44 RUTHENIUM Ru 101.07	45 RHODIUM Rh 102.91	46 PALLADIUM Pd 106.4	47 SILVER Ag 107.87	48 CADMIUM Cd 112.40	49 INDIUM In 114.82	50 TIN Sn 118.69	51 ANTIMONY Sb 121.75	52 TELLURIUM Te 127.60	53 IODINE I 126.90	54 XENON Xe 131.30
55 CAESIUM Cs 132.91	56 BARIUM Ba 137.34	57-71	72 HAFNIUM Hf 178.49	73 TANTALUM Ta 180.95	74 TUNGSTEN W 183.85	75 RHENIUM Re 186.2	76 OSMIUM Os 190.2	77 IRIDIUM Ir 192.22	78 PLATINUM Pt 195.09	79 GOLD Au 196.97	80 MERCURY Hg 200.59	81 THALLIUM Tl 204.37	82 LEAD Pb 207.2	83 BISMUTH Bi 208.98	84 POLONIUM Po [210.0]	85 ASTATINE At [210.0]	86 RADON Rn [222.0]
87 FRANCIUM Fr [223.0]	88 RADIUM Ra [226.0]	89-103	104 RUTHERFORDIUM Rf [261]	105 DUBNIUM Db [262]	106 SEABORGIUM Sg [266]	107 BOHRIUM Bh [262]	108 HASSIUM Hs [265]	109 MEITNERIUM Mt [266]									

LANTHANIDES

57 LANTHANUM La 138.91	58 CERIUM Ce 140.12	59 PRASEODYMIUM Pr 140.91	60 NEODYMIUM Nd 144.24	61 PROMETHIUM Pm [144.9]	62 SAMARIUM Sm 150.4	63 EUROPIUM Eu 151.96	64 GADOLINIUM Gd 157.25	65 TERBIUM Tb 158.93	66 DYSPROSIUM Dy 162.50	67 HOLMIUM Ho 164.93	68 ERBIUM Er 167.26	69 THULIUM Tm 168.93	70 YTTERBIUM Yb 173.04	71 LUTETIUM Lu 174.97
--	-------------------------------------	---	--	--	--------------------------------------	---------------------------------------	---	--------------------------------------	---	--------------------------------------	-------------------------------------	--------------------------------------	--	---------------------------------------

ACTINIDES

89 ACTINIUM Ac [227.0]	90 THORIUM Th 232.04	91 PROTACTINIUM Pa [231.0]	92 URANIUM U 238.03	93 NEPTUNIUM Np [237.0]	94 PLUTONIUM Pu [239.1]	95 AMERICIUM Am [243.1]	96 CURIUM Cm [247.1]	97 BERKELIUM Bk [247.1]	98 CALIFORNIUM Cf [252.1]	99 EINSTEINIUM Es [252.1]	100 FERMIUM Fm [257.1]	101 MENDELEVIUM Md [256.1]	102 NOBELIUM No [259.1]	103 LAWRENCIUM Lr [260.1]
--	--------------------------------------	--	-------------------------------------	---	---	---	--------------------------------------	---	---	---	--	--	---	---

22/08(b)