

CHEMISTRY 1B - CHEM1102SECOND SEMESTER EXAMINATION**CONFIDENTIAL****NOVEMBER 2008****TIME ALLOWED: THREE HOURS**

GIVE THE FOLLOWING INFORMATION IN BLOCK LETTERS

<b>FAMILY NAME</b>		<b>SID NUMBER</b>	
<b>OTHER NAMES</b>		<b>TABLE NUMBER</b>	

**INSTRUCTIONS TO CANDIDATES**

- All questions are to be attempted. There are 22 pages of examinable material.
- Complete the written section of the examination paper in **INK**.
- Read each question carefully. Report the appropriate answer and show all relevant working in the space provided.
- The total score for this paper is 100. The possible score per page is shown in the adjacent tables.
- Each new question of the short answer section begins with a •.
- Electronic calculators, including programmable calculators, may be used. Students are warned, however, that credit may not be given, even for a correct answer, where there is insufficient evidence of the working required to obtain the solution.
- Numerical values required for any question, standard electrode reduction potentials, a Periodic Table and some useful formulas may be found on the separate data sheet.
- Page 24 is for rough working only.

**OFFICIAL USE ONLY****Multiple choice section**

Pages	Marks	
	Max	Gained
2-10	30	

**Short answer section**

Page	Marks		Marker
	Max	Gained	
10	5		
11	7		
12	8		
13	5		
14	3		
15	3		
16	2		
17	4		
18	5		
19	8		
20	5		
21	5		
22	5		
23	5		
<b>Total</b>	<b>70</b>		

- Briefly explain how the concept of electronegativity can rationalise the existence of acidic, basic and amphoteric oxides.

**Marks**  
**3**

Draw the face-centred cubic unit cell.

**2**

A solution is prepared that is 0.10 M in potassium bromide and 0.10 M in potassium chromate. A concentrated aqueous solution of silver nitrate is added with stirring. What is the concentration of  $\text{Ag}^+(\text{aq})$  ions when silver bromide first appears?  
 $K_{\text{sp}}$  of  $\text{AgBr} = 5.0 \times 10^{-13}$

**Marks**  
**4**

Answer:


What is the concentration of  $\text{Ag}^+(\text{aq})$  ions when silver chromate first appears?  
 $K_{\text{sp}}$  of  $\text{Ag}_2\text{CrO}_4 = 2.6 \times 10^{-12}$

Answer:

What is the concentration of  $\text{Br}^-(\text{aq})$  ions when silver chromate first appears?

Answer:

- Calculate the equilibrium constant for the following reaction.



Data:  $K_{\text{stab}}$  of  $[\text{Ag}(\text{CN})_2]^{-} = 3 \times 10^{20}$ ;  $K_{\text{sp}}$  of  $\text{AgI} = 8.3 \times 10^{-17}$

Answer:

**3**

**Marks**  
**2**

- Which of the cations,  $[\text{Fe}(\text{OH}_2)_6]^{3+}$  and  $[\text{Fe}(\text{OH}_2)_6]^{2+}$ , has the larger  $\text{p}K_a$ ? Briefly explain why.

--

**3**

- Consider the compound  $[\text{CrCl}(\text{OH}_2)_4(\text{NCS})]\text{Cl}\cdot 2\text{H}_2\text{O}$ .

What is the oxidation state of the transition metal ion?

--

What is the coordination number of the transition metal ion?

--

How many *d*-electrons in the transition metal ion?

--

List all the ligand donor atoms.

--

- Consider the complexes *cis*- $[\text{PtCl}_2(\text{NH}_3)_2]$  and *trans*- $[\text{PtCl}_2(\text{NH}_3)_2]$ . Draw the structures of the two isomers, clearly illustrating the stereochemistry.

**3**

--

Briefly suggest why *cis*- $[\text{PtCl}_2(\text{NH}_3)_2]$  is an effective anti-cancer drug, but *trans*- $[\text{PtCl}_2(\text{NH}_3)_2]$  is not.

--

**Marks**  
**5**

- Buffers made of mixtures of  $\text{H}_2\text{PO}_4^-$  and  $\text{HPO}_4^{2-}$  are used to control the pH of soft drinks. What is the pH of a 350 mL drink containing 6.0 g of  $\text{NaH}_2\text{PO}_4$  and 4.0 g of  $\text{Na}_2\text{HPO}_4$ ?

For phosphoric acid,  $\text{H}_3\text{PO}_4$ ,  $\text{p}K_{\text{a}1} = 2.15$ ,  $\text{p}K_{\text{a}2} = 7.20$  and  $\text{p}K_{\text{a}3} = 12.38$ .

Briefly describe how this buffer system functions. Use equations where appropriate.

Is this buffer better able to resist changes in pH following the addition of acid or of base? Explain your answer.


- The figure below illustrates the phase diagram for water. The points on the diagram correspond to:

**A:** Triple point ( $0.0098\text{ }^{\circ}\text{C}$ ,  $0.610\text{ kPa}$ )

**B:** Normal melting point ( $0\text{ }^{\circ}\text{C}$ ,  $1.01 \times 10^2\text{ kPa}$ )

**C:** Normal boiling point ( $100\text{ }^{\circ}\text{C}$ ,  $1.01 \times 10^2\text{ kPa}$ )

**D:** Critical point ( $374.4\text{ }^{\circ}\text{C}$ ,  $2.18 \times 10^4\text{ kPa}$ )



Describe all of the phase changes that occur when water at  $1.01 \times 10^2\text{ kPa}$  is slowly warmed from  $-20\text{ }^{\circ}\text{C}$  to  $200\text{ }^{\circ}\text{C}$ .


Describe all of the phase changes that occur when water at  $0\text{ }^{\circ}\text{C}$  is slowly compressed from  $0.500\text{ kPa}$  to  $1000\text{ kPa}$ .

**THIS QUESTION CONTINUES ON THE NEXT PAGE**

**Marks**  
**3**

Addition of salt to water raises its boiling point and lowers its melting point. Sketch the phase diagram for water containing salt, showing how it relates to the phase diagram for water (shown as dotted lines below).


**Marks**  
**3**



In terms of the relative entropies of all relevant species, explain why the boiling point of salt water is higher than that of pure water.

**THE REMAINDER OF THIS PAGE IS FOR ROUGH WORKING ONLY.**

- The following data were obtained for the reaction between gaseous nitric oxide and chlorine at  $-10\text{ }^{\circ}\text{C}$ :



Experiment Number	Initial $P_{\text{NO}}$ (atm)	Initial $P_{\text{Cl}_2}$ (atm)	Initial Reaction Rate (atm $\text{s}^{-1}$ )
1	2.16	2.16	0.065
2	2.16	4.32	0.130
3	4.32	4.32	0.518

Derive an expression for the rate law for this reaction and calculate the value of the rate constant.

**Marks**  
**2**


Rate law:

Rate constant:

**THIS QUESTION CONTINUES ON THE NEXT PAGE**



The mechanism for this reaction has been postulated to be that below.




Work out the rate law expected for this mechanism and hence show that it is consistent with the experimental rate law and the chemical equation.

**Marks****4**

The reaction is exothermic. Draw the potential energy *vs* reaction coordinate diagram for this mechanism, labelling all species that can be isolated.

- The structure of the antihistamine, Zyrtec™, is given below.

**Marks**  
**5**



What is the name of the functional group in Box A?

What is the name of the functional group in Box B?

What is the name of the functional group in Box C?


By drawing an arrow on the structure above, clearly indicate the stereocentre in the structure of Zyrtec.

Draw the product of the reaction when Zyrtec is treated with LiAlH<sub>4</sub> followed by dilute acid.

--

**Marks**  
**8**

- Consider compound **F** shown below.



Assign the stereocentre in compound **F** as (*R*) or (*S*), explaining your reasoning.

--

Draw the enantiomer of compound **F**.

--

When compound **F** is reacted with hot KOH solution, a product (**G**) is formed that shows three peaks in the  $^1\text{H}$  NMR spectrum in the region 7-8 ppm and three peaks in the region 5-6 ppm. Draw the structure of this product.

--

When **G** is reacted with dilute sulfuric acid, a further product, **H**, is formed. **H** has a peak at  $3300\text{ cm}^{-1}$  in its IR spectrum. Draw the structure of product **H**.

--


Is **H** formed as a single enantiomer, as a racemate, or is **H** achiral?

--

Assuming an  $\text{S}_{\text{N}}2$  mechanism, draw the product of the substitution reaction between **F** and  $(\text{CH}_3)_2\text{NH}$ , indicating stereochemistry where appropriate.

--

- Consider the compound **J** below.



**J**

What is the systematic name for compound **J**.

**Marks**  
**5**



Draw a constitutional isomer of **J**.

Draw a configurational isomer of **J**.

Draw the structure of the product formed when compound **J** is reacted with hydrogen gas ( $H_2$ ) and a palladium on carbon (Pd/C) catalyst.

- Complete the following mechanism by adding curly arrows to illustrate the bonding changes that take place.

**Marks**  
**5**




**THE REMAINDER OF THIS PAGE IS FOR ROUGH WORKING ONLY.**

**Marks**  
**5**

- An unknown compound **K** with the molecular formula  $C_4H_8O$  gives the following spectroscopic data.

$^1H$  NMR: 1.06 ppm, triplet, integration = 3H  
2.13 ppm, singlet, integration = 3H  
2.47 ppm, quartet, integration = 2H




IR spectroscopy: stretch at  $1715\text{ cm}^{-1}$ .

Use the information above to deduce the structure of compound **K**. Give reasoning for the structure chosen.

- Devise a synthesis of 1,2-dibromobutane from butanal. Provide any intermediate structures and reagents. (Hint: More than one step is required.)

**Marks**  
**5**



**THE REMAINDER OF THIS PAGE IS FOR ROUGH WORKING ONLY.**

**CHEM1102 - CHEMISTRY 1B**  
**DATA SHEET**

*Physical constants*

Avogadro constant,  $N_A = 6.022 \times 10^{23} \text{ mol}^{-1}$

Faraday constant,  $F = 96485 \text{ C mol}^{-1}$

Planck constant,  $h = 6.626 \times 10^{-34} \text{ J s}$

Speed of light in vacuum,  $c = 2.998 \times 10^8 \text{ m s}^{-1}$

Rydberg constant,  $E_R = 2.18 \times 10^{-18} \text{ J}$

Boltzmann constant,  $k_B = 1.381 \times 10^{-23} \text{ J K}^{-1}$

Permittivity of a vacuum,  $\epsilon_0 = 8.854 \times 10^{-12} \text{ C}^2 \text{ J}^{-1} \text{ m}^{-1}$

Gas constant,  $R = 8.314 \text{ J K}^{-1} \text{ mol}^{-1}$   
 $= 0.08206 \text{ L atm K}^{-1} \text{ mol}^{-1}$

Charge of electron,  $e = 1.602 \times 10^{-19} \text{ C}$

Mass of electron,  $m_e = 9.1094 \times 10^{-31} \text{ kg}$

Mass of proton,  $m_p = 1.6726 \times 10^{-27} \text{ kg}$

Mass of neutron,  $m_n = 1.6749 \times 10^{-27} \text{ kg}$

*Properties of matter*

Volume of 1 mole of ideal gas at 1 atm and 25 °C = 24.5 L

Volume of 1 mole of ideal gas at 1 atm and 0 °C = 22.4 L

Density of water at 298 K = 0.997 g cm<sup>-3</sup>

*Conversion factors*

1 atm = 760 mmHg = 101.3 kPa

1 Ci =  $3.70 \times 10^{10}$  Bq

0 °C = 273 K

1 Hz = 1 s<sup>-1</sup>

1 L = 10<sup>-3</sup> m<sup>3</sup>

1 tonne = 10<sup>3</sup> kg

1 Å = 10<sup>-10</sup> m

1 W = 1 J s<sup>-1</sup>

1 eV =  $1.602 \times 10^{-19}$  J

*Decimal fractions*

Fraction	Prefix	Symbol
10 <sup>-3</sup>	milli	m
10 <sup>-6</sup>	micro	μ
10 <sup>-9</sup>	nano	n
10 <sup>-12</sup>	pico	p

*Decimal multiples*

Multiple	Prefix	Symbol
10 <sup>3</sup>	kilo	k
10 <sup>6</sup>	mega	M
10 <sup>9</sup>	giga	G
10 <sup>12</sup>	tera	T



**CHEM1102 - CHEMISTRY 1B****Standard Reduction Potentials,  $E^\circ$** 

Reaction	$E^\circ / \text{V}$
$\text{S}_2\text{O}_8^{2-} + 2\text{e}^- \rightarrow 2\text{SO}_4^{2-}$	+2.01
$\text{Co}^{3+}(\text{aq}) + \text{e}^- \rightarrow \text{Co}^{2+}(\text{aq})$	+1.82
$\text{Ce}^{4+}(\text{aq}) + \text{e}^- \rightarrow \text{Ce}^{3+}(\text{aq})$	+1.72
$\text{Au}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Au}(\text{s})$	+1.50
$\text{Cl}_2 + 2\text{e}^- \rightarrow 2\text{Cl}^-(\text{aq})$	+1.36
$\text{O}_2 + 4\text{H}^+(\text{aq}) + 4\text{e}^- \rightarrow 2\text{H}_2\text{O}$	+1.23
$\text{Br}_2 + 2\text{e}^- \rightarrow 2\text{Br}^-(\text{aq})$	+1.10
$\text{MnO}_2(\text{s}) + 4\text{H}^+(\text{aq}) + \text{e}^- \rightarrow \text{Mn}^{3+} + 2\text{H}_2\text{O}$	+0.96
$\text{Pd}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Pd}(\text{s})$	+0.92
$\text{Ag}^+(\text{aq}) + \text{e}^- \rightarrow \text{Ag}(\text{s})$	+0.80
$\text{Fe}^{3+}(\text{aq}) + \text{e}^- \rightarrow \text{Fe}^{2+}(\text{aq})$	+0.77
$\text{Cu}^+(\text{aq}) + \text{e}^- \rightarrow \text{Cu}(\text{s})$	+0.53
$\text{Cu}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Cu}(\text{s})$	+0.34
$\text{Sn}^{4+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Sn}^{2+}(\text{aq})$	+0.15
$2\text{H}^+(\text{aq}) + 2\text{e}^- \rightarrow \text{H}_2(\text{g})$	0 (by definition)
$\text{Fe}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Fe}(\text{s})$	-0.04
$\text{Pb}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Pb}(\text{s})$	-0.13
$\text{Sn}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Sn}(\text{s})$	-0.14
$\text{Ni}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Ni}(\text{s})$	-0.24
$\text{Co}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Co}(\text{s})$	-0.28
$\text{Fe}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Fe}(\text{s})$	-0.44
$\text{Cr}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Cr}(\text{s})$	-0.74
$\text{Zn}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Zn}(\text{s})$	-0.76
$2\text{H}_2\text{O} + 2\text{e}^- \rightarrow \text{H}_2(\text{g}) + 2\text{OH}^-(\text{aq})$	-0.83
$\text{Cr}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Cr}(\text{s})$	-0.89
$\text{Al}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Al}(\text{s})$	-1.68
$\text{Mg}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Mg}(\text{s})$	-2.36
$\text{Na}^+(\text{aq}) + \text{e}^- \rightarrow \text{Na}(\text{s})$	-2.71
$\text{Ca}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Ca}(\text{s})$	-2.87
$\text{Li}^+(\text{aq}) + \text{e}^- \rightarrow \text{Li}(\text{s})$	-3.04

## CHEM1102 - CHEMISTRY 1B

## Useful formulas

<p><b>Thermodynamics &amp; Equilibrium</b></p> $\Delta U = q + w = q - p\Delta V$ $\Delta_{\text{universe}}S = \Delta_{\text{sys}}S - \frac{\Delta_{\text{sys}}H}{T_{\text{sys}}}$ $\Delta G^{\circ} = \Delta H^{\circ} - T\Delta S^{\circ}$ $\Delta G = \Delta G^{\circ} + RT \ln Q$ $\Delta G^{\circ} = -RT \ln K$ $K_p = K_c (RT)^{\Delta n}$	<p><b>Electrochemistry</b></p> $\Delta G^{\circ} = -nFE^{\circ}$ <p>Moles of <math>e^{-} = It/F</math></p> $E = E^{\circ} - (RT/nF) \times 2.303 \log Q$ $= E^{\circ} - (RT/nF) \times \ln Q$ $E^{\circ} = (RT/nF) \times 2.303 \log K$ $= (RT/nF) \times \ln K$ $E = E^{\circ} - \frac{0.0592}{n} \log Q \text{ (at } 25^{\circ}\text{C)}$
<p><b>Colligative properties</b></p> $\pi = cRT$ $P_{\text{solution}} = X_{\text{solvent}} \times P^{\circ}_{\text{solvent}}$ $p = kc$ $\Delta T_f = K_f m$ $\Delta T_b = K_b m$	<p><b>Quantum Chemistry</b></p> $E = h\nu = hc/\lambda$ $\lambda = h/mv$ $4.5k_B T = hc/\lambda$ $E = -Z^2 E_R (1/n^2)$ $\Delta x \cdot \Delta(mv) \geq h/4\pi$ $q = 4\pi r^2 \times 5.67 \times 10^{-8} \times T^4$
<p><b>Acids and Bases</b></p> $pK_w = \text{pH} + \text{pOH} = 14.00$ $pK_w = \text{p}K_a + \text{p}K_b = 14.00$ $\text{pH} = \text{p}K_a + \log\{[A^{-}] / [\text{HA}]\}$	<p><b>Gas Laws</b></p> $PV = nRT$ $(P + n^2a/V^2)(V - nb) = nRT$
<p><b>Radioactivity</b></p> $t_{1/2} = \ln 2 / \lambda$ $A = \lambda N$ $\ln(N_0/N_t) = \lambda t$ $^{14}\text{C age} = 8033 \ln(A_0/A_t)$	<p><b>Kinetics</b></p> $t_{1/2} = \ln 2 / k$ $k = Ae^{-E_a/RT}$ $\ln[A] = \ln[A]_0 - kt$ $\ln \frac{k_2}{k_1} = \frac{E_a}{R} \left( \frac{1}{T_1} - \frac{1}{T_2} \right)$
<p><b>Miscellaneous</b></p> $A = -\log_{10} \frac{I}{I_0}$ $A = \epsilon cl$ $E = -A \frac{e^2}{4\pi\epsilon_0 r} N_A$	<p><b>Mathematics</b></p> <p>If <math>ax^2 + bx + c = 0</math>, then <math>x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}</math></p> $\ln x = 2.303 \log x$

# PERIODIC TABLE OF THE ELEMENTS

<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>6</b>	<b>7</b>	<b>8</b>	<b>9</b>	<b>10</b>	<b>11</b>	<b>12</b>	<b>13</b>	<b>14</b>	<b>15</b>	<b>16</b>	<b>17</b>	<b>18</b>
1 HYDROGEN <b>H</b> 1.008																	2 HELIUM <b>He</b> 4.003
3 LITHIUM <b>Li</b> 6.941	4 BERYLLIUM <b>Be</b> 9.012											5 BORON <b>B</b> 10.81	6 CARBON <b>C</b> 12.01	7 NITROGEN <b>N</b> 14.01	8 OXYGEN <b>O</b> 16.00	9 FLUORINE <b>F</b> 19.00	10 NEON <b>Ne</b> 20.18
11 SODIUM <b>Na</b> 22.99	12 MAGNESIUM <b>Mg</b> 24.31											13 ALUMINIUM <b>Al</b> 26.98	14 SILICON <b>Si</b> 28.09	15 PHOSPHORUS <b>P</b> 30.97	16 SULFUR <b>S</b> 32.07	17 CHLORINE <b>Cl</b> 35.45	18 ARGON <b>Ar</b> 39.95
19 POTASSIUM <b>K</b> 39.10	20 CALCIUM <b>Ca</b> 40.08	21 SCANDIUM <b>Sc</b> 44.96	22 TITANIUM <b>Ti</b> 47.88	23 VANADIUM <b>V</b> 50.94	24 CHROMIUM <b>Cr</b> 52.00	25 MANGANESE <b>Mn</b> 54.94	26 IRON <b>Fe</b> 55.85	27 COBALT <b>Co</b> 58.93	28 NICKEL <b>Ni</b> 58.69	29 COPPER <b>Cu</b> 63.55	30 ZINC <b>Zn</b> 65.39	31 GALLIUM <b>Ga</b> 69.72	32 GERMANIUM <b>Ge</b> 72.59	33 ARSENIC <b>As</b> 74.92	34 SELENIUM <b>Se</b> 78.96	35 BROMINE <b>Br</b> 79.90	36 KRYPTON <b>Kr</b> 83.80
37 RUBIDIUM <b>Rb</b> 85.47	38 STRONTIUM <b>Sr</b> 87.62	39 YTRIUM <b>Y</b> 88.91	40 ZIRCONIUM <b>Zr</b> 91.22	41 NIوبيUM <b>Nb</b> 92.91	42 MOLYBDENUM <b>Mo</b> 95.94	43 TECHNETIUM <b>Tc</b> [98.91]	44 RUTHENIUM <b>Ru</b> 101.07	45 RHODIUM <b>Rh</b> 102.91	46 PALLADIUM <b>Pd</b> 106.4	47 SILVER <b>Ag</b> 107.87	48 CADMIUM <b>Cd</b> 112.40	49 INDIUM <b>In</b> 114.82	50 TIN <b>Sn</b> 118.69	51 ANTIMONY <b>Sb</b> 121.75	52 TELLURIUM <b>Te</b> 127.60	53 IODINE <b>I</b> 126.90	54 XENON <b>Xe</b> 131.30
55 CAESIUM <b>Cs</b> 132.91	56 BARIUM <b>Ba</b> 137.34	57-71	72 HAFNIUM <b>Hf</b> 178.49	73 TANTALUM <b>Ta</b> 180.95	74 TUNGSTEN <b>W</b> 183.85	75 RHENIUM <b>Re</b> 186.2	76 OSMIUM <b>Os</b> 190.2	77 IRIDIUM <b>Ir</b> 192.22	78 PLATINUM <b>Pt</b> 195.09	79 GOLD <b>Au</b> 196.97	80 MERCURY <b>Hg</b> 200.59	81 THALLIUM <b>Tl</b> 204.37	82 LEAD <b>Pb</b> 207.2	83 BISMUTH <b>Bi</b> 208.98	84 POLONIUM <b>Po</b> [210.0]	85 ASTATINE <b>At</b> [210.0]	86 RADON <b>Rn</b> [222.0]
87 FRANCIUM <b>Fr</b> [223.0]	88 RADIUM <b>Ra</b> [226.0]	89-103	104 RUTHERFORDIUM <b>Rf</b> [261]	105 DUBNIUM <b>Db</b> [262]	106 SEABORGIUM <b>Sg</b> [266]	107 BOHRIUM <b>Bh</b> [262]	108 HASSIUM <b>Hs</b> [265]	109 MEITNERIUM <b>Mt</b> [266]	110 DARMSTADIUM <b>Ds</b> [271]	111 ROENTGENIUM <b>Rg</b> [272]							

	<b>57</b>	<b>58</b>	<b>59</b>	<b>60</b>	<b>61</b>	<b>62</b>	<b>63</b>	<b>64</b>	<b>65</b>	<b>66</b>	<b>67</b>	<b>68</b>	<b>69</b>	<b>70</b>	<b>71</b>
LANTHANOID S	LANTHANUM <b>La</b> 138.91	CERIUM <b>Ce</b> 140.12	PRASEODYMIUM <b>Pr</b> 140.91	NEODYMIUM <b>Nd</b> 144.24	PROMETHIUM <b>Pm</b> [144.9]	SAMARIUM <b>Sm</b> 150.4	EUROPIUM <b>Eu</b> 151.96	GADOLINIUM <b>Gd</b> 157.25	TERBIUM <b>Tb</b> 158.93	DYSPROSIUM <b>Dy</b> 162.50	HOLMIUM <b>Ho</b> 164.93	ERBIUM <b>Er</b> 167.26	THULIUM <b>Tm</b> 168.93	YTTERBIUM <b>Yb</b> 173.04	LUTETIUM <b>Lu</b> 174.97
ACTINOIDS	89 ACTINIUM <b>Ac</b> [227.0]	90 THORIUM <b>Th</b> 232.04	91 PROTACTINIUM <b>Pa</b> [231.0]	92 URANIUM <b>U</b> 238.03	93 NEPTUNIUM <b>Np</b> [237.0]	94 PLUTONIUM <b>Pu</b> [239.1]	95 AMERICIUM <b>Am</b> [243.1]	96 CURIUM <b>Cm</b> [247.1]	97 BERKELLIUM <b>Bk</b> [247.1]	98 CALIFORNIUM <b>Cf</b> [252.1]	99 EINSTEINIUM <b>Es</b> [252.1]	100 FERMIUM <b>Fm</b> [257.1]	101 MENDELEVIUM <b>Md</b> [256.1]	102 NOBELIUM <b>No</b> [259.1]	103 LAWRENCIUM <b>Lr</b> [260.1]