

Confidential

THE UNIVERSITY OF
SYDNEY

SEAT NUMBER:

STUDENT ID:

SURNAME:

GIVEN NAMES:

**CHEM1102
Chemistry 1B**

**Final Examination
Semester 2, 2014**

Time Allowed: Three hours + 10 minutes reading time

This examination paper consists of 24 pages

INSTRUCTIONS TO CANDIDATES


- This is a closed book exam.
- A simple calculator (programmable versions and PDA's not allowed) may be taken into the exam room.

Make	Model

- The total score for this paper is 100. The possible score per page is shown in the adjacent table.
- The paper comprises 30 multiple choice questions and 11 pages of short answer questions.
ANSWER ALL QUESTIONS.
- Follow the instructions on page 2 to record your answers to the multiple choice questions. Use a dark lead pencil so that you can erase errors made on the computer sheet.
- Answer all short answer questions in the spaces provided on this question paper. Credit may not be given where there is insufficient evidence of the working required to obtain the solution.
- Take care to write legibly. Write your final answers in ink, not pencil.
- Numerical values required for any question, standard electrode reduction potentials, a Periodic Table and some useful formulas may be found on the separate data sheet.

Page(s)	Marks		Marker
	Max	Gained	
2-9	30		MCG
10	5		
11	5		
12	6		
13	5		
15	9		
16	7		
17	8		
18	6		
20	7		
21	4		
23	8		
Total	100		
Check Total			

- The cubic form of boron nitride (borazon) is the second-hardest material after diamond and it crystallizes with the structure shown below. The large spheres represent the nitrogen atoms and the smaller spheres represent boron atoms.



From the unit-cell shown above, determine the empirical formula of boron nitride.

Answer:


Determine the oxidation state of the boron atoms.

Answer:

The cubic form of boron nitride is more thermally stable in air than diamond. Provide a reasonable explanation for this observation.

- A simplified phase diagram for iron is shown below, with the solid part divided into the body-centred cubic (BCC) and face-centred cubic (FCC) phases.

Marks
5



Which form of iron is stable at room temperature and pressure?

If molten iron is cooled slowly to around 1200 °C and then cooled rapidly to room temperature, the FCC form is obtained. Draw arrows on the phase diagram to indicate this process and explain why it leads to the FCC form as a metastable phase.

The line dividing the BCC and FCC forms is almost, but not quite vertical. Predict which way this line slopes and explain your answer.

- Solution A consists of a 0.050 M aqueous solution of benzoic acid, C_6H_5COOH , at 25 °C. Calculate the pH of Solution A. The pK_a of benzoic acid is 4.20.

Marks
6

pH =

Other than water, what are the major species present in solution A?

Solution B consists of a 0.050 M aqueous solution of ammonia, NH_3 , at 25 °C. Calculate the pH of Solution B. The pK_a of NH_4^+ is 9.24.

pH =

Other than water, what are the major species present in solution B?

THIS QUESTION CONTINUES ON THE NEXT PAGE.

Page Total:

Write the equation for the reaction that occurs when benzoic acid reacts with ammonia?

Marks
5

Write the expression for the equilibrium constant for the reaction of benzoic acid with ammonia?

What is the value of the equilibrium constant for the reaction of benzoic acid with ammonia?

Answer:

What are the major species in the solution that results from dissolving equimolar amounts of benzoic acid and ammonia in water?

THE REMAINDER OF THIS PAGE IS FOR ROUGH WORKING ONLY.

- The salt calcium oxalate, $\text{CaC}_2\text{O}_4 \cdot \text{H}_2\text{O}$, is sparingly soluble. Write down the chemical equation for its dissolution in water and the expression for K_{sp} .

Marks
9

What is the molar solubility of calcium oxalate? $K_{\text{sp}} = 2.3 \times 10^{-9}$

Answer:


If additional calcium oxalate is added to a saturated solution, what is the effect on $[\text{Ca}^{2+}(\text{aq})]$?

Following blood donation, a solution of sodium oxalate is added to remove $\text{Ca}^{2+}(\text{aq})$ ions which cause the blood to clot. The concentration of $\text{Ca}^{2+}(\text{aq})$ ions in blood is $9.7 \times 10^{-5} \text{ g mL}^{-1}$. If 100.0 mL of 0.1550 M $\text{Na}_2\text{C}_2\text{O}_4$ is added to 100.0 mL of blood, what will be the concentration (in mol L^{-1}) of Ca^{2+} ions remaining in the blood?

Answer:

Page Total:

- The structure below represents the active site in carbonic anhydrase, which features a Zn^{2+} ion bonded to 3 histidine residues and a water molecule.



The $\text{p}K_a$ of uncoordinated water is 15.7, but the $\text{p}K_a$ of the water ligand in carbonic anhydrase is around 7. Suggest an explanation for this large change.

When studying zinc-containing metalloenzymes, chemists often replace Zn^{2+} with Co^{2+} . Using the box notation to represent atomic orbitals, work out how many unpaired electrons are present in the Zn^{2+} and Co^{2+} ions.











Suggest why it is useful to replace Zn^{2+} with Co^{2+} when studying the nature of the active site in carbonic anhydrase.

Suggest two differences in the chemistry of Zn^{2+} and Co^{2+} ions that may affect the reactivity of the cobalt-containing enzyme.

Marks
7

- Complete the following table.

Marks
8

STARTING MATERIAL	REAGENTS/CONDITIONS	THE MAJOR ORGANIC PRODUCT(S)
	1. NaBH ₄ 2. H ⁺ / H ₂ O	
	hot concentrated H ₂ SO ₄	
		
		
	dilute aqueous H ₂ SO ₄	
	Cr ₂ O ₇ ²⁻ / H ⁺	
	2 equivalents of Cl ₂	
	SOCl ₂	




- Draw the structure of (*S*)-pent-4-en-2-ol.

Marks
3

When (*S*)-pent-4-en-2-ol reacts with bromine, Br₂, two stereoisomers are formed. Draw the structure of both products.

- Draw the structure of the organic product(s) formed when each of the following compounds is treated with 4 M sodium hydroxide. The first reaction requires heating.


3

Compound	Organic products
	
	
	

- The elimination of H₂O from alcohol **A** can form the isomeric alkenes **B** and **C**. Elimination of HBr from the alkyl halide **D** can generate the same two alkenes.

Marks

7



Assign the absolute configuration of alcohol **A**. Show your working.

Name compound **B** fully.

A diastereoisomer of **B** is also formed in these reactions. Draw the enantiomer of **A** and the diastereoisomer of **B**.

enantiomer of A	diastereoisomer of B

Propose a mechanism for the formation of **B** from **A** under the conditions shown. Use curly arrows and draw the structures of any intermediates.

THIS QUESTION CONTINUES ON THE NEXT PAGE.

Page Total:

Explain why compound **C** is the minor product of this reaction.

Marks
4

Propose a mechanism for the formation of **C** from **D** under the conditions shown. Use curly arrows and draw the structures of any intermediates.


Compound **C** is the major product formed from **D** under these conditions. What would be the major product if the enantiomer of **D** were exposed to the same reaction conditions?

THE REMAINDER OF THIS PAGE IS FOR ROUGH WORKING ONLY.

Page Total:

- Propene can be converted into 1,2-dimethyl-1-phenylpropene using a sequence of 6 reactions. Demonstrate your knowledge of Grignard reactions by suggesting a plausible sequence. Make sure you draw the correct structure for each intermediate product and clearly indicate the reagent(s) required for each reaction. The following list of suggested reagents is sufficient to accomplish all necessary reactions, but you may use other reagents if you wish. One of the intermediates is shown for you.

Suggested reagents:



HBr

dilute HCl

 $K_2Cr_2O_7 / H^+$

conc. HCl

Mg

 CH_3MgBr Marks
8

↓ reagent(s)

 product

↓ reagent(s)

 product

↓ reagent(s)

 product

↓ reagent(s)

 product

→ reagent(s)

↑ reagent(s)

 product

↑ reagent(s)

 product

THIS PAGE IS FOR ROUGH WORKING ONLY.

DATA SHEET*Physical constants*Avogadro constant, $N_A = 6.022 \times 10^{23} \text{ mol}^{-1}$ Faraday constant, $F = 96485 \text{ C mol}^{-1}$ Planck constant, $h = 6.626 \times 10^{-34} \text{ J s}$ Speed of light in vacuum, $c = 2.998 \times 10^8 \text{ m s}^{-1}$ Rydberg constant, $E_R = 2.18 \times 10^{-18} \text{ J}$ Boltzmann constant, $k_B = 1.381 \times 10^{-23} \text{ J K}^{-1}$ Permittivity of a vacuum, $\epsilon_0 = 8.854 \times 10^{-12} \text{ C}^2 \text{ J}^{-1} \text{ m}^{-1}$ Gas constant, $R = 8.314 \text{ J K}^{-1} \text{ mol}^{-1}$
 $= 0.08206 \text{ L atm K}^{-1} \text{ mol}^{-1}$ Charge of electron, $e = 1.602 \times 10^{-19} \text{ C}$ Mass of electron, $m_e = 9.1094 \times 10^{-31} \text{ kg}$ Mass of proton, $m_p = 1.6726 \times 10^{-27} \text{ kg}$ Mass of neutron, $m_n = 1.6749 \times 10^{-27} \text{ kg}$ *Properties of matter*

Volume of 1 mole of ideal gas at 1 atm and 25 °C = 24.5 L

Volume of 1 mole of ideal gas at 1 atm and 0 °C = 22.4 L

Density of water at 298 K = 0.997 g cm⁻³*Conversion factors*

1 atm = 760 mmHg = 101.3 kPa

0 °C = 273 K

1 L = 10⁻³ m³1 Å = 10⁻¹⁰ m1 eV = 1.602 × 10⁻¹⁹ J1 Ci = 3.70 × 10¹⁰ Bq1 Hz = 1 s⁻¹1 tonne = 10³ kg1 W = 1 J s⁻¹*Decimal fractions*

Fraction	Prefix	Symbol
10 ⁻³	milli	m
10 ⁻⁶	micro	μ
10 ⁻⁹	nano	n
10 ⁻¹²	pico	p

Decimal multiples

Multiple	Prefix	Symbol
10 ³	kilo	k
10 ⁶	mega	M
10 ⁹	giga	G
10 ¹²	tera	T

Standard Reduction Potentials, E°

Reaction	E° / V
$\text{Co}^{3+}(\text{aq}) + \text{e}^- \rightarrow \text{Co}^{2+}(\text{aq})$	+1.82
$\text{Ce}^{4+}(\text{aq}) + \text{e}^- \rightarrow \text{Ce}^{3+}(\text{aq})$	+1.72
$\text{MnO}_4^-(\text{aq}) + 8\text{H}^+(\text{aq}) + 5\text{e}^- \rightarrow \text{Mn}^{2+}(\text{aq}) + 4\text{H}_2\text{O}$	+1.51
$\text{Au}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Au}(\text{s})$	+1.50
$\text{Cl}_2 + 2\text{e}^- \rightarrow 2\text{Cl}^-(\text{aq})$	+1.36
$\text{O}_2 + 4\text{H}^+(\text{aq}) + 4\text{e}^- \rightarrow 2\text{H}_2\text{O}$	+1.23 (+0.82 at pH = 7)
$\text{Pt}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Pt}(\text{s})$	+1.18
$\text{MnO}_2(\text{s}) + 4\text{H}^+(\text{aq}) + \text{e}^- \rightarrow \text{Mn}^{3+} + 2\text{H}_2\text{O}$	+0.96
$\text{NO}_3^-(\text{aq}) + 4\text{H}^+(\text{aq}) + 3\text{e}^- \rightarrow \text{NO}(\text{g}) + 2\text{H}_2\text{O}$	+0.96
$\text{Pd}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Pd}(\text{s})$	+0.92
$\text{Ag}^+(\text{aq}) + \text{e}^- \rightarrow \text{Ag}(\text{s})$	+0.80
$\text{Fe}^{3+}(\text{aq}) + \text{e}^- \rightarrow \text{Fe}^{2+}(\text{aq})$	+0.77
$\text{I}_2(\text{aq}) + 2\text{e}^- \rightarrow 2\text{I}^-(\text{aq})$	+0.62
$\text{Cu}^+(\text{aq}) + \text{e}^- \rightarrow \text{Cu}(\text{s})$	+0.53
$\text{Cu}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Cu}(\text{s})$	+0.34
$\text{BiO}^+(\text{aq}) + 2\text{H}^+(\text{aq}) + 3\text{e}^- \rightarrow \text{Bi}(\text{s}) + \text{H}_2\text{O}$	+0.32
$\text{Sn}^{4+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Sn}^{2+}(\text{aq})$	+0.15
$2\text{H}^+(\text{aq}) + 2\text{e}^- \rightarrow \text{H}_2(\text{g})$	0 (by definition)
$\text{Fe}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Fe}(\text{s})$	-0.04
$\text{Pb}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Pb}(\text{s})$	-0.13
$\text{Sn}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Sn}(\text{s})$	-0.14
$\text{Ni}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Ni}(\text{s})$	-0.24
$\text{Cd}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Cd}(\text{s})$	-0.40
$\text{Fe}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Fe}(\text{s})$	-0.44
$\text{Cr}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Cr}(\text{s})$	-0.74
$\text{Zn}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Zn}(\text{s})$	-0.76
$2\text{H}_2\text{O} + 2\text{e}^- \rightarrow \text{H}_2(\text{g}) + 2\text{OH}^-(\text{aq})$	-0.83 (-0.41 at pH = 7)
$\text{Cr}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Cr}(\text{s})$	-0.89
$\text{Al}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Al}(\text{s})$	-1.68
$\text{Sc}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Sc}(\text{s})$	-2.09
$\text{Mg}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Mg}(\text{s})$	-2.36
$\text{Na}^+(\text{aq}) + \text{e}^- \rightarrow \text{Na}(\text{s})$	-2.71
$\text{Ca}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Ca}(\text{s})$	-2.87
$\text{Li}^+(\text{aq}) + \text{e}^- \rightarrow \text{Li}(\text{s})$	-3.04

Useful formulas

<p>Thermodynamics & Equilibrium</p> $\Delta U = q + w = q - p\Delta V$ $\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$ $\Delta G = \Delta G^\circ + RT \ln Q$ $\Delta G^\circ = -RT \ln K$ $\Delta_{\text{univ}}S^\circ = R \ln K$ $\ln \frac{K_2}{K_1} = \frac{-\Delta H^\circ}{R} \left(\frac{1}{T_2} - \frac{1}{T_1} \right)$	<p>Electrochemistry</p> $\Delta G^\circ = -nFE^\circ$ <p>Moles of $e^- = It/F$</p> $E = E^\circ - (RT/nF) \times 2.303 \log Q$ $= E^\circ - (RT/nF) \times \ln Q$ $E^\circ = (RT/nF) \times 2.303 \log K$ $= (RT/nF) \times \ln K$ $E = E^\circ - \frac{0.0592}{n} \log Q \text{ (at } 25^\circ \text{C)}$
<p>Acids and Bases</p> $pK_w = \text{pH} + \text{pOH} = 14.00$ $pK_w = \text{p}K_a + \text{p}K_b = 14.00$ $\text{pH} = \text{p}K_a + \log \left\{ \frac{[A^-]}{[HA]} \right\}$	<p>Gas Laws</p> $PV = nRT$ $(P + n^2a/V^2)(V - nb) = nRT$ $E_k = \frac{1}{2}mv^2$
<p>Radioactivity</p> $t_{1/2} = \ln 2 / \lambda$ $A = \lambda N$ $\ln(N_0/N_t) = \lambda t$ $^{14}\text{C age} = 8033 \ln(A_0/A_t) \text{ years}$	<p>Kinetics</p> $t_{1/2} = \ln 2 / k$ $k = Ae^{-E_a/RT}$ $\ln[A] = \ln[A]_0 - kt$ $\ln \frac{k_2}{k_1} = \frac{E_a}{R} \left(\frac{1}{T_1} - \frac{1}{T_2} \right)$
<p>Mathematics</p> <p>If $ax^2 + bx + c = 0$, then $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$</p> $\ln x = 2.303 \log x$ <p>Area of circle = πr^2</p> <p>Surface area of sphere = $4\pi r^2$</p> <p>Volume of sphere = $\frac{4}{3} \pi r^3$</p>	<p>Quantum Chemistry</p> $E = h\nu = hc/\lambda$ $\lambda = h/mv$ $E = -Z^2 E_R (1/n^2)$ $\Delta x \cdot \Delta(mv) \geq h/4\pi$ $q = 4\pi r^2 \times 5.67 \times 10^{-8} \times T^4$ $T\lambda = 2.898 \times 10^6 \text{ K nm}$
<p>Miscellaneous</p> $A = -\log \frac{I}{I_0}$ $A = \epsilon cl$ $E = -A \frac{e^2}{4\pi\epsilon_0 r} N_A$	<p>Colligative Properties & Solutions</p> $\Pi = cRT$ $P_{\text{solution}} = X_{\text{solvent}} \times P^\circ_{\text{solvent}}$ $c = kp$ $\Delta T_f = K_f m$ $\Delta T_b = K_b m$

PERIODIC TABLE OF THE ELEMENTS

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	
1 HYDROGEN H 1.008	2 HELIUM He 4.003																	
3 LITHIUM Li 6.941	4 BERYLLIUM Be 9.012											5 BORON B 10.81	6 CARBON C 12.01	7 NITROGEN N 14.01	8 OXYGEN O 16.00	9 FLUORINE F 19.00	10 NEON Ne 20.18	
11 SODIUM Na 22.99	12 MAGNESIUM Mg 24.31											13 ALUMINUM Al 26.98	14 SILICON Si 28.09	15 PHOSPHORUS P 30.97	16 SULFUR S 32.07	17 CHLORINE Cl 35.45	18 ARGON Ar 39.95	
19 POTASSIUM K 39.10	20 CALCIUM Ca 40.08	21 SCANDIUM Sc 44.96	22 TITANIUM Ti 47.88	23 VANADIUM V 50.94	24 CHROMIUM Cr 52.00	25 MANGANESE Mn 54.94	26 IRON Fe 55.85	27 COBALT Co 58.93	28 NICKEL Ni 58.69	29 COPPER Cu 63.55	30 ZINC Zn 65.39	31 GALLIUM Ga 69.72	32 GERMANIUM Ge 72.59	33 ARSENIC As 74.92	34 SELENIUM Se 78.96	35 BROMINE Br 79.90	36 KRYPTON Kr 83.80	
37 BROMINE Rb 85.47	38 STRONTIUM Sr 87.62	39 YTRBIUM Y 88.91	40 ZIRCONIUM Zr 91.22	41 NIOBIUM Nb 92.91	42 MOLYBDENUM Mo 95.94	43 TECHNETIUM Tc [98.91]	44 RHENIUM Ru 101.07	45 RHODIUM Rh 102.91	46 PALLADIUM Pd 106.4	47 SILVER Ag 107.87	48 Cadmium Cd 112.40	49 INDIUM In 114.82	50 TIN Sn 118.69	51 ANTIMONY Sb 121.75	52 TELLURIUM Te 127.60	53 IODINE I 126.90	54 XENON Xe 131.30	
55 CAESIUM Cs 132.91	56 BARIUM Ba 137.34	57-71 LANTHANIDS	72 HAFNIUM Hf 178.49	73 TANTALUM Ta 180.95	74 TUNGSTEN W 183.85	75 RHENIUM Re 186.2	76 OSMIUM Os 190.2	77 IRIDIUM Ir 192.22	78 PLATINUM Pt 195.09	79 GOLD Au 196.97	80 MERCURY Hg 200.59	81 THALLIUM Tl 204.37	82 LEAD Pb 207.2	83 BISMUTH Bi 208.98	84 POLONIUM Po [210.0]	85 ASTATINE At [210.0]	86 RADON Rn [222.0]	
87 FRANCIUM Fr [223.0]	88 RADIUM Ra [226.0]		104 RIFERBERORIUM Rf [261]	105 DUBNIUM Db [262]	106 SEABERGIUM Sg [266]	107 BOHRIUM Bh [262]	108 HASSIUM Hs [265]	109 MEITNERIUM Mt [266]	110 DARSTADTIUM Ds [271]	111 ROSTERIUM Rg [272]	112 COOPERIUM Cn [283]		114 FLEROVIUM Fl [289]					

LANTHANIDS

57 LANTHANUM La 138.91	58 CEURIUM Ce 140.12	59 PRASEODYMIUM Pr 140.91	60 NEODYMIUM Nd 144.24	61 PROMETHIUM Pm [144.91]	62 SAMARIUM Sm 150.4	63 EUROPIUM Eu 151.96	64 GADOLINIUM Gd 157.25	65 TERBIUM Tb 158.93	66 DYSPROSIUM Dy 162.50	67 HOIMIUM Ho 164.93	68 ERBIUM Er 167.26	69 THULIUM Tm 168.93	70 YTERBIUM Yb 173.04	71 LUTETIUM Lu 174.97
89 ACTINIUM Ac [227.0]	90 THORIUM Th 232.04	91 PROTACTINIUM Pa [231.0]	92 URANIUM U 238.03	93 NEPTUNIUM Np [237.0]	94 PLUTONIUM Pu [239.1]	95 AMERICIUM Am [243.1]	96 CURIUM Cm [247.1]	97 BERKELIUM Bk [247.1]	98 CALIFORNIUM Cf [252.1]	99 EINSTEINIUM Es [252.1]	100 FERMIUM Fm [257.1]	101 MENDELEVIUM Md [256.1]	102 NOBELIUM No [259.1]	103 LAWRENCIUM Lr [260.1]

ACTINOIDS