

CHEMISTRY 1B - CHEM1102SECOND SEMESTER EXAMINATION**CONFIDENTIAL****NOVEMBER 2005****TIME ALLOWED: THREE HOURS**

GIVE THE FOLLOWING INFORMATION IN BLOCK LETTERS

FAMILY NAME		SID NUMBER	
OTHER NAMES		TABLE NUMBER	

INSTRUCTIONS TO CANDIDATES

- All questions are to be attempted. There are 20 pages of examinable material.
- Complete the written section of the examination paper in **INK**.
- Read each question carefully. Report the appropriate answer and show all relevant working in the space provided.
- The total score for this paper is 100. The possible score per page is shown in the adjacent tables.
- Each new question of the short answer section begins with a •.
- Electronic calculators, including programmable calculators, may be used. Students are warned, however, that credit may not be given, even for a correct answer, where there is insufficient evidence of the working required to obtain the solution.
- Numerical values required for any question, standard electrode reduction potentials, a Periodic Table and some useful formulas may be found on the separate data sheet.
- Pages 17, 22 & 24 are for rough working only.

OFFICIAL USE ONLY**Multiple choice section**

	Marks	
Pages	Max	Gained
2-11	44	

Short answer section

Page	Marks		Marker
	Max	Gained	
12	5		
13	5		
14	10		
15	3		
16	8		
18	5		
19	5		
20	2		
21	7		
23	6		
Total	56		

Marks
2

- Calcium oxalate is a major constituent of kidney stones. Calculate the solubility product constant for calcium oxalate given that a saturated solution of the salt can be made by dissolving 0.0061 g of $\text{CaC}_2\text{O}_4 \cdot \text{H}_2\text{O}(\text{s})$ in 1.0 L of water.

Answer:


3

- A sample of 2.0 mg of $\text{Cu}(\text{OH})_2$ is added to 1.0 L of a solution buffered at a pH of 8.00. Will all of the $\text{Cu}(\text{OH})_2$ dissolve? Show all working. (The K_{sp} of $\text{Cu}(\text{OH})_2$ is $4.8 \times 10^{-20} \text{ M}^3$.)

Answer:

Marks
5

- The following data have been obtained for the rapid reaction between the hypochlorite and iodide ions.



$[\text{OCl}^-(\text{aq})] / \text{M}$	$[\text{I}^-(\text{aq})] / \text{M}$	Rate / M s^{-1}
1.5×10^{-3}	1.5×10^{-3}	1.36×10^{-4}
3.0×10^{-3}	1.5×10^{-3}	2.72×10^{-4}
1.5×10^{-3}	3.0×10^{-3}	2.72×10^{-4}

What is the order of this reaction with respect to each reagent?

Calculate the rate of the reaction when $[\text{OCl}^-(\text{aq})] = 2.5 \times 10^{-3} \text{ M}$ and $[\text{I}^-(\text{aq})] = 4.0 \times 10^{-3} \text{ M}$.

Answer:

- Explain in terms of their electronic configurations **and** ionisation energies why the alkali metals (Group 1) are powerful *reducing* agents.

Marks
2

- Compounds of *d*-block elements are frequently paramagnetic. Using the box notation to represent atomic orbitals, account for this property in compounds of Ni^{2+} .

2

- Complete the following table.

6

Formula	Oxidation state of transition metal	Coordination number of transition metal	Number of <i>d</i> -electrons in metal in complex ion	Species formed upon dissolving in water
$\text{K}_3[\text{FeF}_6]$				
$[\text{Cr}(\text{NH}_3)_5(\text{H}_2\text{O})]\text{Cl}_3$				
$[\text{Zn}(\text{en})\text{Cl}_2]$				

en = ethylenediamine = $\text{NH}_2\text{CH}_2\text{CH}_2\text{NH}_2$

- Define what is meant by an “allotrope”. Give an example of a pair of allotropes involving phosphorus and a pair *not* involving phosphorus.

Marks
3

THE REMAINDER OF THIS PAGE IS FOR ROUGH WORKING ONLY.

- Solution A consists of a 0.15 M aqueous solution of HF at 25 °C. Calculate the pH of Solution A. The pK_a of HF is 3.17.

Marks
8

pH =

At 25 °C, 1.00 L of Solution B consists of 11.62 g of KF dissolved in water. Calculate the pH of Solution B.

pH =


Solution B (1.00 L) is poured into Solution A (1.00 L) and allowed to equilibrate at 25 °C. Calculate the pH of the final solution.

pH =

If you wanted to adjust the pH of the mixture of Solution A and Solution B to be exactly equal to 3.00, which component in the solution would you need to increase in concentration?

- The structure of methyl 4-hydroxybenzoate, **E**, a constituent of many suntan lotions, is given below.

Marks
5



Give the molecular formula of compound **E**.

Name the functional groups in molecule **E** indicated by the boxes “**a**” and “**b**”.






a:	b:
-----------	-----------

Give the structure(s) of all organic products formed when compound **E** is treated with the following reagents. If no reaction occurs, write “NO REACTION”.

cold NaOH (1 M)	
hot NaOH (4 M)	
hot HCl (4 M)	


Marks
5

- Draw the constitutional formula of the major organic product formed in each of the following reactions.

**THE REMAINDER OF THIS PAGE IS FOR ROUGH WORKING ONLY.**

- Show the structure(s) of all products formed when dipeptide **F** is treated with hot, concentrated HCl.


Marks
2



THE REMAINDER OF THIS PAGE IS FOR ROUGH WORKING ONLY.

- Butanone is treated first with phenylmagnesium bromide in diethyl ether and then with dilute aqueous acid, to yield alcohol **G**.

Marks
7



State whether **G** is obtained as the (*R*)-enantiomer, the (*S*)-enantiomer, a racemic mixture, or is achiral.

--

List below, the substituents on the stereogenic carbon atom in **G**, in decreasing priority (ie. from highest to lowest priority), as determined by the sequence rules.

highest priority

lowest priority

--	--	--	--

Draw the (*R*)-enantiomer of **G**, showing the correct absolute stereochemistry.

--

G is treated with concentrated sulfuric acid to give a mixture of three alkenes, **H**, **I** and **J**. Compounds **H** and **I** are diastereomers, while **H** and **J** (and **I** and **J**) are constitutional isomers. Give the structures and systematic names for **H**, **I** and **J**.


H	I	J
Name:	Name:	Name:

Give the structure of the product obtained when alkene **J** is treated with hydrogen in ethanol, in the presence of palladium on charcoal.

--


- In the space below, devise a synthesis of ethyl butyrate (ethyl butanoate), starting from 1-bromopropane. Note that your synthetic route may require more than one step. Show clearly the reagents you would use and draw constitutional formulas for any intermediate compounds.

Marks
4



- The incomplete proposed mechanism for the reaction of 2-methyl-2-butene with HBr is shown below. Complete the mechanism by adding curly arrows to illustrate the bonding changes that take place.

2



Which one of the two reactants is the electrophile?

CHEM1102 - CHEMISTRY 1B
DATA SHEET

Physical constants

Avogadro constant, $N_A = 6.022 \times 10^{23} \text{ mol}^{-1}$

Faraday constant, $F = 96485 \text{ C mol}^{-1}$

Planck constant, $h = 6.626 \times 10^{-34} \text{ J s}$

Speed of light in vacuum, $c = 2.998 \times 10^8 \text{ m s}^{-1}$

Rydberg constant, $E_R = 2.18 \times 10^{-18} \text{ J}$

Boltzmann constant, $k_B = 1.381 \times 10^{-23} \text{ J K}^{-1}$

Gas constant, $R = 8.314 \text{ J K}^{-1} \text{ mol}^{-1}$
 $= 0.08206 \text{ L atm K}^{-1} \text{ mol}^{-1}$

Charge of electron, $e = 1.602 \times 10^{-19} \text{ C}$

Mass of electron, $m_e = 9.1094 \times 10^{-31} \text{ kg}$

Mass of proton, $m_p = 1.6726 \times 10^{-27} \text{ kg}$

Mass of neutron, $m_n = 1.6749 \times 10^{-27} \text{ kg}$

Properties of matter

Volume of 1 mole of ideal gas at 1 atm and 25 °C = 24.5 L

Volume of 1 mole of ideal gas at 1 atm and 0 °C = 22.4 L

Density of water at 298 K = 0.997 g cm⁻³

Conversion factors

1 atm = 760 mmHg = 101.3 kPa

0 °C = 273 K

1 L = 10⁻³ m³

1 Å = 10⁻¹⁰ m

1 eV = 1.602 × 10⁻¹⁹ J

1 Ci = 3.70 × 10¹⁰ Bq

1 Hz = 1 s⁻¹

Decimal fractions

Fraction	Prefix	Symbol
10 ⁻³	milli	m
10 ⁻⁶	micro	μ
10 ⁻⁹	nano	n
10 ⁻¹²	pico	p

Decimal multiples

Multiple	Prefix	Symbol
10 ³	kilo	k
10 ⁶	mega	M
10 ⁹	giga	G

CHEM1102 - CHEMISTRY 1B**Standard Reduction Potentials, E°**

Reaction	E° / V
$\text{Co}^{3+}(\text{aq}) + \text{e}^- \rightarrow \text{Co}^{2+}(\text{aq})$	+1.82
$\text{Ce}^{4+}(\text{aq}) + \text{e}^- \rightarrow \text{Ce}^{3+}(\text{aq})$	+1.72
$\text{Au}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Au}(\text{s})$	+1.50
$\text{Cl}_2 + 2\text{e}^- \rightarrow 2\text{Cl}^-(\text{aq})$	+1.36
$\text{O}_2 + 4\text{H}^+(\text{aq}) + 4\text{e}^- \rightarrow 2\text{H}_2\text{O}$	+1.23
$\text{MnO}_2(\text{s}) + 4\text{H}^+(\text{aq}) + \text{e}^- \rightarrow \text{Mn}^{3+} + 2\text{H}_2\text{O}$	+0.96
$\text{Pd}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Pd}(\text{s})$	+0.92
$\text{Ag}^+(\text{aq}) + \text{e}^- \rightarrow \text{Ag}(\text{s})$	+0.80
$\text{Fe}^{3+}(\text{aq}) + \text{e}^- \rightarrow \text{Fe}^{2+}(\text{aq})$	+0.77
$\text{Cu}^+(\text{aq}) + \text{e}^- \rightarrow \text{Cu}(\text{s})$	+0.53
$\text{Cu}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Cu}(\text{s})$	+0.34
$\text{Sn}^{4+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Sn}^{2+}(\text{aq})$	+0.15
$2\text{H}^+(\text{aq}) + 2\text{e}^- \rightarrow \text{H}_2(\text{g})$	0 (by definition)
$\text{Fe}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Fe}(\text{s})$	-0.04
$\text{Pb}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Pb}(\text{s})$	-0.13
$\text{Sn}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Sn}(\text{s})$	-0.14
$\text{Ni}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Ni}(\text{s})$	-0.24
$\text{Fe}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Fe}(\text{s})$	-0.44
$\text{Cr}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Cr}(\text{s})$	-0.74
$\text{Zn}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Zn}(\text{s})$	-0.76
$2\text{H}_2\text{O} + 2\text{e}^- \rightarrow \text{H}_2(\text{g}) + 2\text{OH}^-(\text{aq})$	-0.83
$\text{Cr}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Cr}(\text{s})$	-0.89
$\text{Al}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Al}(\text{s})$	-1.68
$\text{Mg}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Mg}(\text{s})$	-2.36
$\text{Na}^+(\text{aq}) + \text{e}^- \rightarrow \text{Na}(\text{s})$	-2.71
$\text{Ca}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Ca}(\text{s})$	-2.87
$\text{Li}^+(\text{aq}) + \text{e}^- \rightarrow \text{Li}(\text{s})$	-3.04

CHEM1102 - CHEMISTRY 1B

Useful formulas

<p>Quantum Chemistry</p> $E = h\nu = hc/\lambda$ $\lambda = h/mv$ $4.5k_B T = hc/\lambda$ $E = Z^2 E_R (1/n^2)$ $\Delta x \cdot \Delta(mv) \geq h/4\pi$	<p>Radioactivity</p> $t_{1/2} = \ln 2 / \lambda$ $A = \lambda N$ $\ln(N_0/N_t) = \lambda t$ $^{14}\text{C age} = 8033 \ln(A_0/A_t)$
<p>Acids and Bases</p> $pK_w = \text{pH} + \text{pOH} = 14.00$ $pK_w = \text{pK}_a + \text{pK}_b = 14.00$ $\text{pH} = \text{pK}_a + \log\{[A^-] / [\text{HA}] \}$	<p>Gas Laws</p> $PV = nRT$ $(P + n^2 a/V^2)(V - nb) = nRT$
<p>Colligative properties</p> $\pi = cRT$ $P_{\text{solution}} = X_{\text{solvent}} \times P^{\circ}_{\text{solvent}}$ $p = kc$ $\Delta T_f = K_f m$ $\Delta T_b = K_b m$	<p>Kinetics</p> $t_{1/2} = \ln 2 / k$ $k = A e^{-E_a/RT}$ $\ln[A] = \ln[A]_0 - kt$ $\ln \frac{k_2}{k_1} = \frac{E_a}{R} \left(\frac{1}{T_1} - \frac{1}{T_2} \right)$
<p>Electrochemistry</p> $\Delta G^{\circ} = -nFE^{\circ}$ <p>Moles of $e^- = It/F$</p> $E = E^{\circ} - (RT/nF) \times 2.303 \log Q$ $= E^{\circ} - (RT/nF) \times \ln Q$ $E^{\circ} = (RT/nF) \times 2.303 \log K$ $= (RT/nF) \times \ln K$ $E = E^{\circ} - \frac{0.0592}{n} \log Q \text{ (at } 25^{\circ}\text{C)}$	<p>Thermodynamics & Equilibrium</p> $\Delta G^{\circ} = \Delta H^{\circ} - T\Delta S^{\circ}$ $\Delta G = \Delta G^{\circ} + RT \ln Q$ $\Delta G^{\circ} = -RT \ln K$ $K_p = K_c (RT)^{\Delta n}$
<p>Polymers</p> $R_g = \sqrt{\frac{nl_0^2}{6}}$	<p>Mathematics</p> <p>If $ax^2 + bx + c = 0$, then $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$</p> $\ln x = 2.303 \log x$

PERIODIC TABLE OF THE ELEMENTS

November 2005

CHEM1102 - CHEMISTRY 1B

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18		
	1 <small>HYDROGEN</small> H 1.008											2 <small>HELIUM</small> He 4.003								
	3 <small>LITHIUM</small> Li 6.941	4 <small>BERYLLIUM</small> Be 9.012										5 <small>BORON</small> B 10.81	6 <small>CARBON</small> C 12.01	7 <small>NITROGEN</small> N 14.01	8 <small>OXYGEN</small> O 16.00	9 <small>FLUORINE</small> F 19.00	10 <small>NEON</small> Ne 20.18			
	11 <small>SODIUM</small> Na 22.99	12 <small>MAGNESIUM</small> Mg 24.31										13 <small>ALUMINIUM</small> Al 26.98	14 <small>SILICON</small> Si 28.09	15 <small>PHOSPHORUS</small> P 30.97	16 <small>SULFUR</small> S 32.07	17 <small>CHLORINE</small> Cl 35.45	18 <small>ARGON</small> Ar 39.95			
	19 <small>POTASSIUM</small> K 39.10	20 <small>CALCIUM</small> Ca 40.08	21 <small>SCANDIUM</small> Sc 44.96	22 <small>TITANIUM</small> Ti 47.88	23 <small>VANADIUM</small> V 50.94	24 <small>CHROMIUM</small> Cr 52.00	25 <small>MANGANESE</small> Mn 54.94	26 <small>IRON</small> Fe 55.85	27 <small>COBALT</small> Co 58.93	28 <small>NICKEL</small> Ni 58.69	29 <small>COPPER</small> Cu 63.55	30 <small>ZINC</small> Zn 65.39	31 <small>GALLIUM</small> Ga 69.72	32 <small>GERMANIUM</small> Ge 72.59	33 <small>ARSENIC</small> As 74.92	34 <small>SELENIUM</small> Se 78.96	35 <small>BROMINE</small> Br 79.90	36 <small>KRYPTON</small> Kr 83.80		
	37 <small>RUBIDIUM</small> Rb 85.47	38 <small>STRONTIUM</small> Sr 87.62	39 <small>YTRIUM</small> Y 88.91	40 <small>ZIRCONIUM</small> Zr 91.22	41 <small>NIوبيUM</small> Nb 92.91	42 <small>MOLYBDENUM</small> Mo 95.94	43 <small>TECHNETIUM</small> Tc [98.91]	44 <small>RUTHENIUM</small> Ru 101.07	45 <small>RHODIUM</small> Rh 102.91	46 <small>PALLADIUM</small> Pd 106.4	47 <small>SILVER</small> Ag 107.87	48 <small>CADMIUM</small> Cd 112.40	49 <small>INDIUM</small> In 114.82	50 <small>TIN</small> Sn 118.69	51 <small>ANTIMONY</small> Sb 121.75	52 <small>TELLURIUM</small> Te 127.60	53 <small>IODINE</small> I 126.90	54 <small>XENON</small> Xe 131.30		
	55 <small>CAESIUM</small> Cs 132.91	56 <small>BARIUM</small> Ba 137.34	57-71		72 <small>HAFNIUM</small> Hf 178.49	73 <small>TANTALUM</small> Ta 180.95	74 <small>TUNGSTEN</small> W 183.85	75 <small>RHENIUM</small> Re 186.2	76 <small>OSMIUM</small> Os 190.2	77 <small>IRIDIUM</small> Ir 192.22	78 <small>PLATINUM</small> Pt 195.09	79 <small>GOLD</small> Au 196.97	80 <small>MERCURY</small> Hg 200.59	81 <small>THALLIUM</small> Tl 204.37	82 <small>LEAD</small> Pb 207.2	83 <small>BISMUTH</small> Bi 208.98	84 <small>POLONIUM</small> Po [210.0]	85 <small>ASTATINE</small> At [210.0]	86 <small>RADON</small> Rn [222.0]	
	87 <small>FRANCIUM</small> Fr [223.0]	88 <small>RADIUM</small> Ra [226.0]	89-103		104 <small>RUTHERFORDIUM</small> Rf [261]	105 <small>DUBNIUM</small> Db [262]	106 <small>SEABORGIUM</small> Sg [266]	107 <small>BOHRIUM</small> Bh [262]	108 <small>HASSIUM</small> Hs [265]	109 <small>MEITNERIUM</small> Mt [266]										

LANTHANIDES

	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71
	<small>LANTHANUM</small> La 138.91	<small>CERIUM</small> Ce 140.12	<small>PRASEODYMIUM</small> Pr 140.91	<small>NEODYMIUM</small> Nd 144.24	<small>PROMETHIUM</small> Pm [144.9]	<small>SAMARIUM</small> Sm 150.4	<small>EUROPIUM</small> Eu 151.96	<small>GADOLINIUM</small> Gd 157.25	<small>TERBIUM</small> Tb 158.93	<small>DYSPROSIUM</small> Dy 162.50	<small>HOLMIUM</small> Ho 164.93	<small>ERBIUM</small> Er 167.26	<small>THULIUM</small> Tm 168.93	<small>YTTERBIUM</small> Yb 173.04	<small>LUTETIUM</small> Lu 174.97
	<small>ACTINIUM</small> Ac [227.0]	<small>THORIUM</small> Th 232.04	<small>PROTACTINIUM</small> Pa [231.0]	<small>URANIUM</small> U 238.03	<small>NEPTUNIUM</small> Np [237.0]	<small>PLUTONIUM</small> Pu [239.1]	<small>AMERICIUM</small> Am [243.1]	<small>CURIUM</small> Cm [247.1]	<small>BERKELIUM</small> Bk [247.1]	<small>CALIFORNIUM</small> Cf [252.1]	<small>EINSTEINIUM</small> Es [252.1]	<small>FERMIUM</small> Fm [257.1]	<small>MENDELEVIUM</small> Md [256.1]	<small>NOBELIUM</small> No [259.1]	<small>LAWRENCIUM</small> Lr [260.1]

ACTINIDES

22/10(b)