

## CHEM1108 - CHEMISTRY 1A LIFE SCIENCES

### FIRST SEMESTER EXAMINATION

**CONFIDENTIAL**

**JUNE 2009**

**TIME ALLOWED: THREE HOURS**

GIVE THE FOLLOWING INFORMATION IN BLOCK LETTERS

<b>FAMILY NAME</b>		<b>SID NUMBER</b>	
<b>OTHER NAMES</b>		<b>TABLE NUMBER</b>	

### INSTRUCTIONS TO CANDIDATES

- All questions are to be attempted. There are 20 pages of examinable material.
- Complete the examination paper in **INK**.
- Read each question carefully. Report the appropriate answer and show all relevant working in the space provided.
- The total score for this paper is 100. The possible score per page is shown in the adjacent tables.
- Each new short answer question begins with a •.
- Electronic calculators, including programmable calculators, may be used. Students are warned, however, that credit may not be given, even for a correct answer, where there is insufficient evidence of the working required to obtain the solution. Logarithms may also be used.
- Numerical values required for any question as well as a Periodic Table are printed on a separate data sheet.
- Pages 16, 22 and 24 are for rough work only.

### OFFICIAL USE ONLY

#### **Multiple choice section**

Pages	Marks	
	Max	Gained
2-11	30	

#### **Short answer section**

Page	Marks		Marker
	Max	Gained	
12	9		
13	8		
14	4		
15	3		
17	11		
18	4		
19	7		
20	6		
21	8		
23	10		
Total	70		
Check Total			

**Marks**  
**2**

- Complete the following table, giving either the systematic name or the molecular formula as required.

Formula	Systematic name
NaHSO <sub>4</sub>	
	arsenic(III) chloride
CrCl <sub>3</sub> ·6H <sub>2</sub> O	
	silver dichromate

**3**

- Complete the following table, providing the ground state electron configuration for each of the following species.

Species	Ground state electron configuration
chlorine atom	
magnesium ion	
arsenic(V) ion	

**4**

- Like most medicines, the platinum complex, cisplatin, *cis*-[PtCl<sub>2</sub>(NH<sub>3</sub>)<sub>2</sub>], is both effective and toxic. What is cisplatin used to treat?

--

What does the cisplatin react with in the body to cause most of the toxicity?


--

Draw a graph showing the relationship between overall health and the level of platinum in the body of a healthy person.

--

**Marks**  
**8**

- The molecular structure of adrenaline (epinephrine), a hormone involved in the "fight or flight" response, is shown below.



List the types of intermolecular interactions that each of the following sites on adrenaline would be involved in if dissolved in water.

**A**

**B**

**C**

Pharmaceuticals with amine groups are frequently supplied as their "hydrochloride salts". Draw the structure that would result if adrenaline were reacted with one equivalent of HCl. What **additional** intermolecular forces would be present if this form of adrenaline were dissolved in water?

Provide the requested information for each of the indicated sites on adrenaline.

Atom	Geometric arrangement of the electron pairs around the atom	Hybridisation of the atom	Geometry around the atom	Approximate angles around the atom
#1O				
#2N				
#3C				


- Cadmium chloride and cadmium sulfate are both soluble in water. Describe, using equations where appropriate, how to convert cadmium chloride into cadmium sulfate.

**Marks**  
**4**

**THE REMAINDER OF THIS PAGE IS FOR ROUGH WORKING ONLY.**

**Marks**  
**3**

- The diagram below shows the general trend for the first ionisation energy for some *s* and *p* block elements.











How will the general trend differ for the second ionisation energy of these elements (*i.e.*  $X^+(g) \rightarrow X^{2+}(g) + e^-$ )? Explain.

**THE REMAINDER OF THIS PAGE IS FOR ROUGH WORKING ONLY.**


**Marks**  
**11**

- Complete the following table. Make sure you complete the name of the starting material or major product where indicated.

STARTING MATERIAL	REAGENTS/ CONDITIONS	CONSTITUTIONAL FORMULA(S) OF MAJOR ORGANIC PRODUCT(S)
	HBr / CCl <sub>4</sub> (solvent)	
		 <b>Name:</b>
$\text{CH}_3\text{CH}_2\underset{\text{Br}}{\text{CH}}\text{CH}_2\text{CH}_3$ <b>Name:</b>		$\text{CH}_3\text{CH}_2\underset{\text{Br}^\ominus \text{N}^\oplus(\text{CH}_3)_3}{\text{CH}}\text{CH}_2\text{CH}_3$
	$[\text{Ag}(\text{NH}_3)_2]^\oplus / \text{OH}^\ominus$	
	3 M NaOH / heat	
 <b>Name:</b>		
	hot conc. KOH in ethanol solvent	

**Marks**  
**4**

- Acyclovir is an analogue of the nucleoside guanosine, and is used clinically as an antiviral agent.




Hydrolysis of acyclovir gives the nucleic base guanine, a diol and a carbonyl compound. Give the structures of guanine, a tautomer of guanine, and the diol and carbonyl compounds formed.

guanine	tautomer of guanine
the diol	the carbonyl compound

**THE REMAINDER OF THIS PAGE IS FOR ROUGH WORKING ONLY.**

**Marks**  
**7**

- Morphine is the principal active agent in opium and is a highly potent analgesic drug. Its structure and conversion into codeine (a moderate analgesic) and pholcodine (a cough suppressant) are shown below.



Give the molecular formula of morphine.

How many stereogenic (chiral) centres are there in morphine?

Identify the functional groups present in morphine.

Draw the structures of codeine and reagent B.


codeine

reagent B




**Marks**  
**6**

- Show clearly the reagents you would use to carry out the following chemical conversions. Note that more than one step is required and you should indicate all necessary steps and the constitutional formulas of any intermediate compounds.




- Shown below are the Haworth structure of  $\beta$ -D-mannopyranose and the Fischer projection of D-galactose.

**Marks**  
**8**



$\beta$ -D-mannopyranose



D-galactose

Draw structures for the following sugars.

Fischer projection of D-mannose

Haworth structure of  $\alpha$ -D-galactopyranose


Give the product(s) obtained when D-mannose is treated with acidified methanol.

Draw the structure of any non-reducing disaccharide formed from D-mannose and D-galactose, indicating the configurations at the anomeric carbon atoms.

How many different non-reducing disaccharides can be formed from D-mannose and D-galactose? What is the relationship between any two of these compounds?

**Marks**  
**10**

- Glutathione is an important tripeptide (Glu-Cys-Gly) which acts as an antioxidant, protecting cells from toxins such as free radicals. It is an unusual peptide in that the peptidic linkage with glutamic acid (Glu) involves the carboxylic acid group in the side chain.



Give the product when glutathione undergoes oxidation.

--

Draw the Fischer projections of the three amino acids (in their natural absolute configurations, where applicable) that result from the vigorous acid hydrolysis (with 6 M HCl) of glutathione.

--	--	--

Draw the major species present when glutamic acid (Glu) is dissolved in water at pH 1 and pH 12. The  $pK_a$  values of glutamic acid are 2.1 ( $\alpha$ -COOH), 9.5 ( $\alpha$ -NH<sub>3</sub><sup>⊕</sup>) and 4.0 (side chain).

pH 1	pH 12
------	-------

Give the constitutional formula for the dipeptide Cys-Gly in its zwitterionic state.

--

**CHEM1108 - CHEMISTRY 1A LIFE SCIENCES****DATA SHEET***Physical constants*Avogadro constant,  $N_A = 6.022 \times 10^{23} \text{ mol}^{-1}$ Faraday constant,  $F = 96485 \text{ C mol}^{-1}$ Planck constant,  $h = 6.626 \times 10^{-34} \text{ J s}$ Speed of light in vacuum,  $c = 2.998 \times 10^8 \text{ m s}^{-1}$ Rydberg constant,  $E_R = 2.18 \times 10^{-18} \text{ J}$ Boltzmann constant,  $k_B = 1.381 \times 10^{-23} \text{ J K}^{-1}$ Permittivity of a vacuum,  $\epsilon_0 = 8.854 \times 10^{-12} \text{ C}^2 \text{ J}^{-1} \text{ m}^{-1}$ Gas constant,  $R = 8.314 \text{ J K}^{-1} \text{ mol}^{-1}$   
 $= 0.08206 \text{ L atm K}^{-1} \text{ mol}^{-1}$ Charge of electron,  $e = 1.602 \times 10^{-19} \text{ C}$ Mass of electron,  $m_e = 9.1094 \times 10^{-31} \text{ kg}$ Mass of proton,  $m_p = 1.6726 \times 10^{-27} \text{ kg}$ Mass of neutron,  $m_n = 1.6749 \times 10^{-27} \text{ kg}$ *Properties of matter*

Volume of 1 mole of ideal gas at 1 atm and 25 °C = 24.5 L

Volume of 1 mole of ideal gas at 1 atm and 0 °C = 22.4 L

Density of water at 298 K = 0.997 g cm<sup>-3</sup>*Conversion factors*

1 atm = 760 mmHg = 101.3 kPa

1 Ci =  $3.70 \times 10^{10}$  Bq

0 °C = 273 K

1 Hz = 1 s<sup>-1</sup>1 L = 10<sup>-3</sup> m<sup>3</sup>1 tonne = 10<sup>3</sup> kg1 Å = 10<sup>-10</sup> m1 W = 1 J s<sup>-1</sup>1 eV =  $1.602 \times 10^{-19}$  J*Decimal fractions*

Fraction	Prefix	Symbol
10 <sup>-3</sup>	milli	m
10 <sup>-6</sup>	micro	μ
10 <sup>-9</sup>	nano	n
10 <sup>-12</sup>	pico	p

*Decimal multiples*

Multiple	Prefix	Symbol
10 <sup>3</sup>	kilo	k
10 <sup>6</sup>	mega	M
10 <sup>9</sup>	giga	G

**CHEM1108 - CHEMISTRY 1A LIFE SCIENCES***Standard Reduction Potentials, E°*

Reaction	$E^\circ / \text{V}$
$\text{Co}^{3+}(\text{aq}) + \text{e}^- \rightarrow \text{Co}^{2+}(\text{aq})$	+1.82
$\text{Ce}^{4+}(\text{aq}) + \text{e}^- \rightarrow \text{Ce}^{3+}(\text{aq})$	+1.72
$\text{MnO}_4^-(\text{aq}) + 8\text{H}^+(\text{aq}) + 5\text{e}^- \rightarrow \text{Mn}^{2+}(\text{aq}) + 4\text{H}_2\text{O}$	+1.51
$\text{Au}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Au}(\text{s})$	+1.50
$\text{Cl}_2 + 2\text{e}^- \rightarrow 2\text{Cl}^-(\text{aq})$	+1.36
$\text{O}_2 + 4\text{H}^+(\text{aq}) + 4\text{e}^- \rightarrow 2\text{H}_2\text{O}$	+1.23
$\text{Pt}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Pt}(\text{s})$	+1.18
$\text{MnO}_2(\text{s}) + 4\text{H}^+(\text{aq}) + \text{e}^- \rightarrow \text{Mn}^{3+} + 2\text{H}_2\text{O}$	+0.96
$\text{NO}_3^-(\text{aq}) + 4\text{H}^+(\text{aq}) + 3\text{e}^- \rightarrow \text{NO}(\text{g}) + 2\text{H}_2\text{O}$	+0.96
$\text{Pd}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Pd}(\text{s})$	+0.92
$\text{Ag}^+(\text{aq}) + \text{e}^- \rightarrow \text{Ag}(\text{s})$	+0.80
$\text{Fe}^{3+}(\text{aq}) + \text{e}^- \rightarrow \text{Fe}^{2+}(\text{aq})$	+0.77
$\text{Cu}^+(\text{aq}) + \text{e}^- \rightarrow \text{Cu}(\text{s})$	+0.53
$\text{Cu}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Cu}(\text{s})$	+0.34
$\text{Sn}^{4+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Sn}^{2+}(\text{aq})$	+0.15
$2\text{H}^+(\text{aq}) + 2\text{e}^- \rightarrow \text{H}_2(\text{g})$	0 (by definition)
$\text{Fe}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Fe}(\text{s})$	-0.04
$\text{Pb}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Pb}(\text{s})$	-0.13
$\text{Sn}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Sn}(\text{s})$	-0.14
$\text{Ni}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Ni}(\text{s})$	-0.24
$\text{Cd}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Cd}(\text{s})$	-0.40
$\text{Fe}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Fe}(\text{s})$	-0.44
$\text{Cr}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Cr}(\text{s})$	-0.74
$\text{Zn}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Zn}(\text{s})$	-0.76
$2\text{H}_2\text{O} + 2\text{e}^- \rightarrow \text{H}_2(\text{g}) + 2\text{OH}^-(\text{aq})$	-0.83
$\text{Cr}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Cr}(\text{s})$	-0.89
$\text{Al}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Al}(\text{s})$	-1.68
$\text{Mg}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Mg}(\text{s})$	-2.36
$\text{Na}^+(\text{aq}) + \text{e}^- \rightarrow \text{Na}(\text{s})$	-2.71
$\text{Ca}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Ca}(\text{s})$	-2.87
$\text{Li}^+(\text{aq}) + \text{e}^- \rightarrow \text{Li}(\text{s})$	-3.04

## CHEM1108 - CHEMISTRY 1A LIFE SCIENCES

## Useful formulas

<p><b>Quantum Chemistry</b></p> $E = h\nu = hc/\lambda$ $\lambda = h/mv$ $E = -Z^2 E_R(1/n^2)$ $\Delta x \cdot \Delta(mv) \geq h/4\pi$ $q = 4\pi r^2 \times 5.67 \times 10^{-8} \times T^4$ $T\lambda = 2.898 \times 10^6 \text{ K nm}$	<p><b>Electrochemistry</b></p> $\Delta G^\circ = -nFE^\circ$ <p>Moles of <math>e^- = It/F</math></p> $E = E^\circ - (RT/nF) \times 2.303 \log Q$ $= E^\circ - (RT/nF) \times \ln Q$ $E^\circ = (RT/nF) \times 2.303 \log K$ $= (RT/nF) \times \ln K$ $E = E^\circ - \frac{0.0592}{n} \log Q \text{ (at } 25^\circ \text{C)}$
<p><b>Acids and Bases</b></p> $pK_w = \text{pH} + \text{pOH} = 14.00$ $pK_w = \text{p}K_a + \text{p}K_b = 14.00$ $\text{pH} = \text{p}K_a + \log \{ [A^-] / [HA] \}$	<p><b>Gas Laws</b></p> $PV = nRT$ $(P + n^2 a/V^2)(V - nb) = nRT$
<p><b>Radioactivity</b></p> $t_{1/2} = \ln 2 / \lambda$ $A = \lambda N$ $\ln(N_0/N_t) = \lambda t$ $^{14}\text{C age} = 8033 \ln(A_0/A_t) \text{ years}$	<p><b>Kinetics</b></p> $t_{1/2} = \ln 2 / k$ $k = Ae^{-E_a/RT}$ $\ln[A] = \ln[A]_0 - kt$ $\ln \frac{k_2}{k_1} = \frac{E_a}{R} \left( \frac{1}{T_1} - \frac{1}{T_2} \right)$
<p><b>Colligative properties</b></p> $\Pi = cRT$ $P_{\text{solution}} = X_{\text{solvent}} \times P^\circ_{\text{solvent}}$ $c = kp$ $\Delta T_f = K_f m$ $\Delta T_b = K_b m$	<p><b>Thermodynamics &amp; Equilibrium</b></p> $\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$ $\Delta G = \Delta G^\circ + RT \ln Q$ $\Delta G^\circ = -RT \ln K$ $\Delta_{\text{univ}} S^\circ = R \ln K$ $K_p = K_c (RT)^{\Delta n}$
<p><b>Miscellaneous</b></p> $A = -\log \frac{I}{I_0}$ $A = \epsilon cl$ $E = -A \frac{e^2}{4\pi\epsilon_0 r} N_A$	<p><b>Mathematics</b></p> <p>If <math>ax^2 + bx + c = 0</math>, then <math>x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}</math></p> $\ln x = 2.303 \log x$ <p>Area of circle = <math>\pi r^2</math></p> <p>Surface area of sphere = <math>4\pi r^2</math></p>

# PERIODIC TABLE OF THE ELEMENTS

<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>6</b>	<b>7</b>	<b>8</b>	<b>9</b>	<b>10</b>	<b>11</b>	<b>12</b>	<b>13</b>	<b>14</b>	<b>15</b>	<b>16</b>	<b>17</b>	<b>18</b>
1 HYDROGEN <b>H</b> 1.008												2 HELIUM <b>He</b> 4.003					
3 LITHIUM <b>Li</b> 6.941	4 BERYLLIUM <b>Be</b> 9.012											5 BORON <b>B</b> 10.81	6 CARBON <b>C</b> 12.01	7 NITROGEN <b>N</b> 14.01	8 OXYGEN <b>O</b> 16.00	9 FLUORINE <b>F</b> 19.00	10 NEON <b>Ne</b> 20.18
11 SODIUM <b>Na</b> 22.99	12 MAGNESIUM <b>Mg</b> 24.31											13 ALUMINIUM <b>Al</b> 26.98	14 SILICON <b>Si</b> 28.09	15 PHOSPHORUS <b>P</b> 30.97	16 SULFUR <b>S</b> 32.07	17 CHLORINE <b>Cl</b> 35.45	18 ARGON <b>Ar</b> 39.95
19 POTASSIUM <b>K</b> 39.10	20 CALCIUM <b>Ca</b> 40.08	21 SCANDIUM <b>Sc</b> 44.96	22 TITANIUM <b>Ti</b> 47.88	23 VANADIUM <b>V</b> 50.94	24 CHROMIUM <b>Cr</b> 52.00	25 MANGANESE <b>Mn</b> 54.94	26 IRON <b>Fe</b> 55.85	27 COBALT <b>Co</b> 58.93	28 NICKEL <b>Ni</b> 58.69	29 COPPER <b>Cu</b> 63.55	30 ZINC <b>Zn</b> 65.39	31 GALLIUM <b>Ga</b> 69.72	32 GERMANIUM <b>Ge</b> 72.59	33 ARSENIC <b>As</b> 74.92	34 SELENIUM <b>Se</b> 78.96	35 BROMINE <b>Br</b> 79.90	36 KRYPTON <b>Kr</b> 83.80
37 RUBIDIUM <b>Rb</b> 85.47	38 STRONTIUM <b>Sr</b> 87.62	39 YTTRIUM <b>Y</b> 88.91	40 ZIRCONIUM <b>Zr</b> 91.22	41 NIOBIUM <b>Nb</b> 92.91	42 MOLYBDENUM <b>Mo</b> 95.94	43 TECHNETIUM <b>Tc</b> [98.91]	44 RUTHENIUM <b>Ru</b> 101.07	45 RHODIUM <b>Rh</b> 102.91	46 PALLADIUM <b>Pd</b> 106.4	47 SILVER <b>Ag</b> 107.87	48 CADMIUM <b>Cd</b> 112.40	49 INDIUM <b>In</b> 114.82	50 TIN <b>Sn</b> 118.69	51 ANTIMONY <b>Sb</b> 121.75	52 TELLURIUM <b>Te</b> 127.60	53 IODINE <b>I</b> 126.90	54 XENON <b>Xe</b> 131.30
55 CAESIUM <b>Cs</b> 132.91	56 BARIUM <b>Ba</b> 137.34	57-71	72 HAFNIUM <b>Hf</b> 178.49	73 TANTALUM <b>Ta</b> 180.95	74 TUNGSTEN <b>W</b> 183.85	75 RHENIUM <b>Re</b> 186.2	76 OSMIUM <b>Os</b> 190.2	77 IRIDIUM <b>Ir</b> 192.22	78 PLATINUM <b>Pt</b> 195.09	79 GOLD <b>Au</b> 196.97	80 MERCURY <b>Hg</b> 200.59	81 THALLIUM <b>Tl</b> 204.37	82 LEAD <b>Pb</b> 207.2	83 BISMUTH <b>Bi</b> 208.98	84 POLONIUM <b>Po</b> [210.0]	85 ASTATINE <b>At</b> [210.0]	86 RADON <b>Rn</b> [222.0]
87 FRANCIUM <b>Fr</b> [223.0]	88 RADIUM <b>Ra</b> [226.0]	89-103	104 RUTHERFORDIUM <b>Rf</b> [261]	105 DUBNIUM <b>Db</b> [262]	106 SEABORGIUM <b>Sg</b> [266]	107 BOHRIUM <b>Bh</b> [262]	108 HASSIUM <b>Hs</b> [265]	109 MEITNERIUM <b>Mt</b> [266]	110 DARMSTADIUM <b>Ds</b> [271]	111 ROENTGENIUM <b>Rg</b> [272]							

LANTHANOID S	57 LANTHANUM <b>La</b> 138.91	58 CERIUM <b>Ce</b> 140.12	59 PRASEODYMIUM <b>Pr</b> 140.91	60 NEODYMIUM <b>Nd</b> 144.24	61 PROMETHIUM <b>Pm</b> [144.9]	62 SAMARIUM <b>Sm</b> 150.4	63 EUROPIUM <b>Eu</b> 151.96	64 GADOLINIUM <b>Gd</b> 157.25	65 TERBIUM <b>Tb</b> 158.93	66 DYSPROSIUM <b>Dy</b> 162.50	67 HOLMIUM <b>Ho</b> 164.93	68 ERBIUM <b>Er</b> 167.26	69 THULIUM <b>Tm</b> 168.93	70 YTTERBIUM <b>Yb</b> 173.04	71 LUTETIUM <b>Lu</b> 174.97
ACTINOIDS	89 ACTINIUM <b>Ac</b> [227.0]	90 THORIUM <b>Th</b> 232.04	91 PROTACTINIUM <b>Pa</b> [231.0]	92 URANIUM <b>U</b> 238.03	93 NEPTUNIUM <b>Np</b> [237.0]	94 PLUTONIUM <b>Pu</b> [239.1]	95 AMERICIUM <b>Am</b> [243.1]	96 CURIUM <b>Cm</b> [247.1]	97 BERKELIUM <b>Bk</b> [247.1]	98 CALIFORNIUM <b>Cf</b> [252.1]	99 EINSTEINIUM <b>Es</b> [252.1]	100 FERMIUM <b>Fm</b> [257.1]	101 MENDELEVIUM <b>Md</b> [256.1]	102 NOBELIUM <b>No</b> [259.1]	103 LAWRENCIUM <b>Lr</b> [260.1]