

CHEM1405 - CHEMISTRY (VETERINARY SCIENCE)

FIRST SEMESTER EXAMINATION

CONFIDENTIAL

JUNE 2009

TIME ALLOWED: THREE HOURS

GIVE THE FOLLOWING INFORMATION IN BLOCK LETTERS

FAMILY NAME		SID NUMBER	
OTHER NAMES		TABLE NUMBER	

INSTRUCTIONS TO CANDIDATES

- All questions are to be attempted. There are 21 pages of examinable material.
- Complete the written section of the examination paper in **INK**.
- Read each question carefully. Report the appropriate answer and show all relevant working in the space provided.
- The total score for this paper is 100. The possible score per page is shown in the adjacent tables.
- Each new short answer question begins with a •.
- Electronic calculators, including programmable calculators, may be used. Students are warned, however, that credit may not be given, even for a correct answer, where there is insufficient evidence of the working required to obtain the solution.
- Numerical values required for any question, standard electrode reduction potentials, a Periodic Table and some useful formulas may be found on the separate data sheet.
- Pages 16 and 24 are for rough working only.

OFFICIAL USE ONLY

Multiple choice section

		Marks	
Pages	Max	Gained	
2-10	32		

Short answer section


Page	Marks		Marker
	Max	Gained	
11	5		
12	6		
13	7		
14	4		
15	6		
17	6		
18	6		
19	6		
20	8		
21	3		
22	7		
23	4		
Total	68		

- In general terms, which elements in the periodic table are likely to be essential elements for living species and which ones are likely to be toxic. Explain.

Marks
2

- The diagram below shows the general trend for the first ionisation energy for some *s* and *p* block elements.

3



How will the general trend differ for the second ionisation energy of these elements (*i.e.* $X^+(g) \rightarrow X^{2+}(g) + e^-$)? Explain.

Marks
6

- Write the balanced chemical equation for the dissolution of solid $\text{Ca}(\text{CH}_3\text{CO}_2)_2$ in water.

What is the pH of a solution that has 158.2 g of $\text{Ca}(\text{CH}_3\text{CO}_2)_2$ dissolved in 1.000 L of water? The pK_a of acetic acid, CH_3COOH , is 4.76.

pH =

Calculate the pH of this solution after the addition of 0.250 mol of HCl gas?

pH =

Marks
2

- What physical state would water adopt under ambient conditions (1 atm and 25 °C) if it did not possess hydrogen bonding? Explain.

3

- Henry's law describes the solubility of a gas in a liquid phase. What methods are possible to ensure a patient receives enough oxygen during surgery? Which method is the most practical? Explain.

2

- A saline solution used to administer drugs intravenously is prepared by dissolving 0.90 g NaCl in 100.0 mL water. What mass of glucose ($C_6H_{12}O_6$) is required to prepare a 100.0 mL solution with the same osmotic pressure?





Answer:

Marks
4

- In the reaction of Cl_2 with Br_2 in CCl_4 solution, BrCl forms according to the equation:



With initial concentrations of $[\text{Br}_2] = 0.6 \text{ M}$, $[\text{Cl}_2] = 0.4 \text{ M}$ and $[\text{BrCl}] = 0.0 \text{ M}$, which of the following concentration versus time graphs represents this reaction? Explain why you rejected each of the other three graphs.



Marks
2

- The radioactive isotope ^{99m}Tc has a half life of 6.0 hours. How much time after production of the ^{99m}Tc isotope do radiologists have to examine a patient if at least 25 % of the original activity is required to get useful exposures?

Answer:

4

- Both $\text{HCO}_3^-(\text{aq})$ and $\text{CO}_2(\text{aq})$ are present in human blood. How does their presence ensure that the pH of blood is maintained at ~ 7.2 , even if $\text{H}^+(\text{aq})$ or $\text{OH}^-(\text{aq})$ are produced by processes in the body?

How does hyperventilation (very rapid breathing) interfere with this balance? What is the effect?

Marks
3

- Briefly explain the two factors necessary for a collision between two molecules to result in a reaction.

Briefly describe the relationship between the rate of a reaction and the activation energy for the reaction.

- The pH value of pure water at 25 °C is 7.00. How, if at all, does that value change when the temperature is changed to 37 °C (a person's body temperature)? Explain.



3

Is pure water at 37 °C acidic, basic or neutral? Circle your choice.

acidic	basic	neutral
--------	-------	---------

Marks
6

- The structures of histamine and polaramine are shown below.



Indicate the hybridisation and geometry of bonds around each of the nitrogen atoms in histamine.

	Hybridisation	Geometry of bonds
N #1:		
N #2:		
N #3:		

Draw a tautomer of histamine.

In histamine, only one of the nitrogen atoms in the ring is easily protonated (basic). Indicate which it is and explain why.

THIS QUESTION CONTINUES ON THE NEXT PAGE.




The release of histamine in the body triggers nasal secretions and constriction of airways. Polaramine is one of many anti-histamine compounds used to treat allergies. Explain what structural features of polaramine might make it a suitable anti-histamine agent.

Marks
3

(+)-2-[*p*-Chloro- α -[2-(dimethylamino)ethyl]benzyl]pyridine is another name for polaramine. What does the (+) in this name mean?


- Indicate the reagents used in the laboratory to effect the following transformations.

3


Reaction	Reagent
 <p>Reaction of cyclopentanecarboxylic acid to cyclopentanecarbonyl chloride.</p>	
 <p>Reaction of 2-bromobutane to N,N-dimethylbutan-2-amine.</p>	
 <p>Reaction of 4-hydroxy-2-pentanone to a cyclic acetal (2,2-dimethyl-1,3-dioxolane).</p>	

- The following species represent some of the building blocks of RNA.


Marks
8




ribofuranose



phosphate



cytosine



guanine

Is the sugar depicted the α - or the β - form? Circle the one correct answer.

α	β
----------	---------

Is the sugar depicted a reducing sugar or a non-reducing sugar? Circle the one correct answer.

reducing	non-reducing
----------	--------------

Indicate on the above structure the 'anomeric' carbon atom that gives rise to the α or the β form.


Draw the Fischer projection of D-ribose.

Using a selection of the species given, draw a nucleoside and a nucleotide.

Nucleoside	Nucleotide
------------	------------

Indicate the likely hydrogen-bonding interactions between complementary strands of RNA containing cytosine and guanine.

- Add curly arrows to complete the following mechanism.

Marks**1**

- Briefly explain Markovnikov's Rule in terms of the mechanism of electrophilic addition; *i.e.* why does Markovnikov's Rule work?


2

THE REMAINDER OF THIS PAGE IS FOR ROUGH WORKING ONLY.

Marks

7

- A peptide has the following structure.




Would you expect this peptide to be soluble in water? Explain your answer.





Give the products formed after treatment of the peptide with Zn/H^+ .

These products are then heated with excess aqueous OH^- . Draw the constitutional formulas of the different amino acids formed. Ensure you represent the amino acids in the correct charge state for the conditions.

Choose one of the amino acids produced on hydrolysis and draw the (*S*) configuration.



- Indicate the major organic product(s) in the following reactions.

Marks**4****THE REMAINDER OF THIS PAGE IS FOR ROUGH WORKING ONLY.**

CHEM1405 - CHEMISTRY (VETERINARY SCIENCE)**DATA SHEET***Physical constants*Avogadro constant, $N_A = 6.022 \times 10^{23} \text{ mol}^{-1}$ Faraday constant, $F = 96485 \text{ C mol}^{-1}$ Planck constant, $h = 6.626 \times 10^{-34} \text{ J s}$ Speed of light in vacuum, $c = 2.998 \times 10^8 \text{ m s}^{-1}$ Rydberg constant, $E_R = 2.18 \times 10^{-18} \text{ J}$ Boltzmann constant, $k_B = 1.381 \times 10^{-23} \text{ J K}^{-1}$ Permittivity of a vacuum, $\epsilon_0 = 8.854 \times 10^{-12} \text{ C}^2 \text{ J}^{-1} \text{ m}^{-1}$ Gas constant, $R = 8.314 \text{ J K}^{-1} \text{ mol}^{-1}$
 $= 0.08206 \text{ L atm K}^{-1} \text{ mol}^{-1}$ Charge of electron, $e = 1.602 \times 10^{-19} \text{ C}$ Mass of electron, $m_e = 9.1094 \times 10^{-31} \text{ kg}$ Mass of proton, $m_p = 1.6726 \times 10^{-27} \text{ kg}$ Mass of neutron, $m_n = 1.6749 \times 10^{-27} \text{ kg}$ *Properties of matter*

Volume of 1 mole of ideal gas at 1 atm and 25 °C = 24.5 L

Volume of 1 mole of ideal gas at 1 atm and 0 °C = 22.4 L

Density of water at 298 K = 0.997 g cm^{-3} *Conversion factors*

1 atm = 760 mmHg = 101.3 kPa

1 Ci = $3.70 \times 10^{10} \text{ Bq}$

0 °C = 273 K

1 Hz = 1 s^{-1} 1 L = 10^{-3} m^3 1 tonne = 10^3 kg 1 Å = 10^{-10} m 1 W = 1 J s^{-1} 1 eV = $1.602 \times 10^{-19} \text{ J}$ *Decimal fractions*

Fraction	Prefix	Symbol
10^{-3}	milli	m
10^{-6}	micro	μ
10^{-9}	nano	n
10^{-12}	pico	p

Decimal multiples

Multiple	Prefix	Symbol
10^3	kilo	k
10^6	mega	M
10^9	giga	G

CHEM1405 - CHEMISTRY (VETERINARY SCIENCE)*Standard Reduction Potentials, E°*

Reaction	E° / V
$\text{Co}^{3+}(\text{aq}) + \text{e}^- \rightarrow \text{Co}^{2+}(\text{aq})$	+1.82
$\text{Ce}^{4+}(\text{aq}) + \text{e}^- \rightarrow \text{Ce}^{3+}(\text{aq})$	+1.72
$\text{MnO}_4^-(\text{aq}) + 8\text{H}^+(\text{aq}) + 5\text{e}^- \rightarrow \text{Mn}^{2+}(\text{aq}) + 4\text{H}_2\text{O}$	+1.51
$\text{Au}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Au}(\text{s})$	+1.50
$\text{Cl}_2 + 2\text{e}^- \rightarrow 2\text{Cl}^-(\text{aq})$	+1.36
$\text{O}_2 + 4\text{H}^+(\text{aq}) + 4\text{e}^- \rightarrow 2\text{H}_2\text{O}$	+1.23
$\text{Pt}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Pt}(\text{s})$	+1.18
$\text{MnO}_2(\text{s}) + 4\text{H}^+(\text{aq}) + \text{e}^- \rightarrow \text{Mn}^{3+} + 2\text{H}_2\text{O}$	+0.96
$\text{NO}_3^-(\text{aq}) + 4\text{H}^+(\text{aq}) + 3\text{e}^- \rightarrow \text{NO}(\text{g}) + 2\text{H}_2\text{O}$	+0.96
$\text{Pd}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Pd}(\text{s})$	+0.92
$\text{Ag}^+(\text{aq}) + \text{e}^- \rightarrow \text{Ag}(\text{s})$	+0.80
$\text{Fe}^{3+}(\text{aq}) + \text{e}^- \rightarrow \text{Fe}^{2+}(\text{aq})$	+0.77
$\text{Cu}^+(\text{aq}) + \text{e}^- \rightarrow \text{Cu}(\text{s})$	+0.53
$\text{Cu}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Cu}(\text{s})$	+0.34
$\text{Sn}^{4+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Sn}^{2+}(\text{aq})$	+0.15
$2\text{H}^+(\text{aq}) + 2\text{e}^- \rightarrow \text{H}_2(\text{g})$	0 (by definition)
$\text{Fe}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Fe}(\text{s})$	-0.04
$\text{Pb}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Pb}(\text{s})$	-0.13
$\text{Sn}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Sn}(\text{s})$	-0.14
$\text{Ni}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Ni}(\text{s})$	-0.24
$\text{Cd}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Cd}(\text{s})$	-0.40
$\text{Fe}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Fe}(\text{s})$	-0.44
$\text{Cr}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Cr}(\text{s})$	-0.74
$\text{Zn}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Zn}(\text{s})$	-0.76
$2\text{H}_2\text{O} + 2\text{e}^- \rightarrow \text{H}_2(\text{g}) + 2\text{OH}^-(\text{aq})$	-0.83
$\text{Cr}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Cr}(\text{s})$	-0.89
$\text{Al}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Al}(\text{s})$	-1.68
$\text{Mg}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Mg}(\text{s})$	-2.36
$\text{Na}^+(\text{aq}) + \text{e}^- \rightarrow \text{Na}(\text{s})$	-2.71
$\text{Ca}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Ca}(\text{s})$	-2.87
$\text{Li}^+(\text{aq}) + \text{e}^- \rightarrow \text{Li}(\text{s})$	-3.04

CHEM1405 - CHEMISTRY (VETERINARY SCIENCE)

Useful formulas

<p>Quantum Chemistry</p> $E = h\nu = hc/\lambda$ $\lambda = h/mv$ $E = -Z^2 E_R(1/n^2)$ $\Delta x \cdot \Delta(mv) \geq h/4\pi$ $q = 4\pi r^2 \times 5.67 \times 10^{-8} \times T^4$ $T\lambda = 2.898 \times 10^6 \text{ K nm}$	<p>Electrochemistry</p> $\Delta G^\circ = -nFE^\circ$ <p>Moles of $e^- = It/F$</p> $E = E^\circ - (RT/nF) \times 2.303 \log Q$ $= E^\circ - (RT/nF) \times \ln Q$ $E^\circ = (RT/nF) \times 2.303 \log K$ $= (RT/nF) \times \ln K$ $E = E^\circ - \frac{0.0592}{n} \log Q \text{ (at } 25^\circ \text{C)}$
<p>Acids and Bases</p> $pK_w = \text{pH} + \text{pOH} = 14.00$ $pK_w = \text{p}K_a + \text{p}K_b = 14.00$ $\text{pH} = \text{p}K_a + \log \left\{ \frac{[A^-]}{[HA]} \right\}$	<p>Gas Laws</p> $PV = nRT$ $(P + n^2 a/V^2)(V - nb) = nRT$
<p>Radioactivity</p> $t_{1/2} = \ln 2/\lambda$ $A = \lambda N$ $\ln(N_0/N_t) = \lambda t$ $^{14}\text{C age} = 8033 \ln(A_0/A_t) \text{ years}$	<p>Kinetics</p> $t_{1/2} = \ln 2/k$ $k = Ae^{-E_a/RT}$ $\ln[A] = \ln[A]_0 - kt$ $\ln \frac{k_2}{k_1} = \frac{E_a}{R} \left(\frac{1}{T_1} - \frac{1}{T_2} \right)$
<p>Colligative properties</p> $\Pi = cRT$ $P_{\text{solution}} = X_{\text{solvent}} \times P^\circ_{\text{solvent}}$ $c = kp$ $\Delta T_f = K_f m$ $\Delta T_b = K_b m$	<p>Thermodynamics & Equilibrium</p> $\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$ $\Delta G = \Delta G^\circ + RT \ln Q$ $\Delta G^\circ = -RT \ln K$ $\Delta_{\text{univ}} S^\circ = R \ln K$ $K_p = K_c (RT)^{\Delta n}$
<p>Miscellaneous</p> $A = -\log \frac{I}{I_0}$ $A = \epsilon cl$ $E = -A \frac{e^2}{4\pi\epsilon_0 r} N_A$	<p>Mathematics</p> <p>If $ax^2 + bx + c = 0$, then $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$</p> $\ln x = 2.303 \log x$ <p>Area of circle = πr^2</p> <p>Surface area of sphere = $4\pi r^2$</p>

