

CHEM1405 - CHEMISTRY (VETERINARY SCIENCE)FIRST SEMESTER EXAMINATION**CONFIDENTIAL****JUNE 2007****TIME ALLOWED: THREE HOURS**

GIVE THE FOLLOWING INFORMATION IN BLOCK LETTERS

FAMILY NAME		SID NUMBER	
OTHER NAMES		TABLE NUMBER	

INSTRUCTIONS TO CANDIDATES

- All questions are to be attempted. There are 20 pages of examinable material.
- Complete the written section of the examination paper in **INK**.
- Read each question carefully. Report the appropriate answer and show all relevant working in the space provided.
- The total score for this paper is 100. The possible score per page is shown in the adjacent tables.
- Each new short answer question begins with a •.
- Electronic calculators, including programmable calculators, may be used. Students are warned, however, that credit may not be given, even for a correct answer, where there is insufficient evidence of the working required to obtain the solution.
- Numerical values required for any question, standard electrode reduction potentials, a Periodic Table and some useful formulas may be found on the separate data sheet.
- Pages 17, 23 and 24 are for rough working only.

OFFICIAL USE ONLY**Multiple choice section**

	Marks	
Pages	Max	Gained
2-12	45	

Short answer section

Page	Marks		Marker
	Max	Gained	
13	5		
14	9		
15	5		
16	5		
18	6		
19	6		
20	7		
21	6		
22	6		
Total	55		

Marks
5

- In the spaces provided, explain the meanings of the following terms. You may use an equation or diagram where appropriate.

(a) hydrogen bonding

(b) colligative properties

(c) hypotonic solution

(d) isoelectric point

(e) half life

Marks
9

- Complete the table below showing the number of valence electrons, the Lewis structure and the VSEPR predicted shape of each of the following species.

Formula	Number of valence electrons	Lewis structure	Geometry of species
e.g. NH ₃	8	$\begin{array}{c} \text{H}-\ddot{\text{N}}-\text{H} \\ \\ \text{H} \end{array}$	trigonal pyramidal
CH ₄			
CO ₂			
PF ₅			
NO ₃ ⁻			

Which one of the species above displays resonance, and how many resonance forms are possible?



--

- Give the oxidation number of carbon in each of the following.

CF ₂ Cl ₂ (g)	
Na ₂ C ₂ O ₄ (s)	
HCO ₃ ⁻ (aq)	
C(s)	

Marks
2

- Consider a voltaic cell that uses the following half-reactions:



Write a balanced equation for the overall reaction.

3

Which species is the oxidising agent?


Which species is the reducing agent?

Calculate the standard cell potential. (Refer to the table of standard reduction potentials.)

Answer:

Marks
5

- Stearic acid, $C_{18}H_{36}O_2$, is a fatty acid common in animal fats and vegetable oils and is a valuable energy source for mammals. The net reaction for its metabolism in humans is:



Calculate ΔH° for this reaction given the following heats of formation.

$$\Delta H_f^\circ (C_{18}H_{36}O_2(s)) = -948 \text{ kJ mol}^{-1}, \quad \Delta H_f^\circ (CO_2(g)) = -393 \text{ kJ mol}^{-1} \text{ and}$$

$$\Delta H_f^\circ (H_2O(l)) = -285 \text{ kJ mol}^{-1}$$

Answer:

If the combustion of stearic acid is carried out in air, water is produced as a vapour. Calculate the ΔH° for the combustion of stearic acid in air given that



Answer:

Will ΔS be different for the two oxidation reactions? If so, how will it differ and why?

Calculate the mass of carbon dioxide produced by the complete oxidation of 1.00 g of stearic acid.

Answer:

Marks
6

- Lactic acid, $\text{CH}_3\text{CHOHCOOH}$, is produced in the body during normal exercise. It is a monoprotic acid with a $\text{p}K_a$ of 3.86.

(a) What is the pH of a 0.10 M water solution of lactic acid?

Answer:


(b) Calculate the pH of the solution formed when 0.02 mol of $\text{Ca}(\text{OH})_2(\text{s})$ is added to 1.0 L of 0.10 M lactic acid.

Answer:

(c) Using equations, comment on how the final solution in (b) will respond to additions of small amounts (e.g. less than 0.01 mol) of acid or base in comparison to additions of the same amounts of acid or base to 1 L of water.

Marks
6

- Orlistat (shown below) is a drug for obesity management which acts by inhibiting the absorption of dietary fats.



Is it likely to be soluble in water? Why?


Indicate on the above structure all stereogenic centres. Select one of these centres and clearly assign its stereochemical configuration.

Name the functional groups present in orlistat.


THE REMAINDER OF THIS PAGE IS FOR ROUGH WORKING ONLY

Marks**7**

- Glycine, $\text{NH}_2\text{CH}_2\text{COOH}$, is the simplest of all naturally occurring amino acids. The $\text{p}K_a$ of the acid group is 2.35 and the $\text{p}K_a$ associated with the amino group is 9.78. Draw a structure that indicates the charges on the molecule at the physiological pH of 7.4.




Use your structure to illustrate the concept of resonance.



What are the hybridisation states and geometries of the two carbon atoms and the nitrogen atom in glycine?



Propionic acid, $\text{CH}_3\text{CH}_2\text{COOH}$, has a melting point of -20.7°C while glycine has a melting point of 292°C . Suggest a reason why these two molecules have such different melting points.



Marks
6


- Alanine ($R = \text{CH}_3$) and lysine ($R = \text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{NH}_2$) are two common amino acids. Using *ala* and *lys* to represent the two amino acids, represent all constitutional isomers of the tripeptide formed from one *ala* and two *lys* units.

Comment, giving your reason, on whether the tripeptide(s) will be acidic, neutral or basic in character.

Draw the constitutional formulas, indicating the correct ionic state, of the products formed from acid hydrolysis of one of your tripeptides.

THE REMAINDER OF THIS PAGE IS FOR ROUGH WORKING ONLY

- NAD^+ and NADH are coenzymes used by animals in oxidation and reduction reactions. They are related by the following half-reactions.



Which of these coenzymes is used in the biological oxidation of ethanol, $\text{CH}_3\text{CH}_2\text{OH}$?

What is the product of the biological oxidation of ethanol, $\text{CH}_3\text{CH}_2\text{OH}$?

Which of NAD^+ and NADH is aromatic? Give reasons for your answer.

Marks
4

- Benzoic acid has a low solubility in water at pH 7, but is very soluble in aqueous solutions of greater pH. Explain this observation, using chemical equations where appropriate.

2

CHEM1405 - CHEMISTRY (VETERINARY SCIENCE)**DATA SHEET***Physical constants*Avogadro constant, $N_A = 6.022 \times 10^{23} \text{ mol}^{-1}$ Faraday constant, $F = 96485 \text{ C mol}^{-1}$ Planck constant, $h = 6.626 \times 10^{-34} \text{ J s}$ Speed of light in vacuum, $c = 2.998 \times 10^8 \text{ m s}^{-1}$ Rydberg constant, $E_R = 2.18 \times 10^{-18} \text{ J}$ Boltzmann constant, $k_B = 1.381 \times 10^{-23} \text{ J K}^{-1}$ Permittivity of a vacuum, $\epsilon_0 = 8.854 \times 10^{-12} \text{ C}^2 \text{ J}^{-1} \text{ m}^{-1}$ Gas constant, $R = 8.314 \text{ J K}^{-1} \text{ mol}^{-1}$
 $= 0.08206 \text{ L atm K}^{-1} \text{ mol}^{-1}$ Charge of electron, $e = 1.602 \times 10^{-19} \text{ C}$ Mass of electron, $m_e = 9.1094 \times 10^{-31} \text{ kg}$ Mass of proton, $m_p = 1.6726 \times 10^{-27} \text{ kg}$ Mass of neutron, $m_n = 1.6749 \times 10^{-27} \text{ kg}$ *Properties of matter*

Volume of 1 mole of ideal gas at 1 atm and 25 °C = 24.5 L

Volume of 1 mole of ideal gas at 1 atm and 0 °C = 22.4 L

Density of water at 298 K = 0.997 g cm^{-3} *Conversion factors*

1 atm = 760 mmHg = 101.3 kPa

1 Ci = $3.70 \times 10^{10} \text{ Bq}$

0 °C = 273 K

1 Hz = 1 s^{-1} 1 L = 10^{-3} m^3 1 tonne = 10^3 kg 1 Å = 10^{-10} m 1 W = 1 J s^{-1} 1 eV = $1.602 \times 10^{-19} \text{ J}$ *Decimal fractions*

Fraction	Prefix	Symbol
10^{-3}	milli	m
10^{-6}	micro	μ
10^{-9}	nano	n
10^{-12}	pico	p

Decimal multiples

Multiple	Prefix	Symbol
10^3	kilo	k
10^6	mega	M
10^9	giga	G

CHEM1405 - CHEMISTRY (VETERINARY SCIENCE)*Standard Reduction Potentials, E°*

Reaction	E° / V
$\text{Co}^{3+}(\text{aq}) + \text{e}^{-} \rightarrow \text{Co}^{2+}(\text{aq})$	+1.82
$\text{Ce}^{4+}(\text{aq}) + \text{e}^{-} \rightarrow \text{Ce}^{3+}(\text{aq})$	+1.72
$\text{MnO}_4^{-}(\text{aq}) + 8\text{H}^{+}(\text{aq}) + 5\text{e}^{-} \rightarrow \text{Mn}^{2+}(\text{aq}) + 4\text{H}_2\text{O}$	+1.51
$\text{Au}^{3+}(\text{aq}) + 3\text{e}^{-} \rightarrow \text{Au}(\text{s})$	+1.50
$\text{Cl}_2 + 2\text{e}^{-} \rightarrow 2\text{Cl}^{-}(\text{aq})$	+1.36
$\text{O}_2 + 4\text{H}^{+}(\text{aq}) + 4\text{e}^{-} \rightarrow 2\text{H}_2\text{O}$	+1.23
$\text{Pt}^{2+}(\text{aq}) + 2\text{e}^{-} \rightarrow \text{Pt}(\text{s})$	+1.18
$\text{MnO}_2(\text{s}) + 4\text{H}^{+}(\text{aq}) + \text{e}^{-} \rightarrow \text{Mn}^{3+} + 2\text{H}_2\text{O}$	+0.96
$\text{Pd}^{2+}(\text{aq}) + 2\text{e}^{-} \rightarrow \text{Pd}(\text{s})$	+0.92
$\text{Ag}^{+}(\text{aq}) + \text{e}^{-} \rightarrow \text{Ag}(\text{s})$	+0.80
$\text{Fe}^{3+}(\text{aq}) + \text{e}^{-} \rightarrow \text{Fe}^{2+}(\text{aq})$	+0.77
$\text{Cu}^{+}(\text{aq}) + \text{e}^{-} \rightarrow \text{Cu}(\text{s})$	+0.53
$\text{Cu}^{2+}(\text{aq}) + 2\text{e}^{-} \rightarrow \text{Cu}(\text{s})$	+0.34
$\text{Sn}^{4+}(\text{aq}) + 2\text{e}^{-} \rightarrow \text{Sn}^{2+}(\text{aq})$	+0.15
$2\text{H}^{+}(\text{aq}) + 2\text{e}^{-} \rightarrow \text{H}_2(\text{g})$	0 (by definition)
$\text{Fe}^{3+}(\text{aq}) + 3\text{e}^{-} \rightarrow \text{Fe}(\text{s})$	-0.04
$\text{Pb}^{2+}(\text{aq}) + 2\text{e}^{-} \rightarrow \text{Pb}(\text{s})$	-0.13
$\text{Sn}^{2+}(\text{aq}) + 2\text{e}^{-} \rightarrow \text{Sn}(\text{s})$	-0.14
$\text{Ni}^{2+}(\text{aq}) + 2\text{e}^{-} \rightarrow \text{Ni}(\text{s})$	-0.24
$\text{Cd}^{2+}(\text{aq}) + 2\text{e}^{-} \rightarrow \text{Cd}(\text{s})$	-0.40
$\text{Fe}^{2+}(\text{aq}) + 2\text{e}^{-} \rightarrow \text{Fe}(\text{s})$	-0.44
$\text{Cr}^{3+}(\text{aq}) + 3\text{e}^{-} \rightarrow \text{Cr}(\text{s})$	-0.74
$\text{Zn}^{2+}(\text{aq}) + 2\text{e}^{-} \rightarrow \text{Zn}(\text{s})$	-0.76
$2\text{H}_2\text{O} + 2\text{e}^{-} \rightarrow \text{H}_2(\text{g}) + 2\text{OH}^{-}(\text{aq})$	-0.83
$\text{Cr}^{2+}(\text{aq}) + 2\text{e}^{-} \rightarrow \text{Cr}(\text{s})$	-0.89
$\text{Al}^{3+}(\text{aq}) + 3\text{e}^{-} \rightarrow \text{Al}(\text{s})$	-1.68
$\text{Mg}^{2+}(\text{aq}) + 2\text{e}^{-} \rightarrow \text{Mg}(\text{s})$	-2.36
$\text{Na}^{+}(\text{aq}) + \text{e}^{-} \rightarrow \text{Na}(\text{s})$	-2.71
$\text{Ca}^{2+}(\text{aq}) + 2\text{e}^{-} \rightarrow \text{Ca}(\text{s})$	-2.87
$\text{Li}^{+}(\text{aq}) + \text{e}^{-} \rightarrow \text{Li}(\text{s})$	-3.04

CHEM1405 - CHEMISTRY (VETERINARY SCIENCE)

Useful formulas

<p>Quantum Chemistry</p> $E = h\nu = hc/\lambda$ $\lambda = h/mv$ $4.5k_B T = hc/\lambda$ $E = -Z^2 E_R (1/n^2)$ $\Delta x \cdot \Delta(mv) \geq h/4\pi$ $q = 4\pi r^2 \times 5.67 \times 10^{-8} \times T^4$	<p>Electrochemistry</p> $\Delta G^\circ = -nFE^\circ$ <p>Moles of $e^- = It/F$</p> $E = E^\circ - (RT/nF) \times 2.303 \log Q$ $= E^\circ - (RT/nF) \times \ln Q$ $E^\circ = (RT/nF) \times 2.303 \log K$ $= (RT/nF) \times \ln K$ $E = E^\circ - \frac{0.0592}{n} \log Q \text{ (at } 25^\circ \text{C)}$
<p>Acids and Bases</p> $pK_w = \text{pH} + \text{pOH} = 14.00$ $pK_w = \text{p}K_a + \text{p}K_b = 14.00$ $\text{pH} = \text{p}K_a + \log \{ [A^-] / [HA] \}$	<p>Gas Laws</p> $PV = nRT$ $(P + n^2 a/V^2)(V - nb) = nRT$
<p>Colligative properties</p> $\pi = cRT$ $P_{\text{solution}} = X_{\text{solvent}} \times P^\circ_{\text{solvent}}$ $p = kc$ $\Delta T_f = K_f m$ $\Delta T_b = K_b m$	<p>Kinetics</p> $t_{1/2} = \ln 2/k$ $k = Ae^{-E_a/RT}$ $\ln[A] = \ln[A]_0 - kt$ $\ln \frac{k_2}{k_1} = \frac{E_a}{R} \left(\frac{1}{T_1} - \frac{1}{T_2} \right)$
<p>Radioactivity</p> $t_{1/2} = \ln 2/\lambda$ $A = \lambda N$ $\ln(N_0/N_t) = \lambda t$ $^{14}\text{C age} = 8033 \ln(A_0/A_t) \text{ years}$	<p>Thermodynamics & Equilibrium</p> $\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$ $\Delta G = \Delta G^\circ + RT \ln Q$ $\Delta G^\circ = -RT \ln K$ $K_p = K_c (RT)^{\Delta n}$
<p>Miscellaneous</p> $A = -\log_{10} \frac{I}{I_0}$ $A = \epsilon cl$ $E = -A \frac{e^2}{4\pi\epsilon_0 r} N_A$	<p>Mathematics</p> <p>If $ax^2 + bx + c = 0$, then $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$</p> $\ln x = 2.303 \log x$

PERIODIC TABLE OF THE ELEMENTS

June 2007

CHEM1405 - CHEMISTRY (VETERINARY SCIENCE)

22/25(b)

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
1 HYDROGEN H 1.008																	2 HELIUM He 4.003
3 LITHIUM Li 6.941	4 BERYLLIUM Be 9.012											5 BORON B 10.81	6 CARBON C 12.01	7 NITROGEN N 14.01	8 OXYGEN O 16.00	9 FLUORINE F 19.00	10 NEON Ne 20.18
11 SODIUM Na 22.99	12 MAGNESIUM Mg 24.31											13 ALUMINIUM Al 26.98	14 SILICON Si 28.09	15 PHOSPHORUS P 30.97	16 SULFUR S 32.07	17 CHLORINE Cl 35.45	18 ARGON Ar 39.95
19 POTASSIUM K 39.10	20 CALCIUM Ca 40.08	21 SCANDIUM Sc 44.96	22 TITANIUM Ti 47.88	23 VANADIUM V 50.94	24 CHROMIUM Cr 52.00	25 MANGANESE Mn 54.94	26 IRON Fe 55.85	27 COBALT Co 58.93	28 NICKEL Ni 58.69	29 COPPER Cu 63.55	30 ZINC Zn 65.39	31 GALLIUM Ga 69.72	32 GERMANIUM Ge 72.59	33 ARSENIC As 74.92	34 SELENIUM Se 78.96	35 BROMINE Br 79.90	36 KRYPTON Kr 83.80
37 RUBIDIUM Rb 85.47	38 STRONTIUM Sr 87.62	39 YTRIUM Y 88.91	40 ZIRCONIUM Zr 91.22	41 NOBIUM Nb 92.91	42 MOLYBDENUM Mo 95.94	43 TECHNETIUM Tc [98.91]	44 RUTHENIUM Ru 101.07	45 RHODIUM Rh 102.91	46 PALLADIUM Pd 106.4	47 SILVER Ag 107.87	48 CADMIUM Cd 112.40	49 INDIUM In 114.82	50 TIN Sn 118.69	51 ANTIMONY Sb 121.75	52 TELLURIUM Te 127.60	53 IODINE I 126.90	54 XENON Xe 131.30
55 CAESIUM Cs 132.91	56 BARIUM Ba 137.34	57-71	72 HAFNIUM Hf 178.49	73 TANTALUM Ta 180.95	74 TUNGSTEN W 183.85	75 RHENIUM Re 186.2	76 OSMIUM Os 190.2	77 IRIDIUM Ir 192.22	78 PLATINUM Pt 195.09	79 GOLD Au 196.97	80 MERCURY Hg 200.59	81 THALLIUM Tl 204.37	82 LEAD Pb 207.2	83 BISMUTH Bi 208.98	84 POLONIUM Po [210.0]	85 ASTATINE At [210.0]	86 RADON Rn [222.0]
87 FRANCIUM Fr [223.0]	88 RADIUM Ra [226.0]	89-103	104 RUTHERFORDIUM Rf [261]	105 DUBNIUM Db [262]	106 SEABORGIUM Sg [266]	107 BOHRIUM Bh [262]	108 HASSIUM Hs [265]	109 MEITNERIUM Mt [266]	110 DARMSTADTIUM Ds [271]	111 ROENTGENIUM Rg [272]							

LANTHANIDES	57 LANTHANUM La 138.91	58 CERIUM Ce 140.12	59 PRASEODYMIUM Pr 140.91	60 NEODYMIUM Nd 144.24	61 PROMETHIUM Pm [144.9]	62 SAMARIUM Sm 150.4	63 EUROPIUM Eu 151.96	64 GADOLINIUM Gd 157.25	65 TERBIUM Tb 158.93	66 DYSPROSIUM Dy 162.50	67 HOLMIUM Ho 164.93	68 ERBIUM Er 167.26	69 THULIUM Tm 168.93	70 YTTERBIUM Yb 173.04	71 LUTETIUM Lu 174.97
ACTINIDES	89 ACTINIUM Ac [227.0]	90 THORIUM Th 232.04	91 PROFACINIUM Pa [231.0]	92 URANIUM U 238.03	93 NEPTUNIUM Np [237.0]	94 PLUTONIUM Pu [239.1]	95 AMERICIUM Am [243.1]	96 CURIUM Cm [247.1]	97 BERKELIUM Bk [247.1]	98 CALIFORNIUM Cf [252.1]	99 EINSTEINIUM Es [252.1]	100 FERMIUM Fm [257.1]	101 MENDELEVIUM Md [256.1]	102 NOBELIUM No [259.1]	103 LAWRENCIUM Lr [260.1]