

Topics in the June 2013 Exam Paper for CHEM1611

Click on the links for resources on each topic.

2013-J-2:

- [Atomic Structure](#)
- [Chemical Bonding](#)

2013-J-3:

- [Chemical Bonding](#)

2013-J-4:

- [Chemical Bonding](#)
- [The Shapes of Molecules](#)

2013-J-5:

- [Atomic Structure](#)

2013-J-6:

- [Alkenes](#)
- [Alcohols, Phenols, Ethers and Thiols](#)
- [Carboxylic Acids and Derivatives](#)
- [Amines](#)

2013-J-7:

- [Introduction to Organic Chemistry](#)
- [Stereochemistry](#)

2013-J-8:

- [Alkenes](#)
- [Alcohols, Phenols, Ethers and Thiols](#)
- [Carboxylic Acids and Derivatives](#)

2013-J-9:

- [Carbohydrates](#)

2013-J-10:

- [Amino Acids, Peptides and Proteins](#)

2013-J-11:

- [Amino Acids, Peptides and Proteins](#)

2013-J-12:

- [DNA and Nucleic Acids](#)

2216(a)

THE UNIVERSITY OF SYDNEY

CHEM1611 - CHEMISTRY 1A (PHARMACY)

FIRST SEMESTER EXAMINATION

CONFIDENTIAL

JUNE 2013

TIME ALLOWED: THREE HOURS

GIVE THE FOLLOWING INFORMATION IN BLOCK LETTERS

FAMILY NAME		SID NUMBER	
OTHER NAMES		TABLE NUMBER	

OFFICIAL USE ONLY

- All questions are to be attempted. There are 19 pages of examinable material.
- Complete the examination paper in **INK**.
- Read each question carefully. Report the appropriate answer and show all relevant working in the space provided.
- The total score for this paper is 100. The possible score per page is shown in the adjacent tables.
- Each new short answer question begins with a •.
- Only non-programmable, University-approved calculators may be used.
- Students are warned that credit may not be given, even for a correct answer, where there is insufficient evidence of the working required to obtain the solution.
- Numerical values required for any question, standard electrode reduction potentials, a Periodic Table and some useful formulas may be found on the separate data sheets.
- Pages 14, 16, 22 and 24 are for rough work only.

~~Multiple choice section~~

		Marks	
Pages	Max	Gained	
2-9	30		

Short answer section

Page	Marks		Marker
	Max	Gained	
10	4		
11	5		
12	10		
13	5		
15	11		
17	6		
18	6		
19	6		
20	6		
21	6		
23	5		
Total	70		
Check Total			

- In the spaces provided, briefly explain the meaning of the following terms.

Marks
4

Effective nuclear charge

Atomic emission spectrum

Ionic bonding

Core electrons


THE REMAINDER OF THIS PAGE IS FOR ROUGH WORKING ONLY.

- Silicon and carbon are both in Group 14 and form dioxides. Carbon dioxide is a gas at room temperature while silicon dioxide (sand) is a solid with a high melting point. Describe the bonding in these two materials and explain the differences in properties they show.

Marks
3

- Complete the following nuclear equations by filling in the missing particle.

2



THE REMAINDER OF THIS PAGE IS FOR ROUGH WORKING ONLY.

- Draw the Lewis structure of the acetate ion (CH_3COO^-) showing all appropriate resonance structures.

Marks
10

Indicate the hybridisation, molecular geometry and approximate bond angle about each of the carbon atoms in the acetate ion.

	$-\text{CH}_3$	$-\text{COO}^-$
Hybridisation of C		
Molecular geometry about C		
Approximate bond angles about C		

The actual structure of the acetate ion is a weighted combination of all resonance structures. Sketch the σ -bond framework of the acetate ion and indicate the p -orbitals that are involved in the π -bonding of the acetate ion.

How many electrons are involved with the π -bonding?

What is the hybridisation of the oxygen atoms in the acetate ion?


- The yellow light emitted from an excited sodium atom has a wavelength of 590 nm. What is the energy of one photon of this light and one mole of photons? Specify appropriate units with your answers.

Marks
5

Energy	of one photon:	of 1 mol of photons:
--------	----------------	----------------------

The yellow light is associated with the longest wavelength transition as the atom returns to the ground state electron configuration. Fill in the following energy level diagram for sodium and indicate the transition associated with the emission of yellow light.










Energy



A quantum mechanical model of an atom can explain the emission spectrum of sodium, but the Bohr model of the atom cannot. Why?

- Complete the following table. Make sure you complete the name of the starting material where indicated.


Marks
11

STARTING MATERIAL	REAGENTS/ CONDITIONS	CONSTITUTIONAL FORMULA(S) OF MAJOR ORGANIC PRODUCT(S)
	HBr / CCl ₄ (solvent)	
 Name:		
	1. Mg / dry ether 2. CO ₂ 3. H ⁺ / H ₂ O	
	3 M NaOH	
	excess (CH ₃) ₂ NH	
 Name:		
	2 M HCl	

- Oseltamivir, marketed under the trade name Tamiflu, is an antiviral drug, which may slow the spread of influenza (flu) virus between cells in the body by stopping the virus from chemically cutting ties with its host cell.

Marks
6

Tamiflu




Give the molecular formula of Tamiflu.

List the functional groups present in Tamiflu.

How many stereogenic centres are there in Tamiflu?



How many possible stereoisomers can exist for Tamiflu?

Add the NH₂ and H groups to the stereogenic centre indicated below to give the (*R*)-configuration of that centre.





- Show clearly the reagents you would use to carry out the following chemical conversions. Note that more than one step is required and you should indicate all necessary steps and the constitutional formulas of any intermediate compounds.

Marks
6



- Consider the following two monosaccharides, **(A)** and **(B)**.

 β -D-altropyranose α -D-xylofuranose

Draw Fischer projections of the open chain forms of **(A)** and **(B)**.

(A)	(B)
------------	------------


Draw the major organic product of the reaction of D-altropyranose with the following reagents.

1. NaBH_4 2. $\text{H}^{\oplus} / \text{H}_2\text{O}$	$[\text{Ag}(\text{NH}_3)_2]^{\oplus} / \text{OH}^{\ominus}$
--	---

Draw the Haworth stereoformula of a non-reducing disaccharide formed from **(A)** and **(B)**.

Marks
6

- The amino acid, asparagine, was isolated from asparagus juice in 1806. The uncharged form, **Y**, is given below.



Draw the constitutional formula of the product(s) formed in the reaction of **Y** with the following reagents.

Marks
6

Cold, dilute hydrochloric acid	Cold, dilute sodium hydroxide
Hot, 6 M hydrochloric acid	Hot, 6 M sodium hydroxide

THE REMAINDER OF THIS PAGE IS FOR ROUGH WORKING ONLY.

Marks
6

- Alanine ($R = \text{CH}_3$) and lysine ($R = \text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{NH}_2$) are two common amino acids. Using *ala* and *lys* to represent the two amino acids, represent all constitutional isomers of the tripeptide formed from one *ala* and two *lys* units.

--

Comment, giving your reason(s), on whether the tripeptide(s) will be acidic, neutral or basic in character.

--

The pK_a values of lysine are 1.82 ($\alpha\text{-COOH}$), 8.95 ($\alpha\text{-NH}_3^+$) and 10.53 (side chain). What is the value of the isoelectric point of lysine?

--


pI =

Draw the Fischer projection of the zwitterionic form of lysine.

--

- Consider the following structure.

Marks
5



Is it a fragment of DNA or RNA? Give two reasons.

Clearly identify on the above structure one example of each of the following subunits.

nucleic base

nucleoside

nucleotide

CHEM1611 - CHEMISTRY 1A (PHARMACY)**DATA SHEET***Physical constants*Avogadro constant, $N_A = 6.022 \times 10^{23} \text{ mol}^{-1}$ Faraday constant, $F = 96485 \text{ C mol}^{-1}$ Planck constant, $h = 6.626 \times 10^{-34} \text{ J s}$ Speed of light in vacuum, $c = 2.998 \times 10^8 \text{ m s}^{-1}$ Rydberg constant, $E_R = 2.18 \times 10^{-18} \text{ J}$ Boltzmann constant, $k_B = 1.381 \times 10^{-23} \text{ J K}^{-1}$ Permittivity of a vacuum, $\epsilon_0 = 8.854 \times 10^{-12} \text{ C}^2 \text{ J}^{-1} \text{ m}^{-1}$ Gas constant, $R = 8.314 \text{ J K}^{-1} \text{ mol}^{-1}$
 $= 0.08206 \text{ L atm K}^{-1} \text{ mol}^{-1}$ Charge of electron, $e = 1.602 \times 10^{-19} \text{ C}$ Mass of electron, $m_e = 9.1094 \times 10^{-31} \text{ kg}$ Mass of proton, $m_p = 1.6726 \times 10^{-27} \text{ kg}$ Mass of neutron, $m_n = 1.6749 \times 10^{-27} \text{ kg}$ *Properties of matter*

Volume of 1 mole of ideal gas at 1 atm and 25 °C = 24.5 L

Volume of 1 mole of ideal gas at 1 atm and 0 °C = 22.4 L

Density of water at 298 K = 0.997 g cm⁻³*Conversion factors*

1 atm = 760 mmHg = 101.3 kPa = 1.013 bar

1 Ci = 3.70×10^{10} Bq

0 °C = 273 K

1 Hz = 1 s⁻¹1 L = 10⁻³ m³1 tonne = 10³ kg1 Å = 10⁻¹⁰ m1 W = 1 J s⁻¹1 eV = 1.602 × 10⁻¹⁹ J*Decimal fractions*

Fraction	Prefix	Symbol
10 ⁻³	milli	m
10 ⁻⁶	micro	μ
10 ⁻⁹	nano	n
10 ⁻¹²	pico	p

Decimal multiples

Multiple	Prefix	Symbol
10 ³	kilo	k
10 ⁶	mega	M
10 ⁹	giga	G
10 ¹²	tera	T

CHEM1611 - CHEMISTRY 1A (PHARMACY)*Standard Reduction Potentials, E°*

Reaction	E° / V
$\text{Co}^{3+}(\text{aq}) + \text{e}^- \rightarrow \text{Co}^{2+}(\text{aq})$	+1.82
$\text{Ce}^{4+}(\text{aq}) + \text{e}^- \rightarrow \text{Ce}^{3+}(\text{aq})$	+1.72
$\text{MnO}_4^-(\text{aq}) + 8\text{H}^+(\text{aq}) + 5\text{e}^- \rightarrow \text{Mn}^{2+}(\text{aq}) + 4\text{H}_2\text{O}$	+1.51
$\text{Au}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Au}(\text{s})$	+1.50
$\text{Cl}_2 + 2\text{e}^- \rightarrow 2\text{Cl}^-(\text{aq})$	+1.36
$\text{O}_2 + 4\text{H}^+(\text{aq}) + 4\text{e}^- \rightarrow 2\text{H}_2\text{O}$	+1.23
$\text{Pt}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Pt}(\text{s})$	+1.18
$\text{MnO}_2(\text{s}) + 4\text{H}^+(\text{aq}) + \text{e}^- \rightarrow \text{Mn}^{3+} + 2\text{H}_2\text{O}$	+0.96
$\text{NO}_3^-(\text{aq}) + 4\text{H}^+(\text{aq}) + 3\text{e}^- \rightarrow \text{NO}(\text{g}) + 2\text{H}_2\text{O}$	+0.96
$\text{Pd}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Pd}(\text{s})$	+0.92
$\text{NO}_3^-(\text{aq}) + 10\text{H}^+(\text{aq}) + 8\text{e}^- \rightarrow \text{NH}_4^+(\text{aq}) + 3\text{H}_2\text{O}$	+0.88
$\text{Ag}^+(\text{aq}) + \text{e}^- \rightarrow \text{Ag}(\text{s})$	+0.80
$\text{Fe}^{3+}(\text{aq}) + \text{e}^- \rightarrow \text{Fe}^{2+}(\text{aq})$	+0.77
$\text{Cu}^+(\text{aq}) + \text{e}^- \rightarrow \text{Cu}(\text{s})$	+0.53
$\text{Cu}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Cu}(\text{s})$	+0.34
$\text{BiO}^+(\text{aq}) + 2\text{H}^+(\text{aq}) + 3\text{e}^- \rightarrow \text{Bi}(\text{s}) + \text{H}_2\text{O}$	+0.32
$\text{Sn}^{4+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Sn}^{2+}(\text{aq})$	+0.15
$2\text{H}^+(\text{aq}) + 2\text{e}^- \rightarrow \text{H}_2(\text{g})$	0 (by definition)
$\text{Fe}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Fe}(\text{s})$	-0.04
$\text{Pb}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Pb}(\text{s})$	-0.126
$\text{Sn}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Sn}(\text{s})$	-0.136
$\text{Ni}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Ni}(\text{s})$	-0.24
$\text{Co}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Co}(\text{s})$	-0.28
$\text{Cd}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Cd}(\text{s})$	-0.40
$\text{Fe}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Fe}(\text{s})$	-0.44
$\text{Cr}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Cr}(\text{s})$	-0.74
$\text{Zn}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Zn}(\text{s})$	-0.76
$2\text{H}_2\text{O} + 2\text{e}^- \rightarrow \text{H}_2(\text{g}) + 2\text{OH}^-(\text{aq})$	-0.83
$\text{Cr}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Cr}(\text{s})$	-0.89
$\text{Al}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Al}(\text{s})$	-1.68
$\text{Sc}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Sc}(\text{s})$	-2.09
$\text{Mg}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Mg}(\text{s})$	-2.36
$\text{Na}^+(\text{aq}) + \text{e}^- \rightarrow \text{Na}(\text{s})$	-2.71
$\text{Ca}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Ca}(\text{s})$	-2.87
$\text{Li}^+(\text{aq}) + \text{e}^- \rightarrow \text{Li}(\text{s})$	-3.04

CHEM1611 - CHEMISTRY 1A (PHARMACY)

Useful formulas

<p>Quantum Chemistry</p> $E = h\nu = hc/\lambda$ $\lambda = h/mv$ $E = -Z^2 E_R(1/n^2)$ $\Delta x \cdot \Delta(mv) \geq h/4\pi$ $q = 4\pi r^2 \times 5.67 \times 10^{-8} \times T^4$ $T\lambda = 2.898 \times 10^6 \text{ K nm}$	<p>Electrochemistry</p> $\Delta G^\circ = -nFE^\circ$ $\text{Moles of } e^- = It/F$ $E = E^\circ - (RT/nF) \times \ln Q$ $E^\circ = (RT/nF) \times \ln K$ $E = E^\circ - \frac{0.0592}{n} \log Q \text{ (at } 25^\circ \text{C)}$
<p>Acids and Bases</p> $\text{pH} = -\log[\text{H}^+]$ $\text{p}K_w = \text{pH} + \text{pOH} = 14.00$ $\text{p}K_w = \text{p}K_a + \text{p}K_b = 14.00$ $\text{pH} = \text{p}K_a + \log\{[\text{A}^-] / [\text{HA}]\}$	<p>Gas Laws</p> $PV = nRT$ $(P + n^2a/V^2)(V - nb) = nRT$ $E_k = \frac{1}{2}mv^2$
<p>Radioactivity</p> $t_{1/2} = \ln 2 / \lambda$ $A = \lambda N$ $\ln(N_0/N_t) = \lambda t$ $^{14}\text{C age} = 8033 \ln(A_0/A_t) \text{ years}$	<p>Kinetics</p> $t_{1/2} = \ln 2 / k$ $k = Ae^{-E_a/RT}$ $\ln[A] = \ln[A]_0 - kt$ $\ln \frac{k_2}{k_1} = \frac{E_a}{R} \left(\frac{1}{T_1} - \frac{1}{T_2} \right)$
<p>Colligative Properties & Solutions</p> $\Pi = cRT$ $P_{\text{solution}} = X_{\text{solvent}} \times P^\circ_{\text{solvent}}$ $c = kp$ $\Delta T_f = K_f m$ $\Delta T_b = K_b m$	<p>Thermodynamics & Equilibrium</p> $\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$ $\Delta G = \Delta G^\circ + RT \ln Q$ $\Delta G^\circ = -RT \ln K$ $\Delta_{\text{univ}} S^\circ = R \ln K$ $K_p = K_c \left(\frac{RT}{100} \right)^{\Delta n}$
<p>Miscellaneous</p> $A = -\log \frac{I}{I_0}$ $A = \epsilon cl$ $E = -A \frac{e^2}{4\pi\epsilon_0 r} N_A$	<p>Mathematics</p> <p>If $ax^2 + bx + c = 0$, then $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$</p> $\ln x = 2.303 \log x$ <p>Area of circle = πr^2</p> <p>Surface area of sphere = $4\pi r^2$</p>

PERIODIC TABLE OF THE ELEMENTS

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
1 HYDROGEN H 1.008	2 HELIUM He 4.003	3 LITHIUM Li 6.941	4 BERYLLIUM Be 9.012	5 BORON B 10.81	6 CARBON C 12.01	7 NITROGEN N 14.01	8 OXYGEN O 16.00	9 FLUORINE F 19.00	10 NEON Ne 20.18	11 SODIUM Na 22.99	12 MAGNESIUM Mg 24.31	13 ALUMINUM Al 26.98	14 SILICON Si 28.09	15 PHOSPHORUS P 30.97	16 SULFUR S 32.07	17 CHLORINE Cl 35.45	18 ARGON Ar 39.95
19 POTASSIUM K 39.10	20 CALCIUM Ca 40.08	21 SCANDIUM Sc 44.96	22 TITANIUM Ti 47.88	23 VANADIUM V 50.94	24 CHROMIUM Cr 52.00	25 MANGANESE Mn 54.94	26 IRON Fe 55.85	27 COBALT Co 58.93	28 NICKEL Ni 58.69	29 COPPER Cu 63.55	30 ZINC Zn 65.39	31 GALLIUM Ga 69.72	32 GERMANIUM Ge 72.59	33 ARSENIC As 74.92	34 SELENIUM Se 78.96	35 BROMINE Br 79.90	36 KRYPTON Kr 83.80
37 BROMINE Rb 85.47	38 STRONTIUM Sr 87.62	39 YTRBIUM Y 88.91	40 ZIRCONIUM Zr 91.22	41 NIOBIUM Nb 92.91	42 MOLYBDENUM Mo 95.94	43 TECHNETIUM Tc [98.91]	44 RHENIUM Ru 101.07	45 RHODIUM Rh 102.91	46 PALLADIUM Pd 106.4	47 SILVER Ag 107.87	48 Cadmium Cd 112.40	49 INDIUM In 114.82	50 TIN Sn 118.69	51 ANTIMONY Sb 121.75	52 TELLURIUM Te 127.60	53 IODINE I 126.90	54 XENON Xe 131.30
55 CAESIUM Cs 132.91	56 BARIUM Ba 137.34	57-71 LANTHANOID S	72 HAFNIUM Hf 178.49	73 TANTALUM Ta 180.95	74 TUNGSTEN W 183.85	75 RHENIUM Re 186.2	76 OSMIUM Os 190.2	77 IRIDIUM Ir 192.22	78 PLATINUM Pt 195.09	79 GOLD Au 196.97	80 MERCURY Hg 200.59	81 THALLIUM Tl 204.37	82 LEAD Pb 207.2	83 BISMUTH Bi 208.98	84 POLONIUM Po [210.0]	85 ASTATINE At [210.0]	86 RADON Rn [222.0]
87 FRANCIUM Fr [223.0]	88 RADIUM Ra [226.0]	89-103 ACTINOID S	104 RIFERBERORIUM Rf [263]	105 DUBNIUM Db [268]	106 SEABERGIUM Sg [271]	107 BOHRIUM Bh [274]	108 HASSIUM Hs [270]	109 MEITNERIUM Mt [278]	110 DARSTADTIUM Ds [281]	111 ROSTENIUM Rg [281]	112 COOPERIUM Cn [285]	113 FLEROVIUM Fl [289]	114 LIVERMORIUM Lv [293]				

57	58	59	60	61	62	63	64	65	66	67	68	69	70	71
LANTHANUM La 138.91	CERBIUM Ce 140.12	PRASEODYMIUM Pr 140.91	NEODYMIUM Nd 144.24	PROMETHIUM Pm [144.91]	SAMARIUM Sm 150.4	EUROPIUM Eu 151.96	GADOLINIUM Gd 157.25	TERBIUM Tb 158.93	DYSPROSIUM Dy 162.50	HOLIUM Ho 164.93	ERBIUM Er 167.26	THULIUM Tm 168.93	YTERBIUM Yb 173.04	LUTETIUM Lu 174.97
89 ACTINIUM Ac [227.0]	90 THORIUM Th 232.04	91 PROTACTINIUM Pa [231.0]	92 URANIUM U 238.03	93 NEPTUNIUM Np [237.0]	94 PLUTONIUM Pu [239.1]	95 AMERICIUM Am [243.1]	96 CURIUM Cm [247.1]	97 BERKELIUM Bk [247.1]	98 CALIFORNIUM Cf [252.1]	99 EINSTEINIUM Es [252.1]	100 FERMIUM Fm [257.1]	101 MENDELEVIUM Md [256.1]	102 NOBELIUM No [259.1]	103 LAWRENCIUM Lr [260.1]