

## CHEM1612 (Pharmacy) - November 2008

### 2008-N-2

- $-14 \text{ kJ mol}^{-1}$   
Reaction is non-spontaneous ( $\Delta G^\circ = +3.2 \text{ kJ mol}^{-1}$ )

### 2008-N-3

- $567 \text{ }^\circ\text{C}$

### 2008-N-4

- $0.578 \text{ M}$   
 $291 \text{ m}$

### 2008-N-5

- $2.23 \times 10^{-5}$   
 $51.2 \text{ kJ mol}^{-1}$

### 2008-N-6

$0.0139 \text{ M}$   
 $7.82 \times 10^{-10} \text{ M}$

### 2008-N-7

- $2.09$   
 $0.28 \text{ g}$

Cl is a much larger atom than F and is less electronegative. The H–Cl bond is therefore much longer and weaker than the H–F bond. The H–Cl bond is therefore easier to break and it is the stronger acid.

HF is actually a weak acid. F is smaller and more electronegative than O, so the H–F bond is stronger than the O–H bond. There is consequently little dissociation of HF when it is dissolved in water.

### 2008-N-8

- $1.8 \times 10^{-5} \text{ M}$   
no change  
decrease

**2008-N-9**

- 1.47 V

$$5.9 \times 10^{52}$$

$$-301 \text{ kJ mol}^{-1}$$

Reaction is spontaneous as  $\Delta G^\circ$  is negative




**2008-N-10**

- 186 years
- $1.2 \times 10^{-5} \text{ M}$

**2008-N-11**


- Soap molecules consist of a long hydrophobic tail and a charged hydrophilic head. The molecules are able to form micelles (see diagram) in which the tails interact with the oil particles and the heads interact with the water. In this way, the oil is dissolved in the water and can be removed.



- mass of Na = 1.8 g      volume of  $\text{Cl}_2(\text{g}) = 0.98 \text{ L}$

### 2008-N-12

•



• From Step 1:  $K = [\text{N}_2\text{O}_4(\text{g})]/[\text{NO}_2(\text{g})]^2$   
 $\Rightarrow [\text{N}_2\text{O}_4(\text{g})] = K [\text{NO}_2(\text{g})]^2$

From Step 2:  $\text{Rate} = k [\text{N}_2\text{O}_4(\text{g})]$   
 $= k K [\text{NO}_2(\text{g})]^2$

which is consistent with the experimental result.

### 2008-N-13

- 0.06 M
- $\text{Cr}_2\text{O}_7^{2-}(\text{aq}) + 6\text{Fe}^{2+}(\text{aq}) + 14\text{H}^+(\text{aq}) \rightarrow 6\text{Fe}^{3+}(\text{aq}) + 2\text{Cr}^{3+}(\text{aq}) + 7\text{H}_2\text{O}(\text{l})$   
It would decrease.

### 2008-N-14

- $\text{Rate} = k[\text{NO}][\text{O}_3]$   
 $3.6 \times 10^6 \text{ M}^{-1} \text{ s}^{-1}$
- The small size of the colloidal particles means that they have a very large total surface area. The colloid can be stabilised by steric and/or electrostatic effects. If surface interactions are unfavourable, they are minimised by flocculation and coagulation.