

22/09(a)

The University of Sydney

CHEM1907 - CHEMISTRY 1 LIFE SCIENCES A MOLECULAR (ADVANCED)

CHEM1908 - CHEMISTRY 1 LIFE SCIENCES A (ADVANCED)

FIRST SEMESTER EXAMINATION

CONFIDENTIAL

JUNE 2005

TIME ALLOWED: THREE HOURS

GIVE THE FOLLOWING INFORMATION IN BLOCK LETTERS

FAMILY NAME		SID NUMBER	
OTHER NAMES		TABLE NUMBER	

INSTRUCTIONS TO CANDIDATES

- All questions are to be attempted. There are 16 pages of examinable material.
- Complete the examination paper in **INK**.
- Read each question carefully. Report the appropriate answer and show all relevant working in the space provided.
- The total score for this paper is 100. The possible score per page is shown in the adjacent tables.
- Each new short answer question begins with a •.
- Electronic calculators, including programmable calculators, may be used. Students are warned, however, that credit may not be given, even for a correct answer, where there is insufficient evidence of the working required to obtain the solution. Logarithms may also be used.
- Numerical values required for any question as well as a Periodic Table are printed on a separate data sheet.
- Pages 12, 15 and 20 are for rough work only.

OFFICIAL USE ONLY

Multiple choice section

	Marks	
Pages	Max	Gained
2-8	39	

Short answer section

Page	Marks		Marker
	Max	Gained	
9	11		
10	4		
11	8		
13	13		
14	4		
16	6		
17	3		
18	8		
19	4		
Total	61		
Check Total			

- Complete the following table. Give, as required, the formula, the systematic name, the oxidation number of the underlined atom and, where indicated, the number of *d* electrons for the element in this oxidation state.

Marks
5

FORMULA	SYSTEMATIC NAME	OXIDATION NUMBER	NUMBER OF <i>d</i> ELECTRONS
$\underline{\text{S}}\text{O}_3$			
$\text{K}\underline{\text{Mn}}\text{O}_4$			
$\underline{\text{Co}}\text{Cl}_2 \cdot 6\text{H}_2\text{O}$			
	ammonium sulfate		

- Draw the Lewis structures, showing all valence electrons for the following species. Indicate which of the species have contributing resonance structures.

4

NO_3^-	CO_2	N_2H_2
Resonance: YES / NO	Resonance: YES / NO	Resonance: YES / NO

- A sample of carboxypeptidase (an enzyme) was purified and found on analysis to contain 0.191% by weight of zinc. What is the *minimum* molecular weight of the enzyme if we assume it is a monomer?

2

Answer:

- Solutions of lead(II) nitrate (0.080 M, 60 mL) and potassium iodide (0.080 M, 40 mL) are mixed. What amount (in mol) of $\text{PbI}_2(\text{s})$ precipitates?

Marks
4

Answer:



What is the final concentration of $\text{K}^+(\text{aq})$ ions remaining in solution after the reaction?

Answer:

THE REMAINDER OF THIS PAGE IS FOR ROUGH WORKING ONLY

- Shown here are the classical and the zwitterionic forms of the amino acid leucine.

Marks
8



List the types of intermolecular interactions in which each of the indicated sites (**A**, **B** and **C**) in leucine could be involved.

A


B

C

Provide the requested information for each of the indicated atoms in leucine.









Atom	Geometric arrangement of the electron pairs around the atom	Hybridisation of the atom	Geometry/shape of σ -bonding electron pairs around the atom
¹ C			
² C			
³ O			

Given that the pK_a of the carboxylic acid group of leucine is 2.32 and the pK_b of the amine group is 4.24, do you expect the classical or the zwitterionic form to predominate when leucine is dissolved in water? In other words, does the following equilibrium lie to the right or left? Show your reasoning.




Marks
13

- Complete the following table. Make sure you complete the name of the starting material where indicated.


STARTING MATERIAL NAME (where required)	REAGENTS/ CONDITIONS	CONSTITUTIONAL FORMULA(S) OF MAJOR ORGANIC PRODUCT(S)
$\text{CH}_3\text{CH}_2\underset{\text{Br}}{\text{CH}}\text{CH}_2\text{CH}_3$ Name:		$\text{CH}_3\text{CH}_2\underset{\text{Br}^- \oplus \text{N}(\text{CH}_3)_3}{\text{CH}}\text{CH}_2\text{CH}_3$
$\text{CH}_3\text{CH}_2\text{CH}_2\overset{\text{O}}{\parallel}{\text{C}}\text{CH}_3$ Name:	1. LiAlH_4 / dry ether 2. H^+ / H_2O	
 Name:	H_2 / Pd catalyst	
	3 M NaOH / heat	
		
	conc. HCl / heat	
	$[\text{Ag}(\text{NH}_3)_2]^+$ / dil. OH^-	
		

- Two of the following compounds are bases and two are not. Identify the two bases and explain, with the aid of diagrams, why they react with acids and why the other two compounds do not.


Marks
4




pyridine



pyrrole




aniline




acetanilide

- Consider the structures of the two aldohexoses shown below.

Marks
6



β -D-allopyranose



α -D-altropyranose

Draw the Fischer projection of the open chain forms of sugars (A) and (B).

(A)	(B)
-----	-----

Give the products obtained when D-allose is treated with the following reagents.

acidified methanol	NaBH_4 in methanol solvent
--------------------	-------------------------------------

In the case of the product obtained from the reaction with NaBH_4 in methanol, specify the optical rotation as "(+)", "(-)", "zero" or "cannot be predicted". Justify your choice.

THIS QUESTION IS CONTINUED ON THE NEXT PAGE.


Draw the Haworth structure of β -D-allopyranosyl- α -D-altropyranoside.

Marks
3


Will this disaccharide be a reducing sugar? Explain your answer.

THE REMAINDER OF THIS PAGE IS FOR ROUGH WORKING ONLY


- Consider the three amino acids shown below.



threonine (Thr)



aspartic acid (Asp)




lysine (Lys)

Draw the constitutional formulas for the following dipeptides present in water solution at the indicated pH values.

Marks
8

Thr-Lys, pH 1.0	Asp-Thr, pH 12.0
-----------------	------------------

Draw the stereoformula for L-threonine and indicate its absolute configuration using the (*R*)- and (*S*)- convention.

	Absolute configuration <div style="display: inline-block; border: 1px solid black; width: 40px; height: 40px; vertical-align: middle; margin-left: 10px;"></div>	
L-threonine		

The pK_a values of aspartic acid are 1.88 (α -COOH), 9.60 (α -NH₃[⊕]) and 3.65 ($-\text{CH}_2\text{COOH}$). Draw the zwitterionic form of aspartic acid.

--


At what pH will this be the predominant species in aqueous solution?

THIS QUESTION IS CONTINUED ON THE NEXT PAGE.

Give the products obtained when lysine is treated with the following reagents.

Marks
4

excess methanol / HCl



THE REMAINDER OF THIS PAGE IS FOR ROUGH WORKING ONLY

CHEM1907 - CHEMISTRY 1 LIFE SCIENCES A MOLECULAR (ADVANCED)**CHEM1908 - CHEMISTRY 1 LIFE SCIENCES A (ADVANCED)****DATA SHEET***Physical constants*Avogadro constant, $N_A = 6.022 \times 10^{23} \text{ mol}^{-1}$ Faraday constant, $F = 96485 \text{ C mol}^{-1}$ Planck constant, $h = 6.626 \times 10^{-34} \text{ J s}$ Speed of light in vacuum, $c = 2.998 \times 10^8 \text{ m s}^{-1}$ Rydberg constant, $E_R = 2.18 \times 10^{-18} \text{ J}$ Boltzmann constant, $k_B = 1.381 \times 10^{-23} \text{ J K}^{-1}$ Gas constant, $R = 8.314 \text{ J K}^{-1} \text{ mol}^{-1}$
 $= 0.08206 \text{ L atm K}^{-1} \text{ mol}^{-1}$ Charge of electron, $e = 1.602 \times 10^{-19} \text{ C}$ Mass of electron, $m_e = 9.1094 \times 10^{-31} \text{ kg}$ Mass of proton, $m_p = 1.6726 \times 10^{-27} \text{ kg}$ Mass of neutron, $m_n = 1.6749 \times 10^{-27} \text{ kg}$ *Properties of matter*

Volume of 1 mole of ideal gas at 1 atm and 25 °C = 24.5 L

Volume of 1 mole of ideal gas at 1 atm and 0 °C = 22.4 L

Density of water at 298 K = 0.997 g cm⁻³*Conversion factors*

1 atm = 760 mmHg = 101.3 kPa

0 °C = 273 K

1 L = 10⁻³ m³1 Å = 10⁻¹⁰ m1 eV = 1.602 × 10⁻¹⁹ J1 Ci = 3.70 × 10¹⁰ Bq1 Hz = 1 s⁻¹*Decimal fractions*

Fraction	Prefix	Symbol
10 ⁻³	milli	m
10 ⁻⁶	micro	μ
10 ⁻⁹	nano	n
10 ⁻¹²	pico	p

Decimal multiples

Multiple	Prefix	Symbol
10 ³	kilo	k
10 ⁶	mega	M
10 ⁹	giga	G

CHEM1907 - CHEMISTRY 1 LIFE SCIENCES A MOLECULAR (ADVANCED)**CHEM1908 - CHEMISTRY 1 LIFE SCIENCES A (ADVANCED)***Standard Reduction Potentials, E°*

Reaction	E° / V
$\text{Co}^{3+}(\text{aq}) + \text{e}^- \rightarrow \text{Co}^{2+}(\text{aq})$	+1.82
$\text{Ce}^{4+}(\text{aq}) + \text{e}^- \rightarrow \text{Ce}^{3+}(\text{aq})$	+1.72
$\text{Cl}_2(\text{g}) + 2\text{e}^- \rightarrow 2\text{Cl}^-(\text{aq})$	+1.36
$\text{O}_2(\text{g}) + 4\text{H}^+(\text{aq}) + 4\text{e}^- \rightarrow 2\text{H}_2\text{O}(\text{l})$	+1.23
$\text{Pd}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Pd}(\text{s})$	+0.92
$\text{Ag}^+(\text{aq}) + \text{e}^- \rightarrow \text{Ag}(\text{s})$	+0.80
$\text{Fe}^{3+}(\text{aq}) + \text{e}^- \rightarrow \text{Fe}^{2+}(\text{aq})$	+0.77
$\text{Cu}^+(\text{aq}) + \text{e}^- \rightarrow \text{Cu}(\text{s})$	+0.53
$\text{Cu}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Cu}(\text{s})$	+0.34
$\text{Sn}^{4+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Sn}^{2+}(\text{aq})$	+0.15
$2\text{H}^+(\text{aq}) + 2\text{e}^- \rightarrow \text{H}_2(\text{g})$	0 (by definition)
$\text{Fe}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Fe}(\text{s})$	-0.04
$\text{Pb}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Pb}(\text{s})$	-0.13
$\text{Sn}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Sn}(\text{s})$	-0.14
$\text{Ni}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Ni}(\text{s})$	-0.24
$\text{Co}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Co}(\text{s})$	-0.28
$\text{Fe}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Fe}(\text{s})$	-0.44
$\text{Cr}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Cr}(\text{s})$	-0.74
$\text{Zn}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Zn}(\text{s})$	-0.76
$2\text{H}_2\text{O}(\text{l}) + 2\text{e}^- \rightarrow \text{H}_2(\text{g}) + 2\text{OH}^-(\text{aq})$	-0.83
$\text{Cr}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Cr}(\text{s})$	-0.89
$\text{Al}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Al}(\text{s})$	-1.68
$\text{Mg}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Mg}(\text{s})$	-2.36
$\text{Na}^+(\text{aq}) + \text{e}^- \rightarrow \text{Na}(\text{s})$	-2.71

CHEM1907 - CHEMISTRY 1 LIFE SCIENCES A MOLECULAR (ADVANCED)**CHEM1908 - CHEMISTRY 1 LIFE SCIENCES A (ADVANCED)***Useful formulas*

<p>Quantum Chemistry</p> $E = h\nu = hc/\lambda$ $\lambda = h/mv$ $4.5k_B T = hc/\lambda$ $E = Z^2 E_R (1/n^2)$	<p>Radioactivity</p> $t_{1/2} = \ln 2 / \lambda$ $A = \lambda N$ $\ln(N_0/N_t) = \lambda t$ $^{14}\text{C age} = 8033 \ln(A_0/A_t)$
<p>Acids and Bases</p> $pK_w = \text{pH} + \text{pOH} = 14.00$ $pK_w = \text{p}K_a + \text{p}K_b = 14.00$ $\text{pH} = \text{p}K_a + \log\{[A^-] / [\text{HA}] \}$	<p>Gas Laws</p> $PV = nRT$ $(P + n^2 a/V^2)(V - nb) = nRT$
<p>Colligative properties</p> $\pi = cRT$ $P_{\text{solution}} = X_{\text{solvent}} \times P^{\circ}_{\text{solvent}}$ $p = kc$ $\Delta T_f = K_f m$ $\Delta T_b = K_b m$	<p>Kinetics</p> $t_{1/2} = \ln 2 / k$ $k = A e^{-E_a/RT}$ $\ln[A] = \ln[A]_0 - kt$ $\ln \frac{k_2}{k_1} = \frac{E_a}{R} \left(\frac{1}{T_1} - \frac{1}{T_2} \right)$
<p>Electrochemistry</p> $\Delta G^{\circ} = -nFE^{\circ}$ <p>Moles of $e^- = It/F$</p> $E = E^{\circ} - (RT/nF) \times 2.303 \log Q$ $= E^{\circ} - (RT/nF) \times \ln Q$ $E^{\circ} = (RT/nF) \times 2.303 \log K$ $= (RT/nF) \times \ln K$ $E = E^{\circ} - \frac{0.0592}{n} \log Q \text{ (at } 25^{\circ}\text{C)}$	<p>Thermodynamics & Equilibrium</p> $\Delta G^{\circ} = \Delta H^{\circ} - T\Delta S^{\circ}$ $\Delta G = \Delta G^{\circ} + RT \ln Q$ $\Delta G^{\circ} = -RT \ln K$ $K_p = K_c (RT)^{\Delta n}$
<p>Polymers</p> $R_g = \sqrt{\frac{nl_0^2}{6}}$	<p>Mathematics</p> <p>If $ax^2 + bx + c = 0$, then $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$</p> $\ln x = 2.303 \log x$

PERIODIC TABLE OF THE ELEMENTS

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
1 HYDROGEN H 1.008																	2 HELIUM He 4.003
3 LITHIUM Li 6.941	4 BERYLLIUM Be 9.012											5 BORON B 10.81	6 CARBON C 12.01	7 NITROGEN N 14.01	8 OXYGEN O 16.00	9 FLUORINE F 19.00	10 NEON Ne 20.18
11 SODIUM Na 22.99	12 MAGNESIUM Mg 24.31											13 ALUMINIUM Al 26.98	14 SILICON Si 28.09	15 PHOSPHORUS P 30.97	16 SULFUR S 32.07	17 CHLORINE Cl 35.45	18 ARGON Ar 39.95
19 POTASSIUM K 39.10	20 CALCIUM Ca 40.08	21 SCANDIUM Sc 44.96	22 TITANIUM Ti 47.88	23 VANADIUM V 50.94	24 CHROMIUM Cr 52.00	25 MANGANESE Mn 54.94	26 IRON Fe 55.85	27 COBALT Co 58.93	28 NICKEL Ni 58.69	29 COPPER Cu 63.55	30 ZINC Zn 65.39	31 GALLIUM Ga 69.72	32 GERMANIUM Ge 72.59	33 ARSENIC As 74.92	34 SELENIUM Se 78.96	35 BROMINE Br 79.90	36 KRYPTON Kr 83.80
37 RUBIDIUM Rb 85.47	38 STRONTIUM Sr 87.62	39 YTRIUM Y 88.91	40 ZIRCONIUM Zr 91.22	41 NIObIUM Nb 92.91	42 MOLYBDENUM Mo 95.94	43 TECHNETIUM Tc [98.91]	44 RUTHENIUM Ru 101.07	45 RHODIUM Rh 102.91	46 PALLADIUM Pd 106.4	47 SILVER Ag 107.87	48 CADMIUM Cd 112.40	49 INDIUM In 114.82	50 TIN Sn 118.69	51 ANTIMONY Sb 121.75	52 TELLURIUM Te 127.60	53 IODINE I 126.90	54 XENON Xe 131.30
55 CAESIUM Cs 132.91	56 BARIUM Ba 137.34	57-71	72 HAFNIUM Hf 178.49	73 TANTALUM Ta 180.95	74 TUNGSTEN W 183.85	75 RHENIUM Re 186.2	76 OSMIUM Os 190.2	77 IRIDIUM Ir 192.22	78 PLATINUM Pt 195.09	79 GOLD Au 196.97	80 MERCURY Hg 200.59	81 THALLIUM Tl 204.37	82 LEAD Pb 207.2	83 BISMUTH Bi 208.98	84 POLONIUM Po [210.0]	85 ASTATINE At [210.0]	86 RADON Rn [222.0]
87 FRANCIUM Fr [223.0]	88 RADIUM Ra [226.0]	89-103	104 RUTHERFORDIUM Rf [261]	105 DUBNIUM Db [262]	106 SEABORGIUM Sg [266]	107 BOHRIUM Bh [262]	108 HASSIUM Hs [265]	109 MEITNERIUM Mt [266]									

LANTHANIDES

57 LANTHANUM La 138.91	58 CERIUM Ce 140.12	59 PRASEODYMIUM Pr 140.91	60 NEODYMIUM Nd 144.24	61 PROMETHIUM Pm [144.9]	62 SAMARIUM Sm 150.4	63 EUROPIUM Eu 151.96	64 GADOLINIUM Gd 157.25	65 TERBIUM Tb 158.93	66 DYSPROSIUM Dy 162.50	67 HOLMIUM Ho 164.93	68 ERBIUM Er 167.26	69 THULIUM Tm 168.93	70 YTTERBIUM Yb 173.04	71 LUTETIUM Lu 174.97
89 ACTINIUM Ac [227.0]	90 THORIUM Th 232.04	91 PROTACTINIUM Pa [231.0]	92 URANIUM U 238.03	93 NEPTUNIUM Np [237.0]	94 PLUTONIUM Pu [239.1]	95 AMERICIUM Am [243.1]	96 CURIUM Cm [247.1]	97 BERKELIUM Bk [247.1]	98 CALIFORNIUM Cf [252.1]	99 EINSTEINIUM Es [252.1]	100 FERMIUM Fm [257.1]	101 MENDELEVIUM Md [256.1]	102 NOBELIUM No [259.1]	103 LAWRENCIUM Lr [260.1]

ACTINIDES