

CHEM1908 - CHEMISTRY 1 LIFE SCIENCES A (ADVANCED)

FIRST SEMESTER EXAMINATION

CONFIDENTIAL

JUNE 2006

TIME ALLOWED: THREE HOURS

GIVE THE FOLLOWING INFORMATION IN BLOCK LETTERS

FAMILY NAME		SID NUMBER	
OTHER NAMES		TABLE NUMBER	

INSTRUCTIONS TO CANDIDATES

- All questions are to be attempted. There are 18 pages of examinable material.
- Complete the examination paper in **INK**.
- Read each question carefully. Report the appropriate answer and show all relevant working in the space provided.
- The total score for this paper is 100. The possible score per page is shown in the adjacent tables.
- Each new short answer question begins with a •.
- Electronic calculators, including programmable calculators, may be used. Students are warned, however, that credit may not be given, even for a correct answer, where there is insufficient evidence of the working required to obtain the solution. Logarithms may also be used.
- Numerical values required for any question as well as a Periodic Table are printed on a separate data sheet.
- Page 20 is for rough work only.

OFFICIAL USE ONLY

Multiple choice section

Pages	Marks	
	Max	Gained
2-11	36	

Short answer section

Page	Marks		Marker
	Max	Gained	
12	9		
13	3		
14	7		
15	5		
16	11		
17	10		
18	8		
19	11		
Total	64		
Check Total			

Marks
2

- Give the full electron configuration for the ground state K atom.

--

What are the three quantum numbers that describe the orbital that contains the electron furthest from the nucleus in the K atom?

$n =$	$l =$	$m_l =$
-------	-------	---------

4

- Draw the Lewis structures, showing all valence electrons for the following species. Indicate which of the species have contributing resonance structures.

NCO^-	COF_2	NO_3^-
Resonance: YES / NO	Resonance: YES / NO	Resonance: YES / NO

3

- Human haemoglobin has a molar weight of $6.45 \times 10^4 \text{ g mol}^{-1}$ and contains 3.46 g of iron per kg. Calculate the number of iron atoms in each molecule of haemoglobin.

--

Answer:

- If 50 mL of a 0.10 M solution of AgNO_3 is mixed with 50 mL of a 0.040 M solution of BaCl_2 , what mass of $\text{AgCl}(s)$ will precipitate from the reaction?

Marks
3

Answer:


What is the concentration of NO_3^- ions in the final solution from the reaction above?

Answer:

THE REMAINDER OF THIS PAGE IS FOR ROUGH WORKING ONLY

- Tranexamic acid, *trans*-(4-aminomethyl)cyclohexanecarboxylic acid, is used for the treatment of severe haemorrhage in patients with haemophilia.

Marks
4



Provide the requested information for each of the indicated atoms in tranexamic acid.

Atom	Geometric arrangement of the electron pairs around the atom	Hybridisation of the atom	Geometry/shape of σ -bonding electron pairs around the atom
C-1			
C-2			
O-3			
N-4			

- Consider the boiling points of the compounds 1-propanol, 1-propanethiol and 1-propaneselenol shown in the table below?


3

Compound	CH ₃ CH ₂ CH ₂ OH	CH ₃ CH ₂ CH ₂ SH	CH ₃ CH ₂ CH ₂ SeH
Boiling point (° C)	97.2	67.8	147.0

With reference to intermolecular forces, explain briefly why the boiling points increase in the order CH₃CH₂CH₂SH < CH₃CH₂CH₂OH < CH₃CH₂CH₂SeH.

Marks
5

- Consider the following equation.



Name all of the species in this equation.

HBrO

BrO⁻NH₃NH₄⁺










Complete the following table by giving the correct pK_a or pK_b value where it can be calculated. Mark with a cross (✗) those cells for which insufficient data have been given to calculate a value.

Species	HBrO	NH ₃	BrO ⁻	NH ₄ ⁺
pK_a of acid	8.64			
pK_b of base		4.76		


Determine on which side (left or right hand side) the equilibrium for the reaction above will lie. Provide a brief rationale for your answer.

Marks
11

- Complete the following table. Make sure you complete the name of the starting material where indicated.

STARTING MATERIAL NAME (where required)	REAGENTS/ CONDITIONS	CONSTITUTIONAL FORMULA(S) OF MAJOR ORGANIC PRODUCT(S)
 Name:		
 Name:	Br_2 $(\text{CCl}_4 \text{ solvent})$	
 Name:		
	$3 \text{ M NaOH} / \text{heat}$	
		
	$1 \text{ M HCl} / \text{heat}$	

- Trifluridine is an analogue of the nucleoside thymidine and is used clinically as an anti-viral agent. It differs from thymidine in that the methyl group is replaced by a trifluoromethyl group.



Marks
6

Give the molecular formula of trifluridine.

Classify the sugar present in trifluridine as a furanose or pyranose.


Is the sugar present as the α -anomer or β -anomer?

Give an example of a nucleotide derived from trifluridine.

Hydrolysis of trifluridine gives the sugar 2-deoxyribose and a base. Give the structure of the base and the structure of one of its tautomers.

base	tautomer


- Show clearly the reagents you would use to carry out the following chemical conversion. Draw constitutional formulas for any intermediate compounds. Note: More than one step is required.



4

Marks
8

- At equilibrium in aqueous solution, D-xylose exists as a mixture containing the α -pyranose, β -pyranose, α -furanose and β -furanose forms. Draw Haworth formulas in the appropriate boxes below for each of these forms.




Give the formula in Fischer projection, of the product formed when D-xylose is treated with $[\text{Ag}(\text{NH}_3)_2]^+$ / dilute OH^- .

Give the stereoformula of the product(s) formed when β -D-xylopyranose is heated with methanol and an acid catalyst.

**Marks
11**

- The constitutional formula of the naturally occurring tetrapeptide, Tyr-Lys-Ser-Asn, is shown below.



Give the Fischer projection of L-Lys as the zwitterion.	Complete the stereoformula of (<i>S</i>)-Ser.

Give the constitutional formulas in the correct ionic states of the products obtained from the vigorous acidic hydrolysis (6 M HCl) of the tetrapeptide.

--

Give the constitutional formulas for the following dipeptides present in water at the indicated pH values.

Tyr-Ser at pH 12	Asn-Lys at pH 1

CHEM1908 - CHEMISTRY 1 LIFE SCIENCES A (ADVANCED)**DATA SHEET***Physical constants*Avogadro constant, $N_A = 6.022 \times 10^{23} \text{ mol}^{-1}$ Faraday constant, $F = 96485 \text{ C mol}^{-1}$ Planck constant, $h = 6.626 \times 10^{-34} \text{ J s}$ Speed of light in vacuum, $c = 2.998 \times 10^8 \text{ m s}^{-1}$ Rydberg constant, $E_R = 2.18 \times 10^{-18} \text{ J}$ Boltzmann constant, $k_B = 1.381 \times 10^{-23} \text{ J K}^{-1}$ Gas constant, $R = 8.314 \text{ J K}^{-1} \text{ mol}^{-1}$
 $= 0.08206 \text{ L atm K}^{-1} \text{ mol}^{-1}$ Charge of electron, $e = 1.602 \times 10^{-19} \text{ C}$ Mass of electron, $m_e = 9.1094 \times 10^{-31} \text{ kg}$ Mass of proton, $m_p = 1.6726 \times 10^{-27} \text{ kg}$ Mass of neutron, $m_n = 1.6749 \times 10^{-27} \text{ kg}$ *Properties of matter*

Volume of 1 mole of ideal gas at 1 atm and 25 °C = 24.5 L

Volume of 1 mole of ideal gas at 1 atm and 0 °C = 22.4 L

Density of water at 298 K = 0.997 g cm^{-3} *Conversion factors*

1 atm = 760 mmHg = 101.3 kPa

0 °C = 273 K

1 L = 10^{-3} m^3 1 Å = 10^{-10} m 1 eV = $1.602 \times 10^{-19} \text{ J}$ 1 Ci = $3.70 \times 10^{10} \text{ Bq}$ 1 Hz = 1 s^{-1} *Decimal fractions*

Fraction	Prefix	Symbol
10^{-3}	milli	m
10^{-6}	micro	μ
10^{-9}	nano	n
10^{-12}	pico	p

Decimal multiples

Multiple	Prefix	Symbol
10^3	kilo	k
10^6	mega	M
10^9	giga	G

CHEM1908 - CHEMISTRY 1 LIFE SCIENCES A (ADVANCED)*Standard Reduction Potentials, E°*

Reaction	E° / V
$\text{Co}^{3+}(\text{aq}) + \text{e}^{-} \rightarrow \text{Co}^{2+}(\text{aq})$	+1.82
$\text{Ce}^{4+}(\text{aq}) + \text{e}^{-} \rightarrow \text{Ce}^{3+}(\text{aq})$	+1.72
$\text{Au}^{3+}(\text{aq}) + 3\text{e}^{-} \rightarrow \text{Au}(\text{s})$	+1.50
$\text{Cl}_2 + 2\text{e}^{-} \rightarrow 2\text{Cl}^{-}(\text{aq})$	+1.36
$\text{O}_2 + 4\text{H}^{+}(\text{aq}) + 4\text{e}^{-} \rightarrow 2\text{H}_2\text{O}$	+1.23
$\text{MnO}_2(\text{s}) + 4\text{H}^{+}(\text{aq}) + \text{e}^{-} \rightarrow \text{Mn}^{3+} + 2\text{H}_2\text{O}$	+0.96
$\text{Pd}^{2+}(\text{aq}) + 2\text{e}^{-} \rightarrow \text{Pd}(\text{s})$	+0.92
$\text{Ag}^{+}(\text{aq}) + \text{e}^{-} \rightarrow \text{Ag}(\text{s})$	+0.80
$\text{Fe}^{3+}(\text{aq}) + \text{e}^{-} \rightarrow \text{Fe}^{2+}(\text{aq})$	+0.77
$\text{Cu}^{+}(\text{aq}) + \text{e}^{-} \rightarrow \text{Cu}(\text{s})$	+0.53
$\text{Cu}^{2+}(\text{aq}) + 2\text{e}^{-} \rightarrow \text{Cu}(\text{s})$	+0.34
$\text{Sn}^{4+}(\text{aq}) + 2\text{e}^{-} \rightarrow \text{Sn}^{2+}(\text{aq})$	+0.15
$2\text{H}^{+}(\text{aq}) + 2\text{e}^{-} \rightarrow \text{H}_2(\text{g})$	0 (by definition)
$\text{Fe}^{3+}(\text{aq}) + 3\text{e}^{-} \rightarrow \text{Fe}(\text{s})$	-0.04
$\text{Pb}^{2+}(\text{aq}) + 2\text{e}^{-} \rightarrow \text{Pb}(\text{s})$	-0.13
$\text{Sn}^{2+}(\text{aq}) + 2\text{e}^{-} \rightarrow \text{Sn}(\text{s})$	-0.14
$\text{Ni}^{2+}(\text{aq}) + 2\text{e}^{-} \rightarrow \text{Ni}(\text{s})$	-0.24
$\text{Fe}^{2+}(\text{aq}) + 2\text{e}^{-} \rightarrow \text{Fe}(\text{s})$	-0.44
$\text{Cr}^{3+}(\text{aq}) + 3\text{e}^{-} \rightarrow \text{Cr}(\text{s})$	-0.74
$\text{Zn}^{2+}(\text{aq}) + 2\text{e}^{-} \rightarrow \text{Zn}(\text{s})$	-0.76
$2\text{H}_2\text{O} + 2\text{e}^{-} \rightarrow \text{H}_2(\text{g}) + 2\text{OH}^{-}(\text{aq})$	-0.83
$\text{Cr}^{2+}(\text{aq}) + 2\text{e}^{-} \rightarrow \text{Cr}(\text{s})$	-0.89
$\text{Al}^{3+}(\text{aq}) + 3\text{e}^{-} \rightarrow \text{Al}(\text{s})$	-1.68
$\text{Mg}^{2+}(\text{aq}) + 2\text{e}^{-} \rightarrow \text{Mg}(\text{s})$	-2.36
$\text{Na}^{+}(\text{aq}) + \text{e}^{-} \rightarrow \text{Na}(\text{s})$	-2.71
$\text{Ca}^{2+}(\text{aq}) + 2\text{e}^{-} \rightarrow \text{Ca}(\text{s})$	-2.87
$\text{Li}^{+}(\text{aq}) + \text{e}^{-} \rightarrow \text{Li}(\text{s})$	-3.04

CHEM1908 - CHEMISTRY 1 LIFE SCIENCES A (ADVANCED)

Useful formulas

<p>Quantum Chemistry</p> $E = h\nu = hc/\lambda$ $\lambda = h/mv$ $4.5k_B T = hc/\lambda$ $E = Z^2 E_R (1/n^2)$ $\Delta x \cdot \Delta(mv) \geq h/4\pi$ $q = 4\pi r^2 \times 5.67 \times 10^{-8} \times T^4$	<p>Electrochemistry</p> $\Delta G^\circ = -nFE^\circ$ <p>Moles of $e^- = It/F$</p> $E = E^\circ - (RT/nF) \times 2.303 \log Q$ $= E^\circ - (RT/nF) \times \ln Q$ $E^\circ = (RT/nF) \times 2.303 \log K$ $= (RT/nF) \times \ln K$ $E = E^\circ - \frac{0.0592}{n} \log Q \text{ (at 25 }^\circ\text{C)}$
<p>Acids and Bases</p> $pK_w = \text{pH} + \text{pOH} = 14.00$ $pK_w = \text{p}K_a + \text{p}K_b = 14.00$ $\text{pH} = \text{p}K_a + \log\{[A^-] / [HA]\}$	<p>Gas Laws</p> $PV = nRT$ $(P + n^2a/V^2)(V - nb) = nRT$
<p>Colligative properties</p> $\pi = cRT$ $P_{\text{solution}} = X_{\text{solvent}} \times P^\circ_{\text{solvent}}$ $p = kc$ $\Delta T_f = K_f m$ $\Delta T_b = K_b m$	<p>Kinetics</p> $t_{1/2} = \ln 2/k$ $k = Ae^{-E_a/RT}$ $\ln[A] = \ln[A]_0 - kt$ $\ln \frac{k_2}{k_1} = \frac{E_a}{R} \left(\frac{1}{T_1} - \frac{1}{T_2} \right)$
<p>Radioactivity</p> $t_{1/2} = \ln 2/\lambda$ $A = \lambda N$ $\ln(N_0/N_t) = \lambda t$ $^{14}\text{C age} = 8033 \ln(A_0/A_t)$	<p>Thermodynamics & Equilibrium</p> $\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$ $\Delta G = \Delta G^\circ + RT \ln Q$ $\Delta G^\circ = -RT \ln K$ $K_p = K_c (RT)^{\Delta n}$
<p>Polymers</p> $R_g = \sqrt{\frac{nl_0^2}{6}}$	<p>Mathematics</p> <p>If $ax^2 + bx + c = 0$, then $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$</p> $\ln x = 2.303 \log x$

PERIODIC TABLE OF THE ELEMENTS

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
1 HYDROGEN H 1.008																	2 HELIUM He 4.003
3 LITHIUM Li 6.941	4 BERYLLIUM Be 9.012											5 BORON B 10.81	6 CARBON C 12.01	7 NITROGEN N 14.01	8 OXYGEN O 16.00	9 FLUORINE F 19.00	10 NEON Ne 20.18
11 SODIUM Na 22.99	12 MAGNESIUM Mg 24.31											13 ALUMINIUM Al 26.98	14 SILICON Si 28.09	15 PHOSPHORUS P 30.97	16 SULFUR S 32.07	17 CHLORINE Cl 35.45	18 ARGON Ar 39.95
19 POTASSIUM K 39.10	20 CALCIUM Ca 40.08	21 SCANDIUM Sc 44.96	22 TITANIUM Ti 47.88	23 VANADIUM V 50.94	24 CHROMIUM Cr 52.00	25 MANGANESE Mn 54.94	26 IRON Fe 55.85	27 COBALT Co 58.93	28 NICKEL Ni 58.69	29 COPPER Cu 63.55	30 ZINC Zn 65.39	31 GALLIUM Ga 69.72	32 GERMANIUM Ge 72.59	33 ARSENIC As 74.92	34 SELENIUM Se 78.96	35 BROMINE Br 79.90	36 KRYPTON Kr 83.80
37 RUBIDIUM Rb 85.47	38 STRONTIUM Sr 87.62	39 YTRIUM Y 88.91	40 ZIRCONIUM Zr 91.22	41 NOBIUM Nb 92.91	42 MOLYBDENUM Mo 95.94	43 TECHNETIUM Tc [98.91]	44 RUTHENIUM Ru 101.07	45 RHODIUM Rh 102.91	46 PALLADIUM Pd 106.4	47 SILVER Ag 107.87	48 CADMIUM Cd 112.40	49 INDIUM In 114.82	50 TIN Sn 118.69	51 ANTIMONY Sb 121.75	52 TELLURIUM Te 127.60	53 IODINE I 126.90	54 XENON Xe 131.30
55 CAESIUM Cs 132.91	56 BARIUM Ba 137.34	57-71	72 HAFNIUM Hf 178.49	73 TANTALUM Ta 180.95	74 TUNGSTEN W 183.85	75 RHENIUM Re 186.2	76 OSMIUM Os 190.2	77 IRIDIUM Ir 192.22	78 PLATINUM Pt 195.09	79 GOLD Au 196.97	80 MERCURY Hg 200.59	81 THALLIUM Tl 204.37	82 LEAD Pb 207.2	83 BISMUTH Bi 208.98	84 POLONIUM Po [210.0]	85 ASTATINE At [210.0]	86 RADON Rn [222.0]
87 FRANCIUM Fr [223.0]	88 RADIUM Ra [226.0]	89-103	104 RUTHERFORDIUM Rf [261]	105 DUBNIUM Db [262]	106 SEABORGIUM Sg [266]	107 BOHRIUM Bh [262]	108 HASSIUM Hs [265]	109 MEITNERIUM Mt [266]									

LANTHANIDES

57 LANTHANUM La 138.91	58 CERIUM Ce 140.12	59 PRASEODYMIUM Pr 140.91	60 NEODYMIUM Nd 144.24	61 PROMETHIUM Pm [144.9]	62 SAMARIUM Sm 150.4	63 EUROPIUM Eu 151.96	64 GADOLINIUM Gd 157.25	65 TERBIUM Tb 158.93	66 DYSPROSIUM Dy 162.50	67 HOLMIUM Ho 164.93	68 ERBIUM Er 167.26	69 THULIUM Tm 168.93	70 YTTERBIUM Yb 173.04	71 LUTETIUM Lu 174.97
89 ACTINIUM Ac [227.0]	90 THORIUM Th 232.04	91 PROTACTINIUM Pa [231.0]	92 URANIUM U 238.03	93 NEPTUNIUM Np [237.0]	94 PLUTONIUM Pu [239.1]	95 AMERICIUM Am [243.1]	96 CURIUM Cm [247.1]	97 BERKELIUM Bk [247.1]	98 CALIFORNIUM Cf [252.1]	99 EINSTEINIUM Es [252.1]	100 FERMIUM Fm [257.1]	101 MENDELEVIUM Md [256.1]	102 NOBELIUM No [259.1]	103 LAWRENCIUM Lr [260.1]

ACTINIDES