

Chemistry 2

Lecture 3

Particle on a ring approximation



Learning outcomes from Lecture 2

- Be able to explain why confining a particle to a box leads to quantization of its energy levels
- Be able to explain why the lowest energy of the particle in a box is not zero
- Be able to apply the particle in a box approximation as a model for the electronic structure of a conjugated molecule (given equation for E_n).

Assumed knowledge for today

Be able to predict the number of π electrons and the presence of conjugation in a ring containing carbon and/or heteroatoms such as nitrogen and oxygen.

The de Broglie Approach

- The wavelength of the wave associated with a particle is related to its momentum:


$$p = mv = h / \lambda$$

- For a particle with only kinetic energy:

$$E = \frac{1}{2} mv^2 = p^2 / 2m = h^2 / 2m\lambda^2$$

Particle-on-a-ring

- Particle can be anywhere on ring




- Ground state is motionless



Particle-on-a-ring

- Ground state is motionless
- In higher levels, we must fit an integer number of waves around the ring



The Schrödinger equation

- The total energy is extracted by the Hamiltonian operator.
- These are the “observable” energy levels of a quantum particle

$$\hat{H}\Psi(x) = \epsilon_i\Psi(x)$$

Hamiltonian operator points to \hat{H} , Energy eigenvalue points to ϵ_i , and Energy eigenfunction points to $\Psi(x)$.



The Schrödinger equation

- The Hamiltonian has parts corresponding to *Kinetic Energy* and *Potential Energy*. In terms of the angle θ :


$$\hat{H}\Psi = \left(-\frac{\hbar^2}{2mr^2} \frac{\partial^2}{\partial \theta^2} + V(\theta) \right) \Psi$$

Hamiltonian operator Kinetic Energy Potential Energy


"The particle on a ring"

- The ring is a cyclic 1d potential

$$\Psi = \sin(j\theta) \quad \Psi = \cos(j\theta)$$



"The particle on a ring"



π -system of benzene is like a bunch of electrons on a ring

“The particle on a ring”

- On the ring, $V = 0$. Off the ring $V = \infty$.

$$\Psi = \sin(j\theta)$$

$$\begin{aligned}\hat{H}\Psi &= -\frac{\hbar^2}{2mr^2} \frac{\partial^2}{\partial \theta^2} \sin(j\theta) \\ &= \frac{\hbar^2 j^2}{2mr^2} \sin(j\theta) = \epsilon_j \Psi \quad j = 1, 2, 3, \dots\end{aligned}$$

“The particle on a ring”

- On the ring, $V = 0$. Off the ring $V = \infty$.

$$\Psi = \cos(j\theta)$$

$$\begin{aligned}\hat{H}\Psi &= -\frac{\hbar^2}{2mr^2} \frac{\partial^2}{\partial \theta^2} \cos(j\theta) \\ &= \frac{\hbar^2 j^2}{2mr^2} \cos(j\theta) = \epsilon_j \Psi \quad j = 0, 1, 2, 3, \dots\end{aligned}$$

Particle-on-a-ring

- Ground state is motionless




$$\Psi = \text{constant}$$

"The particle on a ring"


- The ring is a cyclic 1d potential

$$\Psi = \sin(j\theta) \quad \Psi = \cos(j\theta)$$



"The particle on a ring"



$$\epsilon_j = \frac{\hbar^2 j^2}{2mr^2} = \frac{2\pi^2 \hbar^2 j^2}{mL^2} \quad j = 0, 1, 2, 3, \dots$$



"The particle on a ring"


All singly degenerate

- n = 4
- n = 3
- n = 2
- n = 1



Doubly degenerate above j=0

- j = 3
- j = 2
- j = 1
- j = 0




ring



Application: benzene


Question: how many π -electrons in benzene?



Answer: Looking at the structure, there are 6 carbon atoms which each contribute one electron each. Therefore, there are 6 electrons.

benzene


Question: what is the length over which the π -electrons are delocalized, if the average bond length is 1.40 Å?




Answer: There are six bonds, which equates to $6 \times 1.40 \text{ \AA} = 8.40 \text{ \AA}$

benzene


Question: if the energy levels of the electrons are given by $\epsilon_j = 2\hbar^2 j^2 \pi^2 / mL^2$, what is the energy of the HOMO in eV?



Answer: since there are 6 π -electrons, and therefore the HOMO must have $j=1$. We know that $L = 6 \times 1.40 \text{ \AA} = 8.4 \text{ \AA}$. From these numbers, we get $\epsilon_j = 3.41 \times 10^{-19} j^2$ in Joules. The energy of the HOMO is thus $\epsilon_1 = 3.41 \times 10^{-19} \text{ J} = 2.13 \text{ eV}$.




benzene



Question: what is the energy of the LUMO, and thus the HOMO-LUMO transition?

Answer: $\epsilon_j = 3.41 \times 10^{-19} j^2$ in Joules. The energy of the LUMO is thus

$\epsilon_{j=2} = 1.365 \times 10^{-18} \text{ J} = 8.52 \text{ eV}$. The energy of the HOMO-LUMO transition is thus 6.39 eV.




benzene


Question: how does the calculated value of the HOMO-LUMO transition compare to experiment?

Answer: The calculated energy of the HOMO-LUMO transition is 6.39 eV. This corresponds to photons of wavelength

$\lambda = hc / (6.39 \times 1.602 \times 10^{-19}) \sim 194 \text{ nm}$, which is not so far from the experimental value (around 200 nm).



Hiraya and Shobatake, J. Chem. Phys. 94, 7700 (1991)



Learning Outcomes

- Be able to explain why confining a particle on a ring leads to quantization of its energy levels
- Be able to explain why the lowest energy of the particle on a ring is zero
- Be able to apply the particle on a ring approximation as a model for the electronic structure of a cyclic conjugated molecule (given equation for E_n).

Next lecture

- Quantitative molecular orbital theory for beginners

Week 10 tutorials

- Schrödinger equation and molecular orbitals for diatomic molecules

Practice Questions

1. The particle on a ring has an infinite number of energy levels (since $j = 0, 1, 2, 3, 4, 5 \dots$) whereas for a ring C_nH_n has only n p-orbitals and so n energy levels.

C_6H_6 , for example, only has levels with $j = 3$ (one level), $j = 1$ (two levels), $j = 2$ (two levels) and $j = 3$ (one level)

- (a) Using the analogy between the particle on a ring waves and the π -orbitals on slide 17, draw the four π molecular orbitals for C_4H_4 and the six π molecular orbitals for C_6H_6
- (b) Using *qualitative* arguments (based on the number of nodes and/or the number of in-phase or out-of-phase interactions between neighbours) construct energy level diagrams and label the orbitals as bonding, non-bonding or antibonding
- (c) Based on your answer to (b), why is C_6H_6 aromatic and C_4H_4 antiaromatic?
