

# Chemistry 2

## Lecture 3

### Particle on a ring approximation



## Assumed knowledge

Be able to predict the number of  $\pi$  electrons and the presence of conjugation in a ring containing carbon and/or heteroatoms such as nitrogen and oxygen

## Learning outcomes



- Be able to explain why confining a particle on a ring leads to quantization of its energy levels
- Be able to explain why the lowest energy of the particle on a ring is zero
- Be able to apply the particle on a ring approximation as a model for the electronic structure of a cyclic conjugated molecule (given equation for  $E_n$ ).

## The de Broglie Approach

- The wavelength of the wave associated with a particle is related to its momentum:

$$p = mv = h / \lambda$$

- For a particle with only kinetic energy:

$$E = \frac{1}{2} mv^2 = p^2 / 2m = h^2 / 2m\lambda^2$$

## Particle-on-a-ring


- Particle can be anywhere on ring




- Ground state is motionless

## Particle-on-a-ring


- Ground state is motionless
- In higher levels, we must fit an integer number of waves around the ring



1 wave  
 $\lambda = 2\pi r$



2 waves  
 $\lambda = 2\pi r/2$



3 waves  
 $\lambda = 2\pi r/3$



## Particle-on-a-ring

- $E = h^2 / 2m\lambda^2$
- $\lambda =$

## The Schrödinger equation

- The total energy is extracted by the Hamiltonian operator.
- These are the “observable” energy levels of a quantum particle

$$\hat{H}\Psi(x) = \epsilon_i\Psi(x)$$

Hamiltonian operator

Energy eigenvalue

Energy eigenfunction

## The Schrödinger equation

- The Hamiltonian has parts corresponding to *Kinetic Energy* and *Potential Energy*. In terms of the angle  $\theta$ :

$$\hat{H}\Psi = \left( -\frac{\hbar^2}{2mr^2} \frac{\partial^2}{\partial \theta^2} + V(\theta) \right) \Psi$$

Hamiltonian operator


Kinetic Energy

Potential Energy


## “The particle on a ring”

- The ring is a cyclic 1d potential

$$\Psi = \sin(j\theta) \quad \Psi = \cos(j\theta)$$



## “The particle on a ring”



$\pi$ -system of benzene is like a bunch of electrons on a ring

## “The particle on a ring”

- On the ring,  $V = 0$ . Off the ring  $V = \infty$ .

$$\Psi = \sin(j\theta)$$

$$\begin{aligned} \hat{H}\Psi &= -\frac{\hbar^2}{2mr^2} \frac{\partial^2}{\partial \theta^2} \sin(j\theta) \\ &= \frac{\hbar^2 j^2}{2mr^2} \sin(j\theta) = \varepsilon_j \Psi \quad j = 1, 2, 3, \dots \end{aligned}$$

## “The particle on a ring”


- On the ring,  $V = 0$ . Off the ring  $V = \infty$ .

$$\Psi = \cos(j\theta)$$

$$\begin{aligned} \hat{H}\Psi &= -\frac{\hbar^2}{2mr^2} \frac{\partial^2}{\partial \theta^2} \cos(j\theta) \\ &= \frac{\hbar^2 j^2}{2mr^2} \cos(j\theta) = \varepsilon_j \Psi \quad j = 0, 1, 2, 3, \dots \end{aligned}$$

## Particle-on-a-ring

- Ground state is motionless




$$\Psi = \text{constant}$$

## "The particle on a ring"

- The ring is a cyclic 1d potential

$$\Psi = \sin(j\theta) \quad \Psi = \cos(j\theta)$$




## "The particle on a ring"

$$\varepsilon_j = \frac{\hbar^2 j^2}{2mr^2} = \frac{2\pi^2 \hbar^2 j^2}{mL^2} \quad j = 0, 1, 2, 3, \dots$$


radius of ring

length of circumference






## "The particle on a ring"


All singly degenerate



Doubly degenerate above  $j=0$




## "The particle on a ring"



## Application: benzene

**Question:** how many pi-electrons in benzene?


**Answer:** Looking at the structure, there are 6 carbon atoms which each contribute one electron each. Therefore, there are 6 electrons.



## benzene

**Question:** what is the length over which the  $\pi$ -electrons are delocalized, if the average bond length is  $1.40\text{\AA}$ ?


**Answer:** There are six bonds, which equates to  $L = 6 \times 1.40\text{\AA} = 8.40\text{\AA}$



## benzene

**Question:** if the energy levels of the electrons are given by  $\epsilon_j = 2\hbar^2 j^2 \pi^2 / mL^2$ , what is the energy of the HOMO in eV?

**Answer:** since there are 6  $\pi$ -electrons, and therefore the HOMO must have  $j=1$ . We know that  $L = 6 \times 1.40\text{\AA} = 8.40\text{\AA}$ . From these numbers, we get  $\epsilon_j = 3.41 \times 10^{-19} j^2$  in Joules. The energy of the HOMO is thus  $\epsilon_1 = 3.41 \times 10^{-19} \text{J} = 2.13 \text{eV}$ .



\_\_\_\_\_  $j=3$



\_\_\_\_\_  $j=2$

↑↓  $j=1$   
↑↓  $j=0$

## benzene

**Question:** what is the energy of the LUMO, and thus the HOMO-LUMO transition?

**Answer:**  $\epsilon_j = 3.41 \times 10^{-19} j^2$  in Joules. The energy of the LUMO is thus  $\epsilon_2 = 1.365 \times 10^{-18} \text{J} = 8.52 \text{ eV}$ . The energy of the HOMO-LUMO transition is thus 6.39 eV.




## benzene


**Question:** how does the calculated value of the HOMO-LUMO transition compare to experiment?

**Answer:** The calculated energy of the HOMO-LUMO transition is 6.39 eV. This corresponds to photons of wavelength

$\lambda = hc / (6.39 \times 1.602 \times 10^{-19}) \sim 194 \text{ nm}$ , which is not so far from the experimental value (around 200nm).



Hiraya and Shobatake, J. Chem. Phys. 94, 7700 (1991)



## Summary

- The particle on a ring problem can be solved exactly and is a good first approximation for the electrons in a cyclically delocalized  $\pi$ -system.
- Confining a particle on a ring leads to quantization of its energy levels due to the condition that an integral number of waves must fit around the ring
- The lowest energy (ZPE) of a particle on a ring is zero
- The particle on a ring approximation can be applied as a model for the electronic structure of a cyclically conjugated molecule (given equation for  $E_n$ ).

## Next lecture

- Quantitative molecular orbital theory for beginners

## Week 10 tutorials

- Particle in a box approximation
  - you solve the Schrödinger equation.

## Practice Questions

1. The particle on a ring has an infinite number of energy levels (since  $j = 0, 1, 2, 3, 4, 5 \dots$ ) whereas for a ring  $C_nH_n$  has only  $n$  p-orbitals and so  $n$  energy levels.

$C_6H_6$ , for example, only has levels with  $j = 3$  (one level),  $j = 1$  (two levels),  $j = 2$  (two levels) and  $j = 3$  (one level)

- (a) Using the analogy between the particle on a ring waves and the  $\pi$ -orbitals on slide 17, draw the four  $\pi$  molecular orbitals for  $C_4H_4$  and the six  $\pi$  molecular orbitals for  $C_6H_6$
- (b) Using *qualitative* arguments (based on the number of nodes and/or the number of in-phase or out-of-phase interactions between neighbours) construct energy level diagrams and label the orbitals as bonding, non-bonding or antibonding
- (c) Based on your answer to (b), why is  $C_6H_6$  aromatic and  $C_4H_4$  antiaromatic?