


## CHEM1101 Worksheet 13: Concentration and Electrochemistry

### Model 1: The Effect of Concentration on the Cell Potential

In worksheet 12, you set up a **voltaic cell** to harness the electrical energy in a redox reaction, such as that below, to make a battery:



During the reaction, tin ions are consumed and zinc ions are produced. The driving force becomes weaker and the cell potential becomes smaller. Eventually, equilibrium is established and the battery is dead.

The **standard cell potential** refers to the reaction in which the reactants are present at **1 M concentrations**.

In Model 2 of worksheet 12, you calculated the standard cell potential for the oxidation of nicotine adenine dinucleotide (NADH) by  $\text{O}_2$ :



### Critical thinking questions

1. What pH does the standard cell potential refer to?
2. If the reaction is performed at a pH of 7.4, will the cell potential be higher or lower?

### Model 2: The Nernst Equation

The actual cell potential,  $E_{\text{cell}}$ , can be calculated from the standard cell potential using the Nernst equation:

$$E_{\text{cell}} = E_{\text{cell}}^{\circ} - \frac{RT}{nF} \ln Q$$

where  $R$  is the gas constant ( $8.314 \text{ J K}^{-1} \text{ mol}^{-1}$ ),  $T$  is the temperature (in Kelvin),  $n$  is the number of electrons transferred in the reaction,  $F$  is Faraday's constant ( $96485 \text{ C mol}^{-1}$ ) and  $Q$  is the reaction quotient. For the reactions in Model 1:

$$(1) \quad Q = \frac{[\text{Zn}^{2+}(\text{aq})]}{[\text{Sn}^{2+}(\text{aq})]}$$


$$(2) \quad Q = \frac{[\text{H}_2\text{O}][\text{NAD}^+]}{[\text{O}_2]^{1/2}[\text{H}^+][\text{NADH}]}$$

### Critical thinking questions

1. What is the value of  $n$  in these reactions? (*Hint*: how many electrons in total are required to change the oxidation number of Zn in (1) and of the two O atoms in  $\text{O}_2$  in (2)?)
2. In the biochemical literature, the biological standard state is used. This has all concentrations as 1 M, except  $[\text{H}^+]$  which is taken to be  $10^{-7}$  as this is closer to its value in the body. Use the Nernst equation to calculate the *biological* standard cell potential for the oxidation of NADH by  $\text{O}_2$  at the typical body temperature of  $37^\circ\text{C}$ .
3. Is your numerical answer to Q2 in Model 2 consistent with your answer to Q2 in Model 1?


### Model 3: Voltaic Cells

In worksheet 12, you saw how the electrical energy in a redox reaction can be harnessed to make a battery, by setting up the **voltaic cell** opposite. The potentials for the two reactions are:



The overall reaction is spontaneous as the reaction has a positive  $E_{\text{cell}}^{\circ}$  value:

$$E_{\text{cell}}^{\circ} = E_{\text{ox}}^{\circ} + E_{\text{red}}^{\circ} = +0.62 \text{ V}.$$



### Critical thinking questions


1. Which electrode (Zn or Sn) will *lose* mass and which one will *gain* mass?
2. What is the overall reaction that occurs when the cells are connected?
3. Oxidation always occurs at the anode. Label the anode and cathode on the cell.
4. Which way do the electrons flow? Draw an arrow on the diagram to show this.
5. In voltaic cells, electrons flow from the *negative* electrode to the *positive* electrode. Which is positive, the anode or the cathode? Label the electrodes as positive or negative.
6. The salt bridge contains  $\text{Na}^{+}(\text{aq})$  and  $\text{SO}_4^{2-}(\text{aq})$ . In which direction(s) do these ions move?

### Model 4: Electrolytic Cells

The *reverse* reaction can be made to happen if power from an external source with potential *greater* than  $E_{\text{cell}}^{\circ}$  is applied.

### Critical thinking questions


1. Which electrode (Zn or Sn) will now *lose* mass and which one will *gain* mass?
2. What is the overall reaction that now occurs when the cells are connected?
3. Oxidation always occurs at the anode. Label the anode and cathode on the cell.
4. Which way do the electrons flow? Draw an arrow on the diagram to show this.
5. The power source supplies electrons to the electrode where reduction occurs, so it becomes negative. The power source removes electrons from the electrode where oxidation occurs, so it becomes positive. Which is positive, the anode or the cathode? Label the electrodes as positive or negative.
6. The salt bridge contains  $\text{Na}^{+}(\text{aq})$  and  $\text{SO}_4^{2-}(\text{aq})$ . In which direction(s) do these ions move?



## Model 5: Electrolysis of Water

Electrolytic cells can be used to perform many useful tasks. A particular useful one is the electrolysis of water as this has the potential to convert electricity generated using solar energy into hydrogen gas, a combustible fuel.

The reactions at the cathode and anode are:



The amount of a substance produced in an electrolytic cell is directly proportional to the amount of electricity that passes through the cell. The number of moles of electrons that pass when a current  $I$  is applied for a time  $t$  is given by:

$$\text{number of moles of electrons} = I \times t / F$$

### Critical thinking questions

1. What is the overall reaction for the electrolysis of water?
2.  $F$  is Faraday's constant. It is the charge of one mole of electrons. The charge of one electron is  $1.602 \times 10^{-19}$  C. What is the charge of one mole?
3. If a current of 10.0 A is applied for 2.00 hours, how many moles of electrons are supplied? (*Hint*: remember to convert  $t$  into seconds).
4. How many moles of  $\text{H}_2\text{(g)}$  will be generated from this amount? (*Hint*: look at the stoichiometry of the reaction at the cathode.)
5. How many moles of  $\text{O}_2\text{(g)}$  will be generated from this amount?
6. Water is a poor conductor so a salt is usually added to increase the conductivity. The salt must contain ions that are harder to reduce or oxidise than water. Using the standard reduction potentials, select a suitable salt.

- How many hours does it take to form 10.0 L of  $O_2$  measured at 99.8 kPa and 28 °C from water if a current of 1.3 A passes through the electrolysis cell?

**Marks**  
**3**

Answer:

- What mass of  $PbSO_4$  is reduced at the cathode when a lead-acid storage battery is charged for 1.5 hours with a constant current of 10.0 A?

**Marks**  
**3**